

Is swimming safe in heart failure? A systematic review

Parin Shah, Pierpaolo Pellicori, Alexandra Macnamara, Alessia Urbinati, Andrew L. Clark

Introduction: Regular exercise is beneficial in patients with chronic heart failure; however, it is not clear whether swimming is safe.

Methods: We conducted a systematic review and identified 10 studies examining the haemodynamic effects of acute water immersion (WI) (155 patients; average age 60 years; 86% male; mean left ventricular ejection fraction (LVEF) 29%) and 6 randomised controlled trials of rehabilitation comparing swimming with either medical treatment only or cycling or aerobic exercise (136 patients, average age 59 years; 84% male, mean LVEF 31%).

Results: In 7 studies of warm WI (30 -35 °C): heart rate (HR) fell (2-15%), and both cardiac output (CO) (7-37%) and stroke volume (SV) increased (13-41%). In 1 study of hot WI (41°C), systemic vascular resistance (SVR) fell (41%) and HR increased (33%). In 2 studies of cold WI (12-22 °C), there were no consistent effects on HR and CO.

Compared with:

1. Medical management (n=3 studies): swimming led to a greater increase in peak VO_2 (7-14%) and 6 minute walk test (6 MWT) (7-13%) with no consistent effects on quality of life.
2. Cycle training only (n=1 studies): combined swimming and cycle training led to greater reduction in resting HR (16%), greater increase in resting SV (23%) and SVR (15%), but no changes in resting CO and a lesser increase in peak VO_2 (6%).
3. Aerobic training only (n=2 studies): combined swimming and aerobic training lead to a reduction in resting HR (19%) and SVR (54%) and a greater increase in SV (34%), resting CO (28%), LVEF (9%) and 6MWT (70%).

Conclusions: Although swimming appears to be safe, the studies conducted have been small, very heterogeneous and inconclusive.

Key words: heart failure, water immersion, swimming

Introduction:

All current guidelines recommend regular exercise as an integral part of the management of patients with chronic heart failure (CHF).^{1,2,3,4} The sorts of exercise that have been shown to be helpful include: aerobic exercise on a treadmill or a cycle at 3-5 times per week at a heart rate of 60-70% of heart rate reserve; and resistance training of peripheral muscles and calisthenics.^{5,6,7} In two meta-analyses, exercise training was associated with a reduction in mortality and hospital admissions as well as improvements in maximum oxygen consumption (peak VO_2), walking distance and quality of life compared to standard medical care.^{6,7} A subsequent randomised trial of more than 2000 patients randomised to supervised exercise training followed by unsupervised exercise training at home (3 sessions per week for 30-35 mins at 70% of heart rate reserve, total of 36 sessions) or usual care. Exercise was associated with an absolute risk reduction of 4% in death or hospitalization from any cause at 3 years.⁵

However, whether swimming is either safe or beneficial is not clear. Most guidelines steer clear of discussing swimming. Only the Scottish guidelines mention swimming in order to warn against it in patients with NYHA class III and IV symptoms.³

Potentially, the pressure exerted by water immersion (WI) can result in the shift of extracellular fluid and peripheral blood back into the venous circulation. This would in turn increase venous return to the failing heart and precipitate pulmonary oedema in susceptible patients.

Patients with many different conditions do swim regularly, especially patients with musculoskeletal conditions.⁸ Whether it is an appropriate treatment for people with

heart failure is not clear: we therefore conducted a systematic review on the effects of water immersion and swimming in patients with chronic heart failure.

Methods:

Search strategy and study selection

We systematically searched for publications on swimming and heart failure on PubMed until January 2016. The words “heart failure”, “ventricular dysfunction” or “cardiomyopathy” in combination with “swimming”, “water immersion”, “hydrotherapy”, “aquatic exercise”, “water exercise” and “water based exercise therapy” were used as search criteria. Randomised controlled trials, controlled trials, and single arm studies were included.

Outcome measures

Haemodynamic variables measured during water immersion to the neck and swimming rehabilitation were: heart rate (HR), SV, CO, systemic vascular resistance (SVR), peak VO_2 , left ventricular end diastolic diameter (LVEDD), left ventricular end systolic diameter (LVESD) and left ventricular ejection fraction (LVEF). A further two outcomes reported for swimming rehabilitation were: quality of life (QOL) and six minute walk test distance (6 MWT).

Results:

We identified 41 publications in English. Studies on animals (n=10), in conditions other than heart failure (n=4), reviews, case reports or letters to editors (n=12) were excluded. Fifteen eligible publications were thus considered (Figure 1).

Acute haemodynamic effects of water immersion:

There were 10 studies examining the acute haemodynamic effects of water immersion with a total of 155 patients (average age 60 years; 86% male; mean LVEF 29%).^{9,10,11,12,13,14,15,16,17,18} Seven were in warm water (32-35°C) with a total of 97 patients (average age of 61 years; 85% male; mean LVEF 30%). Four of these seven compared the responses of patients with normal subjects and the remaining 3 were in patients alone.^{9,10,11,12,13,14,15} Two were in cold water (12-22°C) with a total of 24 patients (average age of 60 years; mean LVEF 31%).^{17,18} One study compared the responses of patients with normal subjects and one was in patients alone. There was a single study of hot water immersion (41°C) with 34 patients (average age 58 years; 76% male; mean LVEF 25%).¹⁶ (Figure 2)

Warm water immersion:

In the 7 studies which compared haemodynamics at baseline to warm WI: 6 studies measured HR, which reduced by a mean of 7% (range: 2 to - 15%).^{10,11,12,13,14,15} Five studies measured SV, which increased in most patients by a mean of 32% (range: 13 to 41%),^{9,11,13,14,15} apart from in a sub-population of patients with severe heart failure in one study in whom SV decreased by 4%.⁹ Five studies measured CO, which

increased by a mean of 22% (range 7 to 37%).^{11,12,13,14,15} Three studies measured SVR, which decreased by a mean of 9% (range 3 to 21%).^{11,14,15} (Figure 3)

In 3 studies which compared echocardiographic changes at baseline to warm WI: 2 studies measured left ventricular end diastolic volume, which increased by a mean of 16% (range 6 to 24%) and LVEF, which increased by a mean of 13% (range 12 to 13%).^{11,13} One study showed a greater increase in left ventricular end diastolic diameter (LVEDD) in patients with moderate heart failure (19%) compared to severe heart failure (5%).⁹ (Figure 3)

4 studies examined the change in haemodynamics from baseline during warm WI in patients with CHF compared with healthy subjects: the changes were of similar magnitude in both groups. In 4 studies, HR decreased in both groups by a mean of 7% (range 2 to -15%) in CHF and by a mean of 11% (range -1 to -22) in healthy subjects.^{10,12,13,14} In 2 studies, SV increased in both groups by a mean of 28% (range 17 to 37%) in CHF and by a mean of 50% (range 30 to 65%) in healthy subjects.^{13,14} In 3 studies, CO increased in both groups by a mean of 20% (range 14 to 37%) in CHF and by a mean of 26% (range 19 to 31%) in healthy subjects.^{12,13,14} In 1 study, SVR decreased in both groups by a mean of 21% in CHF and by a mean of 28% in healthy individuals.¹⁴ (Figure 3)

Hot water immersion:

In the single study of hot water immersion, HR increased (33%), SV increased (16%), CO increased (50%) and SVR decreased (41%).¹⁶ On echocardiography, LVEDV

(2.4%) and left ventricular end systolic diameter (5.4%) both decreased, but LVEF increased (23%).¹⁶ (Figure 3)

Cold water immersion:

In the 2 studies which compared haemodynamics at baseline to cold WI: both studies measured CO which increased, both measured heart rate with inconsistent results and one study measured SVR, which fell.^{17,18} (Figure 3)

Swimming as a form of rehabilitation in patients with heart failure:

There were 6 trials of exercise training with patients. Three studies randomised patients to either swimming or continued medical therapy (67 patients, mean age of 70, 73% male, mean LVEF of 32%).^{11,19,20} Two studies randomised patients to either a combination of swimming and aerobic exercise or aerobic training only (45 patients, mean age of 60, mean LVEF of 31%).^{21,22} One study randomised patients to either gymnastics on land or in water in combination with cycle training in both groups (24 patients, mean age of 54, mean LVEF 30%).²³ (Figure 4)

Swimming training compared to medical management only:

Three studies compared swimming training with medical management only.^{11,19,20} At the end of follow up (8 weeks): in one study, there were no changes in HR, SV, CO or SVR at rest.¹¹ In 2 studies,^{19,20} there was an increase in peak VO₂ by a mean 10% (range: 7 to 14%) in swimming training vs. a reduction by a mean -10% (range -12 to -

7% in medical management) whilst there was no change in peak VO_2 in the third study.¹¹ In 2 studies,^{19,20} there was a greater increase in 6 MWT with swimming, by a mean of 10% (range: 7 to 13%) in swimming training vs. increase by a mean of 1% (range 0 to 2%) in medical management. There was no consistent pattern in quality of life in the 2 studies where it was measured.^{19,20} (Figure 4)

Cycle training compared to a combination of swimming and cycling training:*Error! Bookmark not defined.*

In a study which compared haemodynamics and peak VO_2 between cycle training only or a combination of swimming and cycle training: there was a greater reduction in resting HR (-16% swimming and cycling vs. -8% cycling only), greater increase in resting SV (23% swimming and cycling vs. 10% cycling only), no change in CO at rest (0% swimming and cycling vs. 7% cycling only), an increase in resting SVR (15% swimming and cycling vs. -15% cycling only) and an increased peak VO_2 (6% swimming and cycling vs. 11% cycling).²³ (Figure 4)

Aerobic training compared to a combination of swimming and aerobic training:

In 2 studies which compared haemodynamics between aerobic exercise training only or a combination of swimming and aerobic exercise, at the end of follow up (range 3-24 weeks): there was a greater reduction HR (-19% swimming and aerobic exercise vs. +5% aerobic exercise only) in one study²² but no difference in the other;²¹ greater increase in SV (34% swimming and aerobic exercise vs. 15% aerobic exercise only) in one study²² but no difference in the other.²¹ In one study there was a greater

increased CO at rest (28% swimming and aerobic exercise vs. 9% aerobic exercise only) and greater reduction SVR (54% swimming and aerobic exercise vs. 19% aerobic exercise only).²² (Figure 4)

The echocardiographic findings were: LVEDD reduced (3% swimming and aerobic exercise vs. 5% aerobic exercise only) in one study²² but no difference in the second;²¹ LVESD reduced (9% swimming and aerobic exercise vs. 2% aerobic exercise only) in one study²² no change in the second;²¹ with a similar increase in resting LVEF in both studies (9% swimming and aerobic exercise vs. 6% aerobic exercise). (Figure 4)

One study reported six minute walk test distance, and found that there was a greater increase following swimming and aerobic exercise (70%) than following aerobic exercise alone (32%).²² (Figure 4)

Discussion:

We have found that water immersion to the neck is well tolerated in stable patients with chronic heart failure, particularly when the water is warm. Both cold and hot water immersion caused adverse haemodynamic effects, whereas warm water immersion appeared helpful. The most consistent effects were a fall in resting heart rate and systemic vascular resistance. Exercise in water has effects similar to other forms of exercise training with a similar improvement in exercise capacity. Exercise in water in stable patients with NYHA class II-III symptoms was well tolerated with no adverse effects.

The mechanical properties of water are considerably different to air; there are several changes in haemodynamics and fluid shift even when healthy subjects are immersed in water. During WI to the neck, approximately 700 ml of peripheral blood pools centrally (of which 180-240 ml accumulates in the heart) due to the hydrostatic pressure exerted on the body.^{24,25} As a consequence, in healthy subjects, there is an increase in left ventricular volume and an increase in left ventricular ejection fraction (LVEF) resulting in increased stroke volume and blood pressure.^{26,27} In healthy subjects, cycling in water leads to a greater increase in cardiac output (CO) than cycling in air by an average of 0.7 l/min.²⁸ Stroke volume (SV) increases by 49% in air compared to 34% in water.²⁶

Guidelines are cautious about water immersion and swimming in patients with heart failure because of the potential to precipitate pulmonary oedema secondary to the mobilisation of peripheral fluid into the central circulation. Reports of swimming-induced pulmonary oedema (SIPE) are rare, and confined to athletes usually undertaking severe exercise. The pathophysiology of SIPE is not well understood but factors such as the central shift of fluid during water immersion, vasoconstriction due to cold water and over hydration may increase right-sided cardiac pressures leading to pulmonary oedema.^{29,30}

In a study by Weiler-Ravell and colleagues on 30 young men on a military fitness programme, swimming in water at 23°C caused pronounced shortness of breath in 8 subjects.³⁰ Five subjects stopped swimming early. They needed oxygen supplementation and in the more severe cases intravenous diuretics.³⁰ Subjects with a history of SIPE have higher pulmonary artery pressure (previous SIPE = 34 mmHg vs control = 22 mmHg, $p=0.004$) and pulmonary artery wedge pressure (previous SIPE =

19 mmHg vs control = 11 mmHg, $p=0.028$) when exercised in cold water on a cycle than those who do not.²⁹

None of the studies of water immersion or swimming rehabilitation in patients with heart failure reported SIPE. However, all the studies we found were small, involving at most 34 patients. The majority of patients were male and were young compared to the general population of patients with heart failure. Most of the studies which looked at swimming rehabilitation did not prescribe swimming on its own but in conjunction with aerobic exercise on land and so it is difficult to assess the effects of swimming training alone on cardiac function and exercise performance.

What is perhaps surprising is how few patients have been included in studies of swimming. In the United Kingdom, swimming is the most popular sport with over 3 million people taking part in at least a session per week. This is a million more people per week than play football (which is often considered to be the UK's national sport).³¹ Swimming may be more popular than other forms of exercise because it is a low impact activity suitable for people with disability, poor mobility or frailty.**Error! Bookmark not defined.** Given how common swimming is, it must be the case that patients with heart failure are swimming regularly. The data we have found (and perhaps the lack of reports in the literature) suggest that swimming is probably safe in patients with heart failure. Given that patients with heart failure are often frail and have other co-morbidities (which reduce mobility), swimming is perhaps the ideal way to encourage exercise without the difficulty of weight-bearing exercise on land. However, much larger studies are needed to assess the safety and potential benefits of swimming compared to conventional heart failure rehabilitation.

Conclusion:

Although exercise in water appears to be safe, the studies conducted have been small, very heterogeneous and inconclusive. There is a crying need for further work to establish the effects of immersion and swimming so as to be able give accurate advice to patients with heart failure.

References

- ¹ National Institute for Health and Clinical Excellence (2010) Chronic heart failure: Management of chronic heart failure in adults in primary and secondary care. Clinical Guideline 108. London: NICE
- ² 2013 ACCF/AHA Guideline for the Management of Heart Failure A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2013;**128**:240-327
- ³ Scottish Intercollegiate Guidelines Network. Management of chronic heart failure. Edinburgh: SIGN; 2007. (SIGN Guideline no. 95)
- ⁴ McMurray JJV, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Køber L, Lip GYH, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Rønnevik PK, Rutten FH, Schwitter J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeiher A. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. *European Heart Journal* 2012; **33**:1787-1847.
- ⁵ O'Connor CM, Whellan DJ, Lee KL, Keteyian SJ, Cooper LS, Ellis SJ, Leifer ES, Kraus WE, Kitzman DW, Blumenthal JA, Rendall DS, Miller NH, Fleg JL, Schulman KA, McKelvie RS, Zannad F, Pinˆa IL; HF-ACTION Investigators. Efficacy and safety of exercise training in patients with chronic heart failure: HFACTION randomized controlled trial. *JAMA*. 2009;**301**:1439–1450.
- ⁶ Rees K, Taylor RS, Singh S, Coats AJS, Ebrahim S. Exercise based rehabilitation for heart failure. *Cochrane Database of Systematic Reviews*. 2004, Issue 2
- ⁷ Piepoli MF. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ*. 2004;**328**:189
- ⁸ Chase NL, Sui X, Blair SN. Comparison of the Health Aspects of Swimming With Other Types of Physical Activity and Sedentary Lifestyle Habits. *International Journal of Aquatic Research and Education*. 2008;**2**:151-161
- ⁹ Meyer K and Bucking J. Exercise in Heart Failure: Should Aqua Therapy and Swimming Be Allowed? *Medicine & Science in Sports & Exercise*. 2004;**36**: 2017-23
- ¹⁰ Cider A, Sunnerhagen KS, Schaufelberger M Andersson B. Cardiorespiratory effects of warm water immersion in elderly patients with chronic heart failure. *Clin Physiol Funct Imaging*. 2005;**25**:313–317

-
- ¹¹ Sveälv BG, Cider A, Täng MS, Angwald E, Kardassis D, Andersson B. Benefit of warm water immersion on biventricular function in patients with chronic heart failure. *Cardiovasc Ultrasound*. 2009;**7**:33
- ¹² Gabrielsen A, Sorensen VB, Pump B, Galatius S, Videbaek R, Bie P, Warberg J, Christensen NJ, Wroblewski H, Kastrup J, Norsk P: Cardiovascular and neuroendocrine responses to water immersion in compensated heart failure. *Am J Physiol Heart Circ Physiol*. 2000; **279**:1931-1940.
- ¹³ Cider A, Sveälv BG, Tang MS, Schaufelberger M, Andersson B. Immersion in warm water induces improvement in cardiac function in patients with chronic heart failure. *Eur J Heart Fail* 2006;**8**:308–313
- ¹⁴ Schmid JP, Noveanu M, Morger C, Gaillet R, Capoferri M, Anderegg M, Saner H. Influence of water immersion, water gymnastics and swimming on cardiac output in patients with heart failure. *Heart* 2007;**93**:722–727
- ¹⁵ Mourot L, Teffaha D, Bouhaddi M, Ounissi F, Vernochet P, Dugue B, REgnard J, Monpere C. Exercise rehabilitation restores physiological cardiovascular responses to short-term head-out water immersion in patients with chronic heart failure. *J Cardiopulm Rehabil Prev*. 2010;**30**:22-7
- ¹⁶ Tei C, Horikiri Y, Park JC, Jeong JW, Chang KS, Toyama Y, Tanaka N. Acute Hemodynamic Improvement by Thermal Vasodilation in Congestive Heart Failure. *Circulation* 1995; **91**: 2582-2590
- ¹⁷ Radtke T, Poerschke D, Wilhelm M, Trachsel LD, Tschanz H, Matter F, Jauslin D, Saner H, Schmid JP. Acute effects of Finnish sauna and cold-water immersion on haemodynamic variables and autonomic nervous system activity in patients with heart failure. *Eur J Prev Cardiol*. 2016;**23**:593-601
- ¹⁸ Schmid JP, Morger C, Noveanu M, Binder RK, Anderegg M, and Saner H. Haemodynamic and arrhythmic effects of moderately cold (22⁰C) water immersion and swimming in patients with stable coronary artery disease and heart failure. *European Journal of Heart Failure*. 2009;**11**:903–909
- ¹⁹ Cider A, Schaufelberger M, Sunnerhagen KS, Andersson B. Hydrotherapy—a new approach to improve function in the older patient with chronic heart failure. *The European Journal of Heart Failure*. 2003; **5**: 527–535
- ²⁰ Cider A, Schaufelberger M, Stibrant SK, Andersson B. Aquatic Exercise Is Effective in Improving Exercise Performance in Patients with Heart Failure and Type 2 Diabetes Mellitus. *Evidence-Based Complementary and Alternative Medicine*. 2012; **2012**

-
- ²¹ Teffaha D, Mourot L, Vernochet P, Ounissi F, Regnard J, Monpère C, Dugu B. Relevance of Water Gymnastics in Rehabilitation Programs in Patients With Chronic Heart Failure or Coronary Artery Disease With Normal Left Ventricular Function. *J Cardiac Fail.* 2011;**17**:676-683
- ²² Caminiti G, Volterrani M, Marazzi G, Cerrito A, Massaro R, Sposato B, Arisi A, Rosano G. Hydrotherapy added to endurance training versus endurance training alone in elderly patients with chronic heart failure: A randomized pilot study. *International Journal of Cardiology.* 2011;**148** :199–203
- ²³ Laurent M, Daline T, Malika B, Fawzi O, Philippe V, Benoit D, Catherine M, Jacques R. Training-induced increase in nitric oxide metabolites in chronic heart failure and coronary artery disease: an extra benefit of water-based exercises? *Eur J Cardiovasc Prev Rehabil.* 2009;**16**:(2) 215-21
- ²⁴ Lehmann M and Same L. Recreational Swimming in CHD Patients and Healthy Control Subjects in Relation to Left Heart Function. *Clinical Cardiology* 1990; **13**: 547-554.
- ²⁵ Risch WD, Koubenec HJ, Beckmann U, Lange S, Gauer OH. The effect of graded immersion on heart volume, central venous pressure, pulmonary blood distribution, and heart rate in man. *Pflugers Arch.* 1978;**374**:115-8.
- ²⁶ Smith DE, Kaye AD, Mubarek SK, Kusnisk BA, Anwahr M, Friedman IM, Nossman BD. Cardiac effects of water immersion in health volunteers. *Echocardiography.* 1998;**15**:35-428.
- ²⁷ K Meyer. Left ventricular dysfunction and chronic heart failure: should aqua therapy and swimming be allowed? *Br J Sports Med.* 2006;**40**:817-8
- ²⁸ Sheldahl LM, Tristani FE, Clifford PS, Hughes CV, Sobocinski KA, Moris RD. Effect of Head-Out Water Immersion on Cardiorespiratory Response to Dynamic Exercise. *J Am Coll Cardiol.* 1987;**10**:1254-8
- ²⁹ Moon RE, Martina SD, Peacher DF, Potter JF, Wester TE, Cherry AD, Natoli MJ, Otteni CE, Kernagis DN, White WD, Freiburger JJ. Swimming-Induced Pulmonary Edema: Pathophysiology and Risk Reduction With Sildenafil. *Circulation.* 2016;**133**:988-96
- ³⁰ Weiler-Ravell D1, Shupak A, Goldenberg I, Halpern P, Shoshani O, Hirschhorn G, Margulis A. Pulmonary oedema and haemoptysis induced by strenuous swimming. *BMJ.* 1995;**311**:361-2.
- ³¹ Evaluation of the Impact of Free Swimming. Year 1 report – main report. Department for Culture, Media & Sport. June 2010.

<https://www.gov.uk/government/publications/free-swimming-programme-2009-10-annual-report>




Figure 1: systematic review results




Figure 2: breakdown of eligible studies

Percentage change in haemodynamic, respiratory and echocardiographic findings from rest to water immersion




Figure 3: Percentage change in haemodynamic, respiratory and echocardiographic findings from rest to water immersion