

1 **MIGRATION FLYWAY OF THE MEDITERRANEAN BREEDING LESSER**
2 **CRESTED TERN *Thalasseus bengalensis emigratus***

3 Abdulmaula Hamza^{1,2*}, Nicola Baccetti³, Joe Sultana⁴, Jaber Yahia⁵, Marco Zantello³,
4 Adriano de Faveri³, Nick Cutts², John Borg⁴, Hichem Azafzaf⁶, Pierre Defos du Rau⁷ Esam
5 Bourass⁸, Khaled Etayeb⁹ and Michael Elliott²

6 1 School of Marine and Environmental Sciences, University Malaysia Terengganu 21030,
7 Kuala Terengganu, Malaysia.

8 2 Institute of Estuarine and Coastal Studies (IECS), University of Hull, Hull, HU6 7RX,
9 UK

10 3 Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), via Ca'
11 Fornacetta 9, Ozzano Emilia (Bologna), 40064, Italy.

12 4 Birdlife Malta. 57/28 Triq Abate Rigord, Ta' Xbiex, MT, XBX1120, Malta.

13 5 Zoology Department, Faculty of Science, Omar Almokhtar University, Bayda, Libya.

14 6 Regional Activity Centre for Specially Protected Areas. Tunis, Tunisia.

15 7 Office National de la Chasse et de la Faune Sauvage, Unité Avifaune Migratrice, Le
16 Sambuc, 13200 Arles, France

17 8 Nature Conservation Department. Environment General Authority, Tripoli, Libya

18 9 Zoology Department, Faculty of Sciences, University of Tripoli, Tripoli, Libya

19 *Corresponding author: a.hamza@umt.edu.my Tel. +6096683450

20

21 **Abstract:**

22 The Lesser Crested Tern (*Thalasseus bengalensis emigratus*) breeding population in the
23 Mediterranean is found exclusively in Libya, on the two coastal islands of Gara and Elba and
24 one wetland on the mainland coast at Benghazi. In order to improve knowledge of the species
25 migration to wintering quarters in West Africa, a ringing programme was initiated in 2006 and
26 continued until 2012. From a total of 1354 nestlings ringed using metal and/or colour rings, 64
27 were recovered along their flyway and in their wintering range, representing 6.90% of birds
28 ringed with both colour and metal rings. This provided the opportunity to collect information on
29 post-natal movements (staging and wintering ranges), breeding philopatry and recruitment, in
30 addition to a preliminary estimation of their migration journey duration. This paper indicates

1 sighting and recovery distributions in space and time, highlighting the important areas for the
2 species during its journey between breeding and wintering sites. The findings indicate that
3 several areas where ringed terns stop-over during pre- and post-breeding migration journeys are
4 not protected, causing an additional threat to their survival, as some wintering areas are also not
5 protected. Conservation of this highly localised and threatened population needs not only to
6 address protection at breeding sites but also at migratory stop-overs and wintering strongholds.

7 **Keywords:** Lesser Crested Tern, ringing, migration, breeding, Libya, West Africa.

8

9

10 **Introduction**

11 Mark and recapture techniques, with accurately recorded information within a ringing scheme,
12 are a relatively common and low-cost method to understand population movement and dispersal
13 ecology (Spina, 1999). Bird ringing has provided valuable information, not only through ring
14 sightings of live birds and recoveries from dead ones, but also by determining the migration
15 routes and patterns of several species (Wernham et al., 2002). Bird ringing can also contribute to
16 the monitoring of the population dynamics of a species (Balmer et al., 2008). It provides
17 measures of survival, productivity and dispersal (Viana, 2013; Baillie, 2001) and it is a useful
18 tool for behavioural studies (see review by Sharp, 2009). Ringing is also a support tool for
19 conservation actions and adaptive management (Nichols, 2007). The probability of encountering
20 recoveries decreases when wintering areas are distant from breeding grounds (Shiomi, 2015),
21 therefore long-term ringing campaigns and intensive ring monitoring at targeted areas are central
22 to successful migration studies in such cases.

23 While avian bio-geographical data for most North African countries are available, Libya in
24 particular remains one of the least covered countries in terms of bird studies (Bundy, 1976;
25 Etayeb, 2007; Smart et al., 2006; EGA/RAC/SPA Team, 2012). The Lesser Crested Tern
26 *Thalasseus bengalensis emigratus* is the smallest in population size compared to the other two
27 subspecies *T. b. bengalensis* (Red Sea and East Africa) and *T.b. torresi* (Persian Gulf to southeast
28 Asia and Australia). It is an endemic exclusive summer breeder to Libya and winters in West
29 Africa (Hamza et al., 2012, Hamza, 2014). This localised small-sized population was classified
30 as Endangered under the Mediterranean Action Plan on Seabirds (RAC/SPA, 2003), the action
31 plan calls to conduct studies on ecology and movements of this subspecies being of high
32 conservation relevance.

1 The ringing of young birds at the breeding sites has been conducted during 2006-2008 (Azafaf
2 et al., 2006, Hamza et al., 2007) followed by a second campaign in 2009-2012 (Hamza, 2014).
3 The present study aims to: (a) improve our understanding of post-natal dispersal using ring
4 sightings and field observations; (b) investigate initial recruitment age at breeding colonies, and
5 (c) identify the important staging and wintering sites based on sightings in North and West
6 Africa.

7 **Materials and Methods:**

8 **Study sites**

9 Two coastal islands of Gara and Elba and one wetland on the Benghazi coast are the main
10 breeding sites for this tern population (Fig. 1). Gara Island, is on the eastern side of the Gulf of
11 Sirte at 30°48'N 19°54'E, approx.12 km off the coast near Ajdabiyah and has a total area of 4.5
12 ha and it is 7 m above sea level. Elba island is a small, low-lying Island situated in the Gulf of
13 Bumbah, 32° 14'N 23° 17'E (20 ha). The third breeding site is located inshore at the northern
14 section of Sebkheth Jeliana, Benghazi (a 30 ha. permanently flooded salt-marsh), on the islet of
15 Jeliana (position 32°05'N 20°03'E and 35m²).

16

17 **Results**

18 **Ringling totals and recoveries**

19 A total of 1352 young terns were ringed at their colonies between 2006 and 2012 breeding
20 seasons. Most of the ringing effort occurred at Gara island (72.85%) being the largest colony
21 with a population size of more than 2,000 breeding pairs, 19.9% and 7.25% of rings were used
22 at the Jeliana and Elba colonies respectively (Table. 1). Ringed birds represented 20-60% of
23 crèche size present in each season at both Gara and Jeliana colonies, while at Elba all young birds
24 were ringed. A total of 64 ringed terns have been re-observed/recovered (Table 2) representing
25 4.73% of total ringed birds with metal and colour rings (n=1352) and 7.47 % of the colour ringed
26 Lesser Crested Terns (n= 857) in the present study (Table 2).

27

28 **Post-natal movements (staging areas)**

29 The present study confirmed that fledging occur during late July for Jeliana and during late-
30 August to early September at Elba and Gara. Juveniles continue to depend on their parents for
31 feeding during their first months. Nine recoveries were juveniles accompanying their parents, 6

1 from Gara and 3 from the Jeliana colony. For example, one bird ringed at Gara in 2008 has been
2 sighted in Ceuta, Spain after one month and 27 days of its ringing date. Another two chicks from
3 Gara ringed in were re-sighted at Almadraba beach, Spain within 47 days and 58 days after being
4 ringed in Libya. They roost on small islets, and in estuaries and coastal wetlands during migration
5 to and from wintering areas in West Africa between Senegal, Gambia, Guinea-Bissau and Sierra
6 Leone. During this study, one bird was reported by a fisherman in Tripoli harbour after 144 days
7 since its ringing date in August 4th at Gara Island (Figure 2).

8 The first part of the migration was monitored at several sites along the Libyan coastline. For
9 example, a small group of 3-5 birds was seen in late August 2009 for two days at Al-Ghbeba
10 beach, west of Sirte. Another 18 birds were seen earlier on August 15th 2009 roosting with Little
11 Terns *Sterna albifrons*, and feeding their young on small rocky islets off the coast of Sabratah
12 town (80 km west of Tripoli). The post-natal dispersal is unsynchronised, with some birds leaving
13 the breeding sites much earlier than others.

14 **Post-natal movements (wintering range)**

15 Within-Libya wintering records ranged from three to fifteen individuals, reported during the mid-
16 winter waterbird census between 2007 and 2011. The most distant dispersal recorded for a newly
17 fledged Lesser Crested was reported from the Turtle Islands, Sierra Leone, having travelled some
18 6,700 km within 149 days after being ringed in Libya (average 45 km/day). Six of the 46 sighted
19 birds were observed in Dakar (2), N'Gor Island in Senegal (2) and two at the Tanji Bird Reserve
20 in Gambia.

21 Some birds were sighted more than once after being ringed at their natal sites, for instance an
22 individual sighted in October 2013 after three years and two months of ringing at Gara in Melilla,
23 Southern Spain, then again in January 2014 at Catabao Segundo, Guinea Bissau. The same bird
24 was re-sighted in October 2014, exactly a year later, showing the same pattern in stopping at the
25 same site each year on the way from breeding to wintering areas.

26 Another individual was ringed initially in 2007 at Elba Island, then was first re-sighted after
27 almost 5 years (July 2012) in a different breeding site of Jeliana in Libya, and then it was re-
28 sighted in Massa estuary, Morocco in April 2013.

29 These records confirm the regular migratory pattern of Lesser Crested Tern along the North-
30 West African coastline between breeding and wintering zones.

31

1 **Site fidelity and recruitment**

2 During the 2011 season, one ringed Lesser Crested Tern was observed in flight and bred at Elba
3 Island where it had been ringed in August 2007. In 2012 three additional ringed terns were
4 observed breeding at Elba. One adult bird had a blue ring (ringed in 2009), while the other two
5 had white rings (ringed in 2010). These latter two were the youngest birds to return to a breeding
6 site during this study. At Jeliana, eleven ringed adults were breeding in the 2012 season, eight of
7 which were ringed at the same site in August 2008, one ringed in Jeliana in the 2009 season, the
8 other two were a ringed bird from the Gara Island (ringed in August 2008) and a ringed bird from
9 Elba in August 2007; these latter two records clearly demonstrate inter-colony movements of
10 this species.

11 **Discussion**

12 Ring sightings and recoveries of Lesser Crested Terns in the present study showed that several
13 sites along the North and Northwest African coast are used as staging sites during the migration
14 to and from the breeding sites. Some juveniles spent their first winter with their parents, at some
15 sites along the north and northwest African coast in what is known as ‘a nomadism period’ during
16 the first year (Barlow, 1998) without wintering in West Africa as do most post-breeding migrant
17 terns. In contrast, West Africa is the final wintering destination for this population, as there are
18 no data available on *T. b. emigratus* dispersal to the Eastern Mediterranean basin or the Red Sea,
19 despite some limited observations on the Egyptian Mediterranean coastline (Goodman and
20 Meininger, 1989). The latter could have been foraging or non-breeding individuals from the
21 nominate subspecies *T. b. bengalensis* that breeds in the Red Sea.

22 Some important staging sites were identified by the present study, the Strait of Gibraltar (5
23 sightings within 1-2 months after ringing) is a must-pass area for this species and many other
24 migratory bird species (Hashmi, 2000, Paracuellos and Jerez, 2003) and consequently the
25 Moroccan Mediterranean and Atlantic coastal areas are similarly important migration hubs.

26 Lesser Crested Terns are generally rare in the Iberian Peninsula, however, few individuals are
27 detected around Cadiz, the Doñana National Park, Montijo and La Jara beaches (at the mouth of
28 the Guadalquivir River), Salinas de la Tapa (El Puerto de Santa María) and Odiel marshes
29 (Huelva), associating with Sandwich Tern *Thalasseus sandvicensis* and Common Tern *Sterna*
30 *hirundo* (C. Gutiérrez, SEO/Birdlife, pers. com). During this study four ringed birds were
31 reported from there.

1 The present study has also identified areas that still require more monitoring efforts, particularly
2 northern Tunisia and the Algerian coast during onward and outward migrations (Isenmann et al.,
3 2005). The species is passage migrant in Mauritania, where the Banc d'Arguin National Park is
4 an important hotspot (Isenmann et al., 2010). The peninsula of Nouadhibou to the north of the
5 country represents the northern limit of the wintering range. Similarly, Senegal hosts large
6 concentrations of Lesser Crested Terns. Over 825 birds were counted from 3 to 16 October 2005
7 at N'Gor, with a daily average of 59 birds (Holmström et al., 2005). The Tanji Bird Reserve and
8 Barra Ferry Terminal on the Gambia River are also sites where the species has been frequently
9 reported by several birdwatchers (Skov and Jensen, 2002). Senegal and Gambia are in fact at
10 the centre of the wintering range for the species and more monitoring is essential to understand
11 where the species is concentrated and what conservation measures need to be in place (Schricke
12 et al., 2001).

13 Guinea-Bissau hosts an important numbers of wintering Lesser Crested Tern, a total of 400
14 individuals were counted along the coast and the total number of wintering individuals was
15 estimated at 600-1000 (Meininger, 1988). However, no updated information is available on the
16 species in this country.

17 More monitoring is necessary determine the wintering southern distribution of the species. In the
18 Sierra Leone Western Area National Park, several Lesser Crested Terns were recorded in
19 December 2008, associated with other tern species, such as the Royal Tern
20 *T. maximus* (Valentine, 2008). In the present study one ring recovery was from Turtle Islands in
21 Sierra Leone. This site is the southernmost wintering site known (Gatter, 1988), and although
22 some isolated observations of small number of wintering individuals are mentioned for Ghana
23 (Grimes, 1987) and Nigeria (Meininger, 1988), no recent data are available on these two
24 countries.

25 Eleven out of fifteen individuals previously ringed at Jeliana in 2007-2008 had returned to their natal site
26 to breed in 2012. Site fidelity is a common feature in several gull and tern species (see the review by
27 McNicholl, 1975, Coulson, 2001), to increase breeding success (Greenwood and Harvey, 1982) and
28 reduce the cost of prospecting for a new breeding site (Naves et al., 2006). Inter-colony movements were
29 observed at Jeliana in 2012, this has been reported in several tern species e.g. Sooty Terns *Sterna fuscata*
30 in Seychelles (Feare and Lesperance, 2002) and nominate Crested Terns *T. bergii bergii* (Crawford et al.,
31 2002). The presence of individuals originating from a different site indicates that mate choice was made
32 at either the wintering area or staging site(s) prior to arrival to Libyan colony site.

1 The present study also accurately confirmed the breeding and migration phenology of this
2 localized population. Threats to this species and other migratory seabirds/waterbirds on the
3 flyway and wintering ranges need to be addressed and conservation measures should be taken by
4 different countries. In addition, protection of breeding sites in Libya is also a priority for the
5 maintenance of this limited population that face the threat of extinction from the Mediterranean.

6

7 **Acknowledgements:**

8 Authors would like to present their sincere thanks to the Environment General Authority (EGA)
9 and its offices in Benghazi and Bayda. They thank the directors of the Nature Conservation
10 Department Dr. Ali Alkekli and Mr. Elmakee Alagil for their logistical support during the
11 fieldwork of this project, and the Zuitina Oil Company for providing a boat to visit Gara Island
12 and accommodation for our team. They are grateful to the Regional Activity Centre for Specially
13 Protected areas (UNEP-MAP-RAC/SPA), courtesy of Mrs. Lobna Ben Nakhla, for the technical
14 and financial support. The authors thank both the Italian Institute for Environmental Protection
15 and Research (ISPRA) and Birdlife Malta for providing the colour and metal rings respectively,
16 and the French Conservatoire du Littoral and Tour du Valat for providing support to the field
17 work. Thanks to the several volunteers from the Environment General Authority in Tripoli,
18 Benghazi, Bayda and Tobruk. Finally, we are grateful to the late president of Libyan Birds
19 Society Mr. Wagih Bashimam who participated actively in the ringing campaigns of 2007 and
20 2008. The work from 2009-2012 was supported by a Postgraduate Grant from the Libyan
21 Ministry for Higher Education (Grant no. 235/2008) for the PhD of the main author.

22

23

24 **References:**

25 Azafzaf, H., Etayeb., K. and Hamza., A. 2006. Report on the census of Lesser Crested Tern
26 *Sterna bengalensis* in the Eastern coast of Libya. (1-7 August 2006). Tunis: Regional Activities
27 Centre/Special Protected Areas (MAP/UNEP), Environment General Authority (Libya) and the
28 African-Eurasian Waterbird Agreement (UNEP/AEWA).

29 Balmer, D., Coiffait, L., Clark, J. and Robinson, R. 2008. *Bird Ringing: a Concise Guide.* ,
30 Thetford, British Trust for Ornithology.Pp75.

31 Barlow, M. 1998. Movements of Caspian Terns (*Sterna caspia*) from a colony near
32 Invercargill, New Zealand, and some notes on their behaviour. *Notornis*, 45, 193-219.

- 1 Bundy, G. 1976. The birds of Libya : an annotated check-list. British Ornithologists' Union,
2 London.
- 3 Coulson, J. 2001. Colonial breeding in seabirds. In: SCHREIBER, E. A (ed.) Biology of
4 marine birds. Boca Raton: CRC Marine Biology Series.
- 5 Crawford, R. J. M., Cooper, J., Dyer, B. M., Upfold, L., Venter, A. D., Whittington, P. A.,
6 Williams, A. J. and Wolfaardt, A. C. 2002. Longevity, inter-colony movements and breeding of
7 Crested Terns in South Africa. *Emu*, 102, 265-273.
- 8 EGA-RAC/SPA-WCT 2012. *Atlas of Wintering Waterbirds of Libya, 2005-2010*, Tunisia,
9 Imprimerie COTIM.
- 10 Etayeb, K. and Essghaier, M. 2007. Breeding of marine birds on Farwa Island, western Libya.
11 *Ostrich*, 78, 419-421.
- 12 Feare, C. and Lesperance, C. 2002. Intra- and Inter-colony Movements of Breeding Adult
13 Sooty Terns in Seychelles. *Waterbirds*, 25, 52-55.
- 14 Gatter, W., 1988. The Birds of Liberia (West Africa). A preliminary list with status and open
15 questions. *Verh orn Ges Bayern*, 24, pp.689-723.
- 16 Goodman, S. and Meininger, P. 1989. *The Birds of Egypt*. Oxford, England, Oxford University
17 Press. .
- 18 Greenwood, P. J. And Harvey, P. H. 1982. The Natal and Breeding Dispersal of Birds. *Annual*
19 *Review of Ecology and Systematics*, 13, 1-21.
- 20 Grimes, L. G. 1987. *The birds of Ghana : an annotated check-list*, British Ornithologists'
21 Union.Pp 276
- 22 Hamza, A. 2014. Breeding Ecology, Migration and Population Genetics of Lesser Crested
23 Terns *Thalasseus bengalensis emigrata*. Unpublished PhD Thesis, University of Hull, UK, pp
24 248.
- 25 Hamza, A., H. Azafzaf, N. Baccetti, E. Bourass, J. J.borg, P. Defos du Rau, A. Saied, J. Sultana
26 & Zenatello, M. 2007. Report on census and ringing of Lesser Crested Tern *Sterna bengalensis*
27 in Libya, with a preliminary inventory of Libyan islands (2-10 August 2007). : Regional
28 Activities Centre/Special Protected Areas (MAP/UNEP), Environment General Authority
29 (Libya).

- 1 Hamza, A., Azafzaf, H., J. Yahia .2012. State of knowledge and population trends of the Lesser
2 Crested Tern *Sterna bengalensis emigrata* in the Mediterranean: threats identified and proposed
3 actions for small islands in the Mediterranean, In YÉSOU, BACCETTI and SULTANA (Eds.)
4 Ecology and Conservation of Mediterranean Seabirds and other bird species under the
5 Barcelona Convention - Proceedings of the 13th Medmaravis Pan-Mediterranean Symposium,
6 Alghero 2011: 171-177.
- 7 Hashmi, D. 2000. Opportunities for monitoring seabirds and cetacens in the Strait of Gibraltar.
8 In Yésou P. and Sultana J. (eds.), Monitoring and Conservation of Birds, Mammals and Sea
9 Turtles in the Mediterranean and Black Seas, Proceedings of the 5th Medmaravis Symposium,
10 Gozo, Malta, 29 September-3 October 1998. Environment Protection Department, Malta, pp.
11 176-191.
- 12 Holmström, N., Hägg, J., Johansson, B., Nilsson, C. & Oomen, M. 2005. Seawatching At N'gor
13 Senegal. Retrieved from <http://senegal.seawatching.net/seawatch.html> .
- 14 Isenmann, P., Gaultier, T., El Hili, A., Azafzaf, H., Dlensi, H. and Smart, M. 2005. *Oiseaux de*
15 *Tunisie - Birds of Tunisia.*, Paris, France, SEOF Editions.
- 16 Isenmann, P., Benmergui, M., Browne, P., Diam Ba, A., Hamallah, C., Diawara, Y. and Ould
17 Sidaty, Z. 2010. *Birds of Mauritania*, Paris, Societé d'Études Ornitologiques de France.
- 18 McNicholl, M. K. 1975. Larid Site Tenacity and Group Adherence in Relation to Habitat. *The*
19 *Auk*, 92, 98-104.
- 20 Meininger, P. L. 1988. A Preliminary investigation of Tern catching in Senegal, Winter
21 1987/88, Cambridge, UK, ICBP.
- 22 Meininger, P. L., Wolf, P. A., Hadoud, D. A. and Essghaier, M. F. 1994. Rediscovery of Lesser
23 Crested Terns breeding in Libya. *British Birds*, 87, 160-170.
- 24 Naves, L. C., Monnat, J. Y. and Cam, E. 2006. Breeding performance, mate fidelity, and nest
25 site fidelity in a long-lived seabird: behaving against the current? *Oikos*, 115, 263-276.
- 26 Nichols, J. D., M. C. Runge, F. A. Johnson, and B. K. Williams. 2007. Adaptive harvest
27 management of North American waterfowl populations: a brief history and future prospects.
28 *Journal of Ornithology* 148:343-349.
- 29 Paracuellos, M. and Jerez, D. 2003. A comparison of two seabird communities on opposite
30 coasts of the Alboran Sea (western Mediterranean). *Scientia Marina*, 67, 117-123.

- 1 Schricke, V., Triplet, P. and Yesou, P. 2001. French contribution to the knowledge of western
2 Palaearctic water birds overwintering in the Senegal River delta. *Alauda*, 69, 135-148.
- 3 Sharp, S. P. 2009. Bird ringing as a tool for behavioural studies *Ringing and Migration*, 24,
4 213-219.
- 5 Shiomi, K., U. Lötberg, and S. Akesson. 2015. Seasonal distributions of Caspian Terns
6 *Hydroprogne caspia* from Swedish populations, revealed by recoveries and resightings of
7 ringed birds. *Ringing and Migration* 30:22-36.
- 8 Smart, M., M. F. Essghaier, K. Etayeb, A. Hamza, H. Azafzaf, N. Baccetti, P. D. Rau, And H.
9 and Dlensi. 2006. Wetlands and wintering waterbirds in Libya, January 2005 and 2006.
10 *Wildfowl* 56:172-191.
- 11 Spina, F. 1999. Value of ringing information for bird conservation in Europe. *Ringing and*
12 *Migration* 19:29-40.
- 13 Valentine, K. 2008. RBT Sierra Leone Trip Report: 9th – 22nd December 2008. Cascades,
14 South Africa: Rockjumper Birding Tours.
- 15 Viana, D. S., L. Santamaría, T. C. Michot and J. Figuerola. 2013. Migratory strategies of
16 waterbirds shape the continental-scale dispersal of aquatic organisms. *Ecography* 36:430-438.
- 17 Wernham, C. V., Toms, M. P., Marchant, J. H., Clark, J. A., Siriwardena, G. M. and S. R.
18 Baillie. 2002. *The Migration Atlas: Movements of the Birds of Britain and Ireland* T. and A.D.
19 Poyser, London.