Letter in response to Lovell 2018

Acute-on-chronic breathlessness: recognition and response

Authors: Ann Hutchinson, Miriam Johnson and David Currow

Lovell et al are to be commended for their description of the widespread effects of breathlessness within the construct 'total breathlessness'.(1) Despite the large body of evidence on the experience of people living with chronic breathlessness, clinicians still struggle with its recognition and assessment. Even when recognised, the symptom is undertreated (2) and is experienced by people over many years (3-5) and is associated with repeated unplanned presentations to health services.(6)

In our prospective observational study of 1,212 patients presenting to the emergency department (ED) by ambulance (5), 20% presented due to acute-on-chronic breathlessness (acute worsening of chronic breathlessness). The concept of acute-on-chronic breathlessness builds on the definition of chronic breathlessness syndrome (7) and mirrors concepts of disease (acute-on-chronic renal failure) or symptoms (acute-on-chronic pain). Acute-on-chronic breathlessness is a construct beyond the 'dyspnea crisis' which is limited in its definition to late-stage disease (8) and encompasses all forms of episodic breathlessness, triggered or untriggered.(9)

In our ED study, one third of people with acute-on-chronic breathlessness were discharged home but without evidence of a plan to manage the ongoing chronic breathlessness. Unless clinicians (in any setting) recognise *both* the acute and chronic aspects of acute-on-chronic breathlessness the life-experience of chronic

Acute-on-chronic breathlessness: recognition and response

breathlessness remains invisible, denies the patient access to evidence-based interventions (10, 11) and misses opportunities to lessen the likelihood of representations.(6)

Community clinicians struggle with how to identify, assess and manage chronic breathlessness. There are now useful frameworks to guide clinicians' assessment (12) and management.(13) As described in the Breathing Space concept (14), the clinician plays a pivotal role in recognising background chronic breathlessness – a step often overlooked.

Defining chronic breathlessness syndrome (7) has been an important first step. There is a new imperative: if a patient presents with acute breathlessness, there is now a responsibility for clinicians to determine whether this is an isolated episode of acute breathlessness or a presentation of acute-on-chronic breathlessness.

1. Abernethy, Wheeler. Total dyspnoea. Current Opinion in Supportive and Palliative Care. 2008;2(2):110-3.

2. Ahmadi Z, Sandberg J, Shannon-Honson A, Vandersman Z, Currow DC, Ekström M. Is chronic breathlessness less recognised and treated compared with chronic pain? A case-based randomised controlled trial. European Respiratory Journal. 2018;52(3):1800887.

Acute-on-chronic breathlessness: recognition and response

3. Johnson MJ, Bowden JA, Abernethy AP, Currow DC. To what causes do people attribute their chronic breathlessness? A population survey. Journal of palliative medicine. 2012;15(7):744-50.

4. Currow DC, Dal Grande E, Ferreira D, Johnson MJ, McCaffrey N, Ekström M. Chronic breathlessness associated with poorer physical and mental health-related quality of life (SF-12) across all adult age groups. Thorax. 2017;72(12):1151-3.

 Hutchinson A, Pickering A, Williams P, Bland JM, Johnson MJ. Breathlessness and presentation to the emergency department: a survey and clinical record review.
BMC Pulmonary Medicine. 2017;17(1):53.

6. Nunez S, Hexdall A, Aguirre-Jaime A. Unscheduled returns to the emergency department: an outcome of medical errors? Quality and Safety in Health Care. 2006;15(2):102-8.

7. Johnson M, Yorke J, Hansen-Flaschen J. Towards an expert consensus to delineate a clinical syndrome of chronic breathlessness. Eur Respir J. 2017:13993003.02277-2016.

8. Mularski RA, Reinke LF, Carrieri-Kohlman V, Fischer MD, Campbell ML, Rocker G, et al. An official American Thoracic Society workshop report: assessment and palliative management of dyspnea crisis. Annals of the American Thoracic Society. 2013;10(5):S98-S106.

9. Simon ST, Higginson IJ, Benalia H, Gysels M, Murtagh FE, Spicer J, et al. Episodes of breathlessness: Types and patterns–a qualitative study exploring experiences of patients with advanced diseases. Palliative medicine. 2013;27(6):524-32. 10. Ekström M, Bajwah S, Bland JM, Currow DC, Hussain J, Johnson MJ. One evidence base; three stories: do opioids relieve chronic breathlessness? Thorax. 2018;73(1):88-90.

11. Bausewein C, Schunk M, Schumacher P, Dittmer J, Bolzani A, Booth S. Breathlessness services as a new model of support for patients with respiratory disease. Chronic respiratory disease. 2018;15(1):48-59.

12. Hopkinson NS, Baxter N. Breathing SPACE—a practical approach to the breathless patient. npj Primary Care Respiratory Medicine. 2017;27(1):5.

13. Spathis A, Booth S, Moffat C, Hurst R, Ryan R, Chin C, et al. The Breathing, Thinking, Functioning clinical model: a proposal to facilitate evidence-based breathlessness management in chronic respiratory disease. Primary Care Respiratory Society UK. 2017.

14. Hutchinson A, Barclay-Klingle N, Galvin K, Johnson MJ. Living with breathlessness: a systematic literature review and qualitative synthesis. European Respiratory Journal. 2018;51(2):1701477.