

1 **Refinement of acoustic-tagging protocol for twaite shad *Alosa fallax* (Lacépède), a**
2 **species sensitive to handling and sedation**

3 Jonathan D. Bolland^{1*}, Andrew D. Nunn¹, Natalie V. Angelopoulos¹, Jamie R. Dodd¹, Peter
4 Davies^{1,2}, Catherine Gutmann Roberts²; J. Robert Britton² and Ian G. Cowx¹

5 ¹ University of Hull International Fisheries Institute, University of Hull, England, HU6 7RX

6 ² Department of Life and Environmental Sciences, Faculty of Science and Technology,
7 Bournemouth University, BH12 5BB, UK

8 * Corresponding author: j.bolland@hull.ac.uk

9 **ABSTRACT**

10 Telemetry investigations to gather essential information about fish migrations are reliant on
11 the behaviour, condition and survival of the animals being unaltered by the tagging procedure.
12 Twaite shad (*Alosa fallax* Lacépède; 'shad') is a threatened clupeid fish for which there is a
13 considerable knowledge gap on their anadromous movements. They are also reported to be
14 sensitive to handling and anaesthesia, resulting in practical difficulties in tag implantation;
15 previous investigations externally attached tags without sedation. The aim of this study was
16 to incrementally refine the acoustic-tagging protocol for shad *via* application of a previously
17 un-tried anaesthetic (i.e. tricaine methanesulphonate (MS-222)) and by surgical implantation
18 of the tag in the peritoneal cavity. All captured shad ($n = 25$) survived handling, anaesthesia
19 and tagging, and were detected moving upstream after release. Surgically implantation ($n =$
20 5) was significantly faster than externally mounting the tag ($n = 20$) and time to recover was
21 similar. Total upstream movement, total movement, residence time in receiver array and
22 speed of upstream movement were statistically similar for externally and internally tagged fish.
23 Post-spawning, a large proportion (68 %) of tagged fish returned to the estuary, downstream
24 of the receiver array. Internal tagging under anaesthesia is recommended for studying
25 anadromous movements of shad, given welfare benefits during surgery and once at liberty,
26 thus increasing the likelihood of tagged fish performing natural behaviours. Further,
27 implantation of tags programmed to last many years enables multiple spawning migrations by

28 the same individuals to be studied, which would lead to substantial advances in ecological
29 knowledge and potentially reduce the number of fish tagged.

30 *Keywords*

31 Anadromous; Animal welfare; Iteroparous; Regulated procedure; Surgical implantation;
32 Telemetry

33 **1. Introduction**

34 Fish telemetry investigations are routinely performed to gather essential information about
35 migrations, habitat use, predator–prey interactions and responses to anthropogenic impacts,
36 to help protect species and the environments they inhabit (Hussey et al., 2015). Such studies
37 are reliant on the behaviour, condition and survival of the animals being unaltered by the
38 tagging procedure (Cooke et al., 2013). This has resulted in a considerable amount of work to
39 identify maximum tag burden, optimal tag implantation location and most appropriate methods
40 of anaesthesia (Broadhurst et al., 2009; Ross & Ross, 2009). There have been considerable
41 refinements in internal tagging procedures, with tags often retained for the lifetime of the fish
42 with minimal long-term impact (Jepsen et al., 2002; Bridger and Booth, 2003; Cooke et al.,
43 2011). External tag attachment remains important in some studies and species, including
44 those considered to be sensitive to handling (Jepsen et al., 2015; Johnson et al., 2015).
45 However, tags can become fouled, increase drag during swimming, cause irritation and harm
46 as the fish grow, potentially impairing individual behaviour and increasing mortality risk
47 (Mulcahy, 2003; Cooke et al., 2013; Jepsen et al., 2015).

48 Twaité shad *Alosa fallax* (Lacépède) ('shad' hereafter) is an anadromous clupeid fish
49 species that was once abundant and widespread across Europe (Arahamian et al., 2003).
50 Their populations have, however, declined considerably in the last century. Causal factors
51 relate primarily to anthropogenic disturbances, especially the construction of weirs in the lower
52 reaches of rivers that reduce access to spawning areas (Jolly et al., 2012). The species is
53 listed on Appendix III of the Bern Convention and Annexes II and V of the EU Habitats
54 Directive. Despite their conservation importance, their anadromous spawning migration

55 remains under-studied primarily due to difficulties tagging shad, a species reported to
56 adversely react to handling and sedation (with 2-phenoxyethanol) that results in high mortality
57 rates (Rooney and King, 2014; Breine et al., 2017). To overcome these challenges, recent
58 investigations have externally mounted acoustic tags without sedation because it is less
59 invasive and thought to be quicker than surgical implantation (Rooney and King, 2014; Breine
60 et al., 2017). Although these studies were successful, Breine et al. (2017) recommended
61 further research on the effects of anaesthesia, handling and tagging on shad.

62 The aim of this study was to refine the acoustic-tagging protocol for shad, giving due
63 consideration to their sensitivity to handling and sedation, to provide short-term welfare
64 benefits during surgery and long-term welfare benefits while at liberty, thus enabling
65 expression of natural behaviours. Objectives were to: (1) refine the external tag attachment
66 protocol of Breine et al. (2017) *via* application of previously un-tried anaesthetic (i.e. tricaine
67 methanesulphonate (MS-222)); (2) further refine the procedure by surgically implanting the
68 tag within the peritoneal cavity; and (3) quantify the impacts of the tagging methods through
69 comparison of shad movement. As shad are iteroparous and, potentially, philopatric (King and
70 Roche, 2008), implantation of tags programmed to last many years enables multiple spawning
71 migrations by the same individuals to be studied, which would lead to substantial advances in
72 ecological knowledge.

73 **2. Methods**

74 *2.1. Fish capture and iterative tagging process*

75 The refinement of the shad tagging protocol was completed during the 2017 shad spawning
76 migration in the River Severn, Western England (Fig. 1). Twenty-five shad were captured from
77 two locations, downstream of Maisemore ($n = 8$) and Upper Lode weirs ($n = 17$), with 23
78 captured by angling (small lure with single barbless hook) and two with a seine net (30-m long,
79 2-m deep and 10-mm mesh) (Table 1). Tagging was an iterative process involving small
80 batches of fish to minimise the number of fish with compromised welfare if tagging was
81 unsuccessful and to enable refinements between batches. Thus, the initial 3 captured fish
82 were externally tagged under general anaesthesia (batch 1), with tagging only recommencing

83 once a receiver 14.8-km upstream of the release location revealed the fish had recovered
 84 sufficiently to continue their upstream movement. The decision to commence surgically
 85 implanting tags in the body cavity (batch 4) was only taken after a further 11 shad had been
 86 successfully tagged externally (batch 2 and 3). The final six fish (batch 5) were tagged
 87 externally because there was no opportunity to establish if the internally tagged fish (batch 4)
 88 had been detected on the receiver upstream of the release location.


89 Table 1. Capture date, sample size and capture, release (DS = downstream, US = upstream)
 90 and tag locations of twaite shad tagged in five batches on the River Severn.

Batch	Date	<i>n</i>	Capture location	Release location	Tag location
1	11/5/17	3	DS Maisemore Weir	US Maisemore Weir	External
2	17/5/17	5	DS Upper Lode Weir	US Upper Lode Weir	External
3	17/5/17	6	DS Upper Lode Weir	DS Upper Lode Weir	External
4	22/5/17	5	DS Maisemore Weir	US Maisemore Weir	Internal
5	31/5/17	6	DS Upper Lode Weir	DS Upper Lode Weir	External

91

92 2.2. External and internal tagging procedures

93 Prior to tagging, acoustic tags (20-mm long x 7-mm diameter (V7), 1.6-g weight in air and 29-
 94 mm long x 9-mm diameter (V9), 4.7-g weight in air; www.vemco.com) were activated and
 95 tested with a hand-held detector to verify they were transmitting; weight in air did not exceed
 96 2% of fish mass. Following capture, fish were briefly held in water filled containers (100 L)
 97 prior to their general anaesthesia (MS-222; 0.4-g per 10-L of water). All fish were inspected
 98 for signs of pre-existing injury and disease; no captured fish were excluded from tagging.
 99 Whilst being sedated, the fish were measured (fork length, nearest mm; mean \pm S.D.: 354 \pm
 100 37 mm, range = 302–420 mm), and scale sample and a fin biopsy taken (for use in
 101 complementary studies). The influence of the anaesthetic was visually assessed using body,
 102 opercula and eye movements, with fish only removed following their lack of a response to
 103 touch, loss of ability to balance and the cessation of pectoral fin and eye movements.



104
 105 Figure 1. A map of acoustic receiver locations (black dots) in the River Sever catchment,
 106 including impediments to fish migration (black lines). Maisemore and Llanthony weirs
 107 represent the tidal limit, and Maisemore and Upper Lode weirs were capture locations.

108 Externally mounted tags were attached using surgical thread (Ethilon) passed through
 109 the dorsal musculature using hollow needles and held in place using a rubber plate and

110 aluminium sleeves (as per Breine et al., 2017). Surgically implanted tags were disinfected with
111 providone-iodine and rinsed with saline solution before being implanted into the body cavity
112 through a ventro-lateral incision made with a scalpel, anterior to the muscle bed of the pelvic
113 fins. The incision was closed with an absorbable monofilament suture. Fish were held in a
114 clean V-shaped foam support and their eyes were covered with a damp cloth during surgery.
115 All fish were treated in compliance with the UK ASPA (1986) Home Office licence number PPL
116 60/4400.

117 After surgery, fish were transferred to a damp sling for weighing (to 25 g; mean \pm S.D.
118 = 547 \pm 173 g, range = 300–850 g) and then returned to the river, being held whilst they
119 orientated towards the flow and were only released when they had regained balance, body
120 reflexes and swimming ability. This was considered preferable to holding fish in tanks with
121 water circulation and aeration, as shad have been recorded to die during transportation and
122 at fish farms (Clough et al., 2004). Fish were released upstream of Maisemore Weir ($n = 8$),
123 downstream of Upper Lode Weir ($n = 12$) and upstream of Upper Lode Weir ($n = 5$) as part of
124 the wider investigation (Table 1). Catchment-wide migration was examined using 23
125 strategically located acoustic receivers (Vemco; www.vemco.com) (Fig. 1); no fish were
126 detected on the most upstream receivers.

127 2.3. *Data analysis*

128 Time taken for anaesthesia, surgery and recovery when externally and internally tagging shad
129 was compared using t-tests (non-normal data (Shapiro test) were log-transformed). It was not
130 possible to recapture tagged shad to assess general health and condition, external tag fouling
131 or healing of incisions for internally implanted tags. Instead, movements of fish in the river
132 were used as evidence that the fish had recovered from handling, anaesthesia and surgery.
133 Specifically, the amount of upstream movement (i.e. sum of all upstream movements), total
134 movement (i.e. sum of all up and downstream movements), and residence time in the receiver
135 array (i.e. number of days from release to first detection on last receiver) were calculated for
136 each fish. In addition, the speed of upstream movement between receivers was calculated
137 (distance between receivers / last detection on upstream receiver – first detection on

138 downstream receiver). The movements of fish in batches 1 and 4, captured and released at
 139 the same location but with external and internal tag attachment, were compared using t-tests
 140 (non-normal data (Shapiro test) were log-transformed) to quantify impacts of the tagging
 141 methodology. Both movement and speed metrics represent minimum estimates, as they are
 142 measured at the resolution of receiver separation, thus back and forth movements between
 143 receiver detection area are undetected. The fates of individual fish were broadly separated
 144 into those that returned to the estuary and those that were assumed to have died in the river,
 145 though the latter could not be separated from tag failure or loss, and the potential cause of
 146 death could not be determined (e.g. tagging induced, natural predation event, tagging-induced
 147 predation event or natural mortality after spawning). Data analysis was performed primarily in
 148 Microsoft Excel and statistical comparisons performed using R statistical software (version
 149 3.4.3, R Core Team 2017), with movement speed analysis in the V-Track package (Campbell
 150 et al., 2012).

151 **3. Results**

152 All 25 fish caught during the investigation survived capture, handling, sedation and tagging,
 153 and were assessed as being in satisfactory condition prior to be returned to the river. The time
 154 taken for anaesthesia was similar ($t = -0.054171$, d.f. = 5.5144, $P = 0.959$) whereas internal
 155 implantation was significantly faster than external attachment (t -test on logged data; $t = -88.36$,
 156 d.f. = 32.372, $P < 0.001$), both usually within four minutes (Table 2). The mean time to recover
 157 was also similar (t -test on logged data; $t = -1.9709$, d.f. = 7.8191, $P = 0.085$), and the longest
 158 recovery did not exceed six minutes for either treatment group (Table 2).

159 Table 2. Time (seconds; mean \pm 95% C.I. (min.–max.)) taken for anaesthesia, surgery and
 160 recovery when externally and internally tagging shad with acoustic tags.

Procedure stage	External ($n = 20$)	Internal ($n = 5$)
Anaesthesia	112 \pm 12 (60–182)	113 \pm 28 (70–150)
Surgery	113 \pm 10 (83–179)	117 \pm 12 (104–136)
Recovery	149 \pm 28 (85–356)	196 \pm 54 (140–301)

161

162 All shad were detected moving upstream in fresh water, i.e. against the flow. Of all the
 163 batches, the first batch of fish (external tag) had the greatest mean upstream movement (61.1
 164 \pm 51.7 km) and mean total movement (122.9 \pm 95.2 km), whereas the fourth batch (internal
 165 tag) spent the longest mean time in the river (21.4 \pm 8.8 days) and fastest mean speed of
 166 upstream movement (1.10 \pm 0.32 m/s) (Table 3). Fish in batches 1 and 4 were captured and
 167 released at the same location with external and internal tags, respectively, and had similar
 168 upstream movements (*t*-test on logged data; *t* = 0.095988, d.f. = 3.7202, *P* = 0.926), total
 169 movements (*t*-test on logged data; *t* = 0.31356, d.f. = 4.3419, *P* = 0.768), times in the river (*t*-
 170 test; *t* = -0.61932, d.f. = 5.5427, *P* = 0.560) and speed of upstream movements (*t*-test; *t* =
 171 2.1894, d.f. = 6, *P* = 0.0711) (Table 3). The individual fish with the greatest upstream (138.0
 172 km) and total movements (281.4 km), and longest time in the river (29.8 days) had an internal
 173 tag, whereas the fastest upstream movements (1.79 m/s) was by a fish that had an external
 174 tag.

175 Table 3. Mean \pm 95% C.I. (min.–max.) upstream movement (km), total movement (km),
 176 residence time in the receiver array (days) and speed of upstream movement (m/s) for shad
 177 tagged in five batches on the River Severn.

Batch	Upstream movement (km)	Total movement (km)	Time in river (days)	Speed of upstream movements (m/s)
1	61.1 \pm 51.7 (27.7–113.1)	122.9 \pm 95.2 (60.4–218.5)	18.3 \pm 4.4 (13.9–21.2)	0.60 \pm 0.19 (0.50–0.80)
2	16.4 \pm 11.4 (4.0–37.7)	50.8 \pm 26.5 (19.0–96.5)	12.8 \pm 5.0 (6.6–23.3)	0.54 \pm 0.14 (0.30–0.73)
3	14.4 \pm 11.7 (1.0–33.9)	46.2 \pm 28.7 (5.7–91.4)	8.4 \pm 4.5 (0.2–16.2)	0.51 \pm 0.17 (0.31–0.77)
4	58.0 \pm 39.6 (30.7–138.0)	112.1 \pm 83.6 (51.0–281.4)	21.4 \pm 8.8 (9.3–29.8)	1.10 \pm 0.32 (0.72–1.52)

5	15.5 ± 11.6	49.0 ± 16.4	8.9 ± 5.7	1.09 ± 0.38
	(2.0–38.6)	(28.3–73.7)	(1.5–19.1)	(0.55–1.79)

178

179 Seventeen (68%) of the tagged shad performed a downstream migration to the estuary
180 between 25 May and 21 June 2017, 14.7 ± 3.9 days after tagging. Eight fish were assumed
181 to have died in the river (though tag failure or loss could not be ruled out) but were tracked for
182 a similar amount of time, i.e. 10.6 ± 8.2 days. The one exception (external tag) was last
183 detected 5 h after release, 5.7 km upstream of its release location. Four fish (external = 2 and
184 internal = 2) were last detected in the vicinity of a suspected spawning location 9–27 days
185 after release, three of which moved downstream after release and subsequently returned to
186 fresh water. Three fish (external = 2 and internal = 1) were last detected moving downstream
187 5, 7 and 12 days after release, each having moved a minimum of 18.7, 4.0 and 36.3 km,
188 respectively, in an upstream direction while in fresh water.

189 **4. Discussion**

190 During this investigation, twaite shad, a threatened anadromous fish species that is sensitive
191 to handling and sedation, were successfully anaesthetised which enabled tags to be surgically
192 implanted into the peritoneal cavity. These findings are contrary to Rooney and King (2014)
193 who reported mortality of shad anaesthetised with 2-phenoxyethanol and represents a
194 substantial refinement of an accepted tagging protocol (*cf.* Breine et al., 2017). The novel and
195 successful use of MS-222 for shad might be reflective of high variability in species-specific
196 responses to different anaesthetics (e.g. Readman et al., 2017). These refinements have
197 important welfare, ethical and methodological implications for future shad tracking studies.

198 Twaite shad are anadromous and iteroparous. In this study, a large proportion of the
199 tagged fish (68%) migrated downstream to the estuary after undertaking substantial
200 movements upstream and spent an appreciable amount of time in fresh water. This suggested
201 that tagging had little or no impact on their behaviour and that these fish evaded predators
202 (e.g. pike *Esox lucius* L., zander *Sander lucioperca* (L.), otter *Lutra lutra* (L.) and cormorant
203 *Phalacrocorax carbo* L.) and survived spawning. The assumed mortality of individuals that did

204 not return to the estuary (though tag failure or loss could not be ruled out) was considered a
205 result of either natural predation or post-spawning mortality, rather than a direct consequence
206 of being tagged. This is because they performed substantial upstream movements, entered
207 the estuary and returned to fresh water, were last detected at a suspected spawning location
208 and/or residence time was similar to fish that returned to the estuary.

209 A commonly cited advantage of external tagging over surgical implantation is that
210 attachment can be faster (Jepsen et al., 2015; Breine et al., 2017), but internal implantation
211 was significantly faster than external attachment in this investigation. Although there was no
212 evidence of detrimental impacts of externally mounting tags they may have reduced swimming
213 performance through drag or disequilibrium. There are many other long-term benefits of
214 internal implantation to individual fish post-release, including improved tag retention, reduced
215 tissue damage, zero risk of biofouling and zero tag visibility to predators (Cooke et al, 2013;
216 Jepsen et al., 2015). Surgically implanting long-lived tags will also provide substantial
217 advances in ecological knowledge of iteroparous shad by enabling multiple annual spawning
218 migrations of the same individual to be studied. Consequently, the number of fish that need to
219 be tagged could also be reduced, thereby complying with the reduction principle of animal
220 research (Metcalf and Craig 2011). These refinements should be transferable to other fishes
221 considered sensitive to handling and sedation, and should lead to further refinements in
222 tagging procedures during biotelemetry research.

223 **Acknowledgments**

224 Funding was received from the Unlocking the Severn for LIFE Programme, a partnership
225 between the Canal & River Trust, Severn Rivers Trust, Environment Agency and Natural
226 England through funding from the EU LIFE Nature Programme (LIFE15/NAT/UK/000219) and
227 Heritage Lottery Fund (HLF-15-04573). The authors would like to thank all landowners for
228 access and Tewkesbury Popular Angling Association for assisting with fish capture.

229 **References**

- 230 Aprahamian, M.W., Baglinière, J.L., Sabatié, M.R., Alexandrino, P., Thiel, R., Aprahamian,
231 C.D., 2003. Biology, status, and conservation of the anadromous Atlantic twaite shad
232 *Alosa fallax fallax*. Am. Fish. Soc. Symp. 35, 103–124.
- 233 Breine, J., Pauwels, I.S., Verhelst, P., Vandamme, L., Baeyens, R., Reubens, J. and Coeck,
234 J. 2017. Successful external acoustic tagging of twaite shad *Alosa fallax* (Lacépède
235 1803). Fish. Res. 191, 36–40.
- 236 Bridger, C.J., Booth, R.K., 2003. The effects of biotelemetry transmitter presence and
237 attachment procedures on fish physiology and behavior. Rev. Fish Sci. 11, 13–34.
- 238 Broadhurst, B.T., Ebner, B.C., Clear, R.C., 2009. Radio-tagging flexible-bodied fish: temporary
239 confinement enhances radio-tag retention. Mar. Freshwater Res. 60, 356-360.
- 240 Campbell, H.A., Watts, M.E., Dwyer, R.G., Franklin, C.E. 2012. V-Track: software for
241 analysing and visualising animal movement from acoustic telemetry detections. Mar.
242 Freshwater Res. 63, 815-820.
- 243 Clough, S.C., Lee-Elliott, I.E., Turnpenny, A.W.H., Holden, S.D.J., Hinks, C. 2004. The
244 swimming speeds of twaite shad (*Alosa fallax*). R&D Technical Report W2-049/TR3.
- 245 Cooke, S.J., Midwood, J.D., Thiem, J.D., Klimley, P., Lucas, M.C., Thorstad, E.B., Eiler, J.,
246 Holbrook, C. and Ebner, B.C., 2013. Tracking animals in freshwater with electronic tags:
247 past, present and future. Anim. Biotelem. 1, 5.
- 248 Cooke, S.J., Woodley, C.M., Eppard, M.B., Brown, R.S., Nielsen, J.L., 2011. Advancing the
249 surgical implantation of electronic tags in fish: a gap analysis and research agenda
250 based on a review of trends in intraceolomic tagging effects studies. Rev. Fish. Biol.
251 Fish. 21, 127–51.
- 252 Hussey, N.E., Kessel, S.T., Aarestrup, K., Cooke, S.J., Cowley, P.D., Fisk, A.T., Harcourt,
253 R.G., Holland, K.N., Iverson, S.J., Kocik, J.F., Flemming, J.E.M., Whoriskey, F.G., 2015.
254 Aquatic animal telemetry: a panoramic window into the underwater world. Science 348,
255 6240.

256 Jepsen, N., Koed, A., Thorstad, E., Baras, E. 2002. Surgical implantation of transmitters in
257 fish: how much have we learnt? *Hydrobiologia* 483, 239–48.

258 Jepsen, N., Thorstad, E.B., Havn, T., Lucas, M.C., 2015. The use of external electronic tags
259 on fish: an evaluation of tag retention and tagging effects. *Anim. Biotel.* 3, 49.

260 Johnson, M.W., Diamond, S.L., Stunz, G.W., 2015. External attachment of acoustic tags to
261 deepwater reef fishes: an alternate approach when internal implantation affects
262 experimental design. *Trans. Am. Fish. Soc.* 144, 851–859.

263 Jolly, M.T., Aprahamian, M.W., Hawkins, S.J., Henderson, P.A., Hillman, R., O'Maoiléidigh,
264 N., Maitland, P.S., Piper, R., Genner, M.J., 2012. Population genetic structure of
265 protected allis shad (*Alosa alosa*) and twaite shad (*Alosa fallax*). *Mar. biol.* 159, 675-
266 687.

267 King, J.J., Roche, W.K. 2008. Aspects of anadromous Allis shad (*Alosa alosa* Linnaeus) and
268 Twaite shad (*Alosa fallax* Lacépède) biology in four Irish Special Areas of Conservation
269 (SACs): status, spawning indications and implications for conservation designation.
270 *Hydrobiologia* 602, 145–154.

271 Metcalfe, J.D. and Craig, J.F., 2011. Ethical justification for the use and treatment of fishes in
272 research: an update. *J. Fish Biol.* 78, 393-394.

273 Mulcahy, D.M., 2003. Surgical implantation of transmitters into fish. *ILAR journal* 44, 295-306.

274 R Core Team (2017). R: A language and environment for statistical computing. R Foundation
275 for Statistical Computing, Vienna, Austria.

276 Readman, G.D., Owen, S.F., Knowles, T.G., Murrell, J.C., 2017, Species specific anaesthetics
277 for fish anaesthesia and euthanasia. *Sci. Rep.-UK* 7, 7102.

278 Rooney, S., King, J., 2014. Presentation: Use of acoustic telemetry to monitor behaviour
279 during the upriver spawning migration of a diadromous fish, the twaite shad (*Alosa*
280 *fallax*). IFM Tagging and Telemetry Workshop Leeds, England, 22–23 July 2014.

281 Ross, L.G., Ross, B., 2009. *Anaesthetic and sedative techniques for aquatic animals*. John
282 Wiley & Sons.