

This is an author-submitted, peer-reviewed version of a manuscript that has been accepted for publication in the European Respiratory Journal, prior to copy-editing, formatting and typesetting. This version of the manuscript may not be duplicated or reproduced without prior permission from the copyright owner, the European Respiratory Society. The publisher is not responsible or liable for any errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final, copy-edited, published article, which is the version of record, is available without a subscription 18 months after the date of issue publication

To the Editor,

Monocytes and macrophages in chronic sarcoidosis pathology

Kaiser et al make a compelling case for T lymphocytes in the pathogenesis of sarcoidosis (1), particularly when considering the initiation of granulomatous inflammation and patients with acute, self-resolving sarcoidosis (Lofgren's syndrome). However, in some situations it seems that T cells are less important. For example, SCID mice, which have no lymphocytes, develop macrophage-rich granulomas when infected with mycobacteria (2). In clinical practice, powerful suppression of T lymphocytes with anti-rejection medication cannot prevent recurrence of granulomas in one third of patients who have undergone lung transplantation for pulmonary sarcoidosis (3).

Visually, chronic sarcoid granulomas have a well-defined macrophage-rich core and a sparse peripheral ring of lymphocytes. Indeed, pathologists use the term 'naked granulomas' to contrast macrophage-dominant sarcoid pathology with the lymphocyte-rich lesions of tuberculosis. Recently, animal models and studies in patients have highlighted the importance of macrophages and their bone-marrow derived precursors, blood monocytes, in chronic sarcoidosis pathology. Mice genetically engineered with overactive mTORC1 specifically in their myeloid cells spontaneously developed granulomas in the lungs and skin reminiscent of sarcoidosis, and mTORC1 activation in biopsies correlated with disease progression in sarcoidosis patients (4). Several clinical studies have demonstrated heightened inflammatory responses in circulating blood monocytes from sarcoidosis patients (5-7).

In sarcoidosis it is likely that there is a complex interplay between adaptive and innate immunity, represented by T lymphocytes and monocyte-derived macrophages. An important research focus going forward will be to identify the drivers of non-resolving and progressive sarcoidosis that lead to considerable morbidity, loss of quality of life, and economic hardship.

Simon D. Fraser, Simon P. Hart

Respiratory Research Group, Hull York Medical School, Castle Hill Hospital, Cottingham, East

Yorkshire HU16 5JQ, UK

s.hart@hull.ac.uk

References

1. Kaiser Y, Eklund A, Grunewald J. Moving target: shifting the focus to pulmonary sarcoidosis as an autoimmune spectrum disorder. *Eur Respir J.* 2019;54(1):1802153.
2. Goldmann T, Zissel G, Sen Gupta R, Schlaak M, Vollmer E, Müller-Quernheim J. Formation of granulomas in the lungs of severe combined immunodeficient mice after infection with bacillus Calmette-Guerin. *The American journal of pathology.* 2001;158(5):1890-1.
3. Schultz HH, Andersen CB, Steinbruechel D, Perch M, Carlsen J, Iversen M. Recurrence of sarcoid granulomas in lung transplant recipients is common and does not affect overall survival. *Sarcoidosis Vasc Diffuse Lung Dis.* 2014;31(2):149-53.
4. Linke M, Pham HT, Katholnig K, Schnoller T, Miller A, Demel F, et al. Chronic signaling via the metabolic checkpoint kinase mTORC1 induces macrophage granuloma formation and marks sarcoidosis progression. *Nat Immunol.* 2017;18(3):293-302.
5. Talreja J, Farshi P, Alazizi A, Luca F, Pique-Regi R, Samavati L. RNA-sequencing Identifies Novel Pathways in Sarcoidosis Monocytes. *Sci Rep.* 2017;7(1):2720.

6. Fraser SD, Sadofsky LR, Kaye PM, Hart SP. Reduced expression of monocyte CD200R is associated with enhanced proinflammatory cytokine production in sarcoidosis. *Sci Rep.* 2016;6:38689.

7. Crawshaw A, Kendrick YR, McMichael AJ, Ho LP. Abnormalities in iNKT cells are associated with impaired ability of monocytes to produce IL-10 and suppress T-cell proliferation in sarcoidosis. *Eur J Immunol.* 2014;44(7):2165-74.