

ElMokhallalati, Y., Bradley, S. H., Chapman, E., Ziegler, L., Murtagh, F. E., Johnson, M. J., & Bennett, M. I. (2020). Identification of patients with potential palliative care needs: A systematic review of screening tools in primary care. *Palliative Medicine* (v.34, no.8) pp. 989-1005. Copyright © 2020 The Authors. <https://doi.org/10.1177/0269216320929552>

Identification of patients with potential palliative care needs: a systematic review of screening tools in primary care

Yousuf ElMokhallalati^{1*}, Stephen H. Bradley², Emma Chapman¹, Lucy Ziegler¹, Fliss E.M. Murtagh³, Miriam J Johnson³, Michael I Bennett¹

1. Academic Unit of Palliative Care, Leeds Institute of Health Sciences (LIHS), School of Medicine, University of Leeds, UK
2. Academic Unit of Primary Care, Leeds Institute of Health Sciences (LIHS), School of Medicine University of Leeds, Leeds, UK
3. Wolfson Palliative Care Research Centre, Hull York Medical School, University of Hull, Hull, UK

*Correspondence: y.elmokhallalati@leeds.ac.uk

¹Academic Unit of Palliative Care, Leeds Institute of Health Sciences (LIHS), School of Medicine, University of Leeds, Room 10.39, Level 10, Worsley Building, Clarendon Way, Leeds, LS2 9NL, United Kingdom

Abstract

Background

Despite increasing evidence of the benefits of early access to palliative care, many patients do not receive palliative care in a timely manner. A systematic approach in primary care can facilitate earlier identification of patients with potential palliative care needs and prompt further assessment.

Aim:

To identify existing screening tools for identification of patients with advanced progressive diseases who are likely to have palliative care needs in primary health care and evaluate their accuracy.

Design:

Systematic review (PROSPERO registration number CRD42019111568).

Data sources:

Cochrane, MEDLINE, Embase and CINAHL were searched from inception to March 2019

Results:

From 4,127 unique articles screened, 25 reported the use or development of 10 screening tools. Most tools use prediction of death and/or deterioration as a proxy for the identification of people with potential palliative care needs. The tools are based on a wide range of general and disease-specific indicators. The accuracy of five tools was assessed in eight studies; these tools differed significantly in their ability to identify patients with potential palliative care needs with sensitivity ranged 3-94%, and specificity ranged 26-99%.

Conclusion

The ability of current screening tools to identify patients with advanced progressive diseases who are likely to have palliative care needs in primary care is limited. Further research is needed to identify standardised screening processes which are based not only on predicting mortality and deterioration, but also on anticipating the palliative care needs and predicting the rate and course of functional decline. This would prompt comprehensive assessment to identify and meet their needs on time.

Key Words:

Palliative care, terminal care, mass screening, primary health care, systematic review, advance care planning, symptom assessment, terminally ill

What is already known about the topic?

- Earlier initiation of palliative care can improve quality of care for individuals with advanced diseases.
- However, disease trajectories are highly variable, so it is difficult to identify the appropriate time to initiate palliative care.
- A systematic approach may help to identify patients with advanced progressive disease and potential palliative care needs who could benefit from holistic assessment.

What this paper adds

- Most screening tools use prediction of death and/or deterioration as a proxy for the identification of people who are likely to have unmet palliative care needs.
- The performance metrics for these tools were generally poor.

Implications for practice, theory or policy

- More research is needed to identify a standardised and robust screening tool to identify patients with advanced progressive diseases and potential palliative care needs in primary care.
- Future studies should validate screening tools against an appropriate reference standard such as palliative care interview to evaluate their ability to identify patients with potential palliative care needs.
- Identification of patients with advanced progressive diseases and potential palliative care needs process should be supported by comprehensive and holistic assessment to identify their unmet palliative care needs and determine the appropriate care pathway.

Background

In Europe, 85% of people now die of chronic diseases such as cancer, heart disease, stroke and dementia¹. Chronic diseases are characterised by slow progression, fluctuations in trajectory, long duration and uncertainty in prognoses^{2,3}. During advanced stages of chronic life-limiting illnesses, patients usually suffer high levels of pain and other physical and psychological symptoms^{4,5}. At this stage, patients with any progressive disease could benefit from palliative care⁶.

There is evidence from randomised controlled trials that earlier access to specialist palliative care can promote quality of life, reduce hospital length of stay and hospitalisations, and even prolong survival⁷⁻¹³. However, current evidence shows that palliative care is often delivered late in the illness trajectory and access to palliative care is inequitable¹⁴. In the UK, around 90,000 people with advanced progressive conditions who could benefit from palliative care are estimated not to be receiving such care every year¹⁵.

One of the key barriers to providing palliative care on time is the difficulty in identifying patients who could benefit from it^{16, 17}. Once the patient is identified as having potential palliative care needs, their needs can be assessed and addressed in a timely manner. However, not all patients with advanced progressive diseases have unmet palliative care needs. In addition, busy health care professionals cannot provide holistic assessment for all of these patients¹⁸. It has been suggested that a systematic method could facilitate earlier identification of a subset of patients with advanced progressive diseases who are likely to have unmet palliative care needs and hence benefit from palliative care needs assessment^{16, 19}.

Since most people with chronic diseases live at home in the last phase of their life, primary care teams are in the best position to identify patients with potential palliative care needs who could benefit from palliative care needs assessment^{20 21 22}. Two systematic reviews have assessed the screening tools that can be used for the identification of patients who are likely to have unmet palliative care needs. However, neither of them examined the accuracy of the available tools^{16, 23}. This systematic review aimed to identify the existing screening tools for identification of patients with advanced progressive diseases who are likely to have unmet palliative care needs in primary care and synthesize the available evidence regarding their accuracy.

Review questions:

- What screening tools have been used and studied to identify patients with advanced progressive diseases and potential palliative care needs in primary care?
- What are the main characteristics and differences between these screening tools?
- What is the accuracy of these screening tools?

Methods

A positivist approach was used to undertake this systematic review and narrative synthesis of the evidence. This research design was selected because the evidence incorporated a wide range of screening tools and included data from different study designs not suitable for a meta-analysis²⁴. The details of the systematic review protocol are provided in PROSPERO (CRD42019111568). The systematic review was conducted and reported following Preferred Reporting Items for Systematic Reviews and Meta-Analyses protocols (PRISMA-P) guidelines²⁵.

Criteria for considering studies for this review

Types of studies

We included articles that were published in peer-reviewed journals. Commentaries, abstracts, posters, letters to the editor, case reports, reviews and unpublished studies were excluded.

Types of participants

This review included studies examining adults (18 years or older). Studies that reported mixed populations of children and adults were included if data for adults were reported separately. Only studies which included primary care patients or assessed patients in primary care settings were included. Studies which were conducted in mixed settings were included as long as they included primary care patients.

Types of intervention

We included studies that mentioned the use or development of any screening tool to identify patients with advanced progressive diseases who are likely to have unmet palliative care needs in primary health care. Any type of screening tool (electronic or manual) was considered as long as it has been used to identify primary care patients with potential palliative care needs. We also included studies evaluating the ability of the current screening tools to identify patients who could have unmet palliative care needs.

Language

The search was restricted to articles reported in the English language.

Search strategy and Study selection

We searched Cochrane Library, MEDLINE, Embase and CINAHL. A search strategy for MEDLINE is presented in Supplementary File 1. Databases were searched from inception to the end of September 2018. The search was updated in March 2019 to include articles published after September 2018. We searched the reference lists of the included studies and the relevant review articles to make sure that all relevant articles were captured. The

search strategies were created by one reviewer (Y.E) and peer reviewed by a librarian and an information specialist, not otherwise associated with the project. The search results were imported into a reference management software package (EndNote X7) and duplicated references were removed.

Abstracts of all identified studies were independently screened for inclusion by two reviewers. We obtained the full texts of all abstracts that met the inclusion criteria or where there was insufficient information in the abstract alone to determine eligibility. Final article selection was carried out after reading full papers by two reviewers. Disagreements related to screening were resolved through discussion and where necessary a third researcher was consulted.

Data extraction

The characteristics of the included studies and screening tools were extracted prior to synthesis. For studies which assessed the accuracy of the screening tools, specificity, sensitivity, positive predictive value, and negative predictive value were either extracted from the text or calculated from the reported data. Study authors were contacted to resolve any uncertainties, whenever possible. Data were extracted by one reviewer and double checked for accuracy by a second reviewer.

Assessing the risk of bias

Two independent reviewers assessed the methodological quality and risk of bias in the studies that examined the accuracy of the screening tools. Disagreements were resolved first through discussion and then by involving a third reviewer for arbitration. For observational studies, we used the Newcastle-Ottawa Scale to assess the risk of bias²⁶. The methodological quality of these studies was rated on a scale from 0 stars to 9 stars. Studies were classified into groups of low (less than six stars), moderate (7-8 stars), or high (9 stars) quality studies. The quality of randomised clinical trials (RCTs) was assessed using the Cochrane Collaboration's tool for assessing the risk of bias²⁷. This tool evaluates seven possible sources of bias. For each individual domain, studies were classified into low, unclear and high risk of bias.

Strategy for data synthesis

A narrative synthesis was used with information provided in the tables and text to describe and summarise the main findings and features of the included studies and the identified screening tools.

Results

Selection of studies

We identified 6,203 records through the database search and other sources (Figure 1). Of these, 2,076 duplicates were removed, leaving 4,127 publications for title and abstract

screening. Fifty-seven articles remained following review of title and abstract. A further 32 articles were excluded following full-text review, resulting in a total of 25 articles. Of these, only eight evaluated the accuracy of screening tools. No studies were excluded based on their quality assessment.

Figure 1: PRISMA flow chart of the study selection

Characteristics of the included studies

The main characteristics of the 25 articles included in the review are outlined in Table 1^{20, 21, 28-50}. Most studies were published within the last five years (2015–2019). Of those, 17 studies were carried out in the United Kingdom (7), the Netherlands (6) and Spain (4). Twelve studies were observational (prospective observational and cross-sectional), nine studies incorporated mixed methods, three studies were RCTs, and one was a service evaluation study. The majority of the studies included patients with a variety of both cancer and non-cancer conditions. Seventeen studies were conducted exclusively in primary care settings and the remaining studies in mixed settings, including primary care.

Table 1. Characteristics of the included articles

Tool	Author, year	country	Setting	Study design	Study objectives
SQ §	Barnes, 2008	UK	Primary care	Prospective observational study	To identify predictive factors of mortality of heart failure patients in primary care, and to report the sensitivity and specificity of prognostic information from GPs.
SQ §	Moroni, 2014	Italy	Primary care	Prospective cohort study	To determine the prognostic accuracy of GPs asking the SQ about their patients with advanced cancer.
SQ §	Lakin, 2016	USA	Primary care	Retrospective observational study	To assess the SQ performance in primary care setting.
GSF PIG §	Clifford, 2016	UK	Primary care	Service evaluation	To describe the most recent developments and outline the potential of the updated version of Gold Programme.
GSF PIG (Italian version) §	Scaccabarozzi, 2018	Italy	Primary care and home palliative care units	Prospective observational study	To demonstrate the characteristics of patients with palliative care needs, who early identified by GPs and to explore their care process in home palliative care services.
SPICT (German version) §	Afshar, 2018	Germany	Primary care	Mixed methods	To develop, refine, and evaluate SPICT (German version) for its application in primary care.
SPICT (Japanese version) §	Hamano, 2019	Japan	Primary care	Cross-sectional study	To identify the prevalence and characteristics of primary care patients being at risk of deterioration and dying, as determined by SPICT.
SPICT (Japanese version) §	Hamano, 2018	Japan	Primary care	Cross-sectional study	To explore the prevalence and characteristics of family practice patients who need palliative care approach as determined using Supportive and palliative care Indicators Tool.
SPICT §	Highet, 2013	UK	Primary care and hospital	Mixed methods	To refine and test SPICT tool to help multidisciplinary teams, to identify patients at risk of deteriorating and dying in all care settings.
SPICT (Spanish version) §	Fachado, 2018	Spain	Primary care and socio-sanitary services	Mixed methods	To translate, cross-culturally adapt to Spanish, evaluate the Spanish version of the SPICT.
SPICT (2012 version) & SQ §	Mitchell, 2018	Australia	Primary care	RCT	To test whether screening for likely death within 12 months using SPICT and SQ is more effective than an intuition approach.
NECPAL & SQ §	Gómez-Batist e, 2017	Spain	Primary care, hospitals, social health centres and nursing homes	Prospective cohort study	To investigate the predictive validity of the NECPAL and SQ to determine 12- and 24-month mortality.
NECPAL §	Gómez-Batist e, 2014	Spain	Primary care, hospitals, social health centres and nursing homes	Cross-sectional study.	To determine the prevalence of advanced chronically ill patients limited life prognosis in need of palliative care using NECPAL tool
NECPAL §	Gómez-Batist e, 2013	Spain	Primary care, hospitals, social health centres and nursing homes	Mixed methods	To develop the NECPAL tool to identify patient need of palliative care.

RADPAC §	Thoosen, 2016	The Netherlands	Primary care	Cross-sectional study after RCT	To examine whether trained GPs identified more patients in need of palliative care using RADPAC tool and provided multidisciplinary care more than untrained GPs.
RADPAC §	Thoosen, 2015	The Netherlands	Primary care	RCT	To train GPs in identifying patients in need of palliative care and in structuring anticipatory palliative care planning and studied its effect on the quality of life.
RADPAC §	Thoosen, 2012	The Netherlands	Primary care	Mixed methods	To develop a tool for identification of patients with congestive heart failure, COPD and cancer who could benefit from proactive palliative care in primary care.
PALLI §	Vrijmoeth, 2018 _n	The Netherlands	Primary care, central residential settings and intellectual disability physician clinics.	Mixed methods	To evaluate feasibility, construct validity and predictive validity of PALLI.
PALLI §	Vrijmoeth, 2018 _a	The Netherlands	Primary care, central residential settings and intellectual disability physician clinics.	Mixed methods	To describe development of PALLI and to explore its applicability.
The double SQ §	Weijers, 2018	The Netherlands	Primary care	Pilot RCT with caged vignettes	To pilot test whether adding SQ2 to SQ1 prompts GPs to plan for anticipatory palliative care.
Raincine tool ☿	Raincine, 2007	USA	Primary care	Prospective observational study	To develop a methodology to identify patients who may benefit from palliative care and provide estimates of their prevalence in primary care.
AnticiPal (updated version) ☿	Mason, 2018	UK	Primary care	Mixed methods	To refine and evaluate the utility of an electronic ST to help primary care teams screen their patients for people who could benefit from palliative care.
AnticiPal ☿	Mason, 2015	UK	Primary care	Mixed methods	To develop and test an electronic ST in primary care as a tool to improve patient identification and a palliative care approach.
eFI ☿	Stow, 2018 _n	UK	Primary care	Longitudinal population-based study (case control study)	To identify frailty trajectories that could indicate increased risk of dying and the need to consider palliative care.
eFI ☿	Stow, 2018 _a	UK	Primary care	Prospective case control study	To examine if changes in eFI could indicate which individuals are at increased risk of mortality and may require palliative care.

COPD = chronic obstructive pulmonary disease; **GSF PIG** = Gold Standard Framework Proactive Identification Guidance; **SPICt**= the Supportive Palliative Care Identification Checklist; **ST**= Screening Tool; **SQ**= Surprise Question; **eFI**= Electronic Frailty Index; **GPs**= general practitioner; **PALLI**= PALLiative care: Learning to Identify; **RCT**= randomised control trial.

☿ = Electronic tools; § = Paper-based screening tools.

Characteristics of the screening tools

Ten screening tools, used to identify patients with advanced progressive diseases who are likely to have unmet palliative care needs, were identified in this systematic review. Of these, nine were originally designed to identify patients with potential palliative care needs and one was originally developed to identify patients with frailty (Table 2). Four tools were originally developed in the UK (Gold Standard Framework - Proactive Identification Guidance: GSF PIG, Supportive, and Palliative Care Indicators Tool: SPICT, AnticiPal electronic tool, and Electronic Frailty Index: eFI), three in the Netherlands (RADboud indicators for Palliative Care Needs: RADPAC, Palliative care: Learning to Identify in people with intellectual disabilities: PALLI, and the double Surprise Question), two in the United States (Surprise Question; SQ, and early identification tool for palliative care patients “Rainoe tool”), and one tool in Spain (Necesidades Paliativas [Palliative Needs]: NECPAL tool). Seven of the identified tools were paper-based screening tools and three of them were electronic case finding tools. The screening object for most of the identified tools was to identify patients who are at high risk of deteriorating and dying and might benefit from palliative care. The time frame within which symptoms and clinical indicators are assessed varies across the screening tools. The PALLI tool assesses the health status over the last 3–6 months, but the time period for assessment is unspecified for the majority of the symptoms and clinical indicators in all other screening tools. Reviewing care, assessment of needs and initiating discussions about end-of-life needs are some examples of the recommended actions following the screening (Table 3).

Table 2. Summary of the main features of tools which were designed to identify patients with potential palliative

Tool^a	Screening objectives	languages	Target population	Setting (Primary care/GP, Hospital)	Type: paper-based /electronic tool	Completion time	Time frame of assessment	cut
SPICT	To identify people who are at risk of deteriorating and dying and might benefit from palliative care.	English, Japanese, German, Spanish	All	Primary care/GP, hospital	Paper-based	SPICT™: few minutes SPICT-DE: an average of 7.5 minutes. SPICT-ES™: an average of 4 minutes and 45 seconds.	Unspecified for most variables.	SPIC and SPIC indi SPIC indi
NECPAL	To identify people who are at high risk of dying (who likely in need of palliative care).	Spanish	All	Primary care/GP, hospital	Paper-based	NM (one page)	Unspecified for most variables.	NEC indi
RADPAC	To identify people who could benefit from palliative care based on their clinical indicators.	Dutch	COPD, congestive heart failure and cancer patients	Primary care/GP	Paper-based	NM (one page)	Unspecified for most variables.	No
GSF PIG	To identify people who may be in their final stage of life who could benefit from an early palliative approach.	English, Italian	All	Primary care/GP, hospital	Paper-based	NM (one page)	Unspecified for most variables.	GSF indi
PALLI	To identify patients with intellectual disability who may benefit from palliative care via screening deteriorating health, indicative of a limited life expectancy.	Dutch	Patients with intellectual disabilities	Primary care/GP	Paper-based	Mean time of 10.5 minutes (physicians) and 10.1 minutes (daily care professionals)	Previous 3-6 months for all domains except fragility.	No
Surprise Question	To identify patients with poor prognosis who might benefit from palliative care.	English, Italian	All	Primary care/GP, hospital	Paper-based	NM (one question)	NA	SQ que
The double SQ	To identify patients with poor prognosis who might benefit from palliative care.	Dutch, Slovak	All	Primary care/GP, hospital	Paper-based	NM (two questions)	NA	The of S
AnticiPal	To identify patients who potentially have deteriorating health due to one or more advanced illnesses and a likelihood of unmet supportive and palliative care needs.	English	All	Primary care/GP	Electronic	NA	Unspecified for most variables. Previous 18 months for Codes that indicate malignancy.	Ant incl the Typ pan Typ at Con Diff den
Racine tool	To identify people who are at high risk of death (who may benefit from palliative care).	English	All	Primary care/GP	Electronic	NA	NA	Pati elec leas risk

								e.g. con
--	--	--	--	--	--	--	--	-------------

COPD = chronic obstructive pulmonary disease; **GSF PIG** = Gold Standard Framework Proactive Identification Guidance; **SPICT**= the Supportive & palliative Paliativas [Palliative Needs]; **SQ**= Surprise Question; **GPs**= general practitioner; **PALLI**= palliative care: Learning to Identify in people with intellectual disability positive.

^a The most recent version of the tool

Table 3. Summary of the general and specific indicators of deteriorating health and increasing needs in t with potential palliative care needs.

	GSF PIG	SPICT	NECPAL	RADI
Surprise Question (SQ)	Yes	No (SQ was part of some previous versions of SPICT but was removed from the recent versions of SPICT in different languages)	Yes	No
Nutritional decline	Progressive weight loss (>10%) in the past six months Serum albumen <25g/l	Progressive weight loss or remains underweight Low muscle mass	Weight loss > 10%	NM
Functional decline	In bed or chair 50% of the day General physical and performance status decline (Barthel score) and decreasing activities Increasing dependence and need for support	In bed or chair >50% of the day Poor or deteriorating performance status Dependent and increasing need for support	- Karnofsky or Barthel score > 30% loss of two or more activities of the daily living Severe Dependence (Karnofsky <20)	NM
Cognitive decline	-	NM	Minimental/Pfeiffer Decline	NM
Symptom burden	Unstable, deteriorating, complex symptom burden	Persistent symptoms despite optimal treatment	Persistent symptoms (e.g. Pain, weakness, anorexia, dyspnoea, digestive)	NM
Psychosocial decline	NM	NM	Present of emotional stress (Detection of Emotional Distress Scale (DME) > 9) Severe Social Vulnerability (Social and family assessment)	NM
Multi-morbidity	Significant multi-morbidities.	NM	>2 chronic diseases	NM
Urgent/unplanned admissions	Repeated unplanned hospital admissions	Unplanned hospital admission(s).	• > 2 urgent or not planned admittances in last 6 months	NM
Presence of an adverse event	Sentinel event e.g. serious fall, bereavement, transfer to nursing home	NM	Geriatric syndromes (at least two): Falls, pressure ulcers, dysphagia, delirium, recurrent infections.	NM
Others	Considered eligible for DS 1500 payment ^a			
Choice of no further active treatment/ no	Choice for no further active treatment	Chooses to reduce, stop or not have treatment (patient or family)	Limitations of therapeutic effort were mentioned by patient, family or the team	NM

curative treatment available				
Choosing or requiring palliative care	Asks for palliative care by patient	Asks for palliative care by patient or family	Asks for palliative care by patient, family or the team	NM
Additional specific clinical indicators for	Cancer, heart disease, COPD, kidney disease, liver disease, general neurological diseases, Parkinson's disease, motor neurone disease, multiple sclerosis, frailty, dementia, stroke.	Cancer, heart/ vascular disease, kidney disease, liver disease, neurological disease, respiratory disease, dementia/ frailty.	Cancer, COPD, chronic heart disease, chronic neurological disease (CVA, ALS, motor neurone disease, multiple sclerosis), dementia.	COPD failure

COPD = chronic obstructive pulmonary disease **GSF PIG** = Gold Standard Framework Proactive Identification Guidance; **SPICT**= the Supportive & Palliative Needs]; **SQ**= Surprise Question; **PALLI**= PALLiative care: Learning to Identify in people with intellectual disabilities; **NM**= not mentioned

^a DS 1500 is a Form for patients who are terminally ill who are not expected to live for more than six months to rapidly access benefits in the U

Table 3 summarises the general and specific indicators of the screening tools for identification of people with potential palliative care needs in primary care. The Surprise Question is part of all of the paper-based tools (except the RADPAC and the current versions of SPICT). Five tools (GSF PIG, SPICT, NECPAL, PALLI and AnticiPal) contain general indicators for decline and increasing needs such as repeated unplanned hospital admissions, progressive weight loss and functional decline. Only NECPAL and PALLI contain indicators for psychological and cognitive decline. Six tools (GSF PIG, SPICT, NECPAL, RADPAC, PALLI and AnticiPal) contain additional disease-specific clinical indicators of decline for a number of medical conditions. In the paper-based tools, the number of items or questions varied significantly and ranged from 1 to 42. The remainder of this section describes the included tools which used to identify patients who may benefit from palliative care in primary care.

1. The Surprise Question (SQ), which was originally developed by Lynn, is the first tool that has been used for this purpose^{37, 56}. It is utilised as a part of some screening tools or used in isolation. The SQ asks whether the respondent would be surprised if the patient died within a specified time period (usually the next year). The SQ has been widely validated in different settings^{35, 37, 45}. The proportion of patients identified by SQ as having potential palliative care needs across studies ranged from 1.6 to 79%. In those studies applied to patients with advanced progressive diseases, the percentage of patients identified by SQ as having potential palliative care needs ranged from 41% to 79% whereas that applied SQ to more general populations reported percentages between 1.6% and 11.7%^{35, 37, 45}.
2. The double Surprise Question was developed by adding an additional question (SQ2) that asks whether the respondent would be surprised if the patient is still alive after 12 months?" when SQ1 is answered in the negative⁵⁰. The purpose of adding the second SQ was to increase the predictive value of SQ1. The validity of this tool has not been explored yet, although a pilot study concluded that the majority of GPs considered it a useful addition to SQ1⁵⁰.
3. The GSF PIG was developed in the UK^{28, 29}. The tool, which is applicable across care settings, uses the Surprise Question, along with general and disease-specific indicators of decline and increasing need. To the authors' knowledge, there is no underlying research about the development of GSF PIG, and no validation studies have been performed in primary care settings in the UK. The GSF PIG has been translated and adapted for the Italian context²⁸. An Italian study which utilised the GSF PIG among primary care patients found that 0.67% of the patients identified as having a low life expectancy, and palliative care needs²⁸.
4. The SPICT was developed in the UK using a process of literature review, peer review and a prospective case-finding study³³. It is a one-page tool which consists of a

combination of general indicators of deteriorating health and disease-specific indicators. The SPICT had been translated and adapted to Japanese, German, and Spanish settings^{20, 30, 32}. These translated versions (in addition to the original English version) have been validated in a wide range of inpatient and outpatient clinical settings^{20, 30, 32, 33, 37}. Various cut-off scores were used in different versions of SPICT (Table 3). Studies in Australia and Japan that used SPICT among old patients in primary care showed that between 5.1% and 17.3% of these patients could benefit from palliative care^{37, 20, 31}.

5. Mason et al. developed an electronic tool called AnticiPal based on the SPICT criteria⁴³. This electronic tool was developed initially through an iterative process of designing, implementation and testing. In a recent study to evaluate the utility of AnticiPal in Scotland, around 0.8% of 62,708 registered patients at eight GP practices were identified as having potential palliative care needs⁴⁴.
6. The NECPAL tool was developed in Catalonia, Spain based on SPICT and the GSF PIG tools³⁶. This instrument, the NECPAL, is a checklist which combines the SQ with general clinical indicators of severity and progression (e.g. co-morbidity and resource use); and specific indicators for some medical conditions. NECPAL has been validated in a wide variety of care settings^{15, 34-36}. Recent Spanish observational studies which conducted in multiple setting including primary care settings found that 1.5% of primary care patients and 73.7% of patients with advanced progressive diseases met the NECPAL criteria and could benefit from palliative care^{34, 35}.
7. The RADPAC tool was developed in the Netherlands through a three-step process comprising a literature search, focus group interviews and a Delphi study with GPs³⁸. The RADPAC tool contains specific indicators for congestive heart failure, chronic obstructive pulmonary disease (COPD), and cancer; although it does not include the Surprise Question or general clinical indicators which can be applied to all patients. A Dutch RCT on the effects of training GPs in early identification of patients who could benefit from palliative care using the RADPAC tool did not find any differences between the intervention and control group in out-of-hours contacts, contacts with their GP, hospitalisations and place of death⁴⁰. The study also revealed that only one in four patients who died had been identified as in need of palliative care.
8. The PALLI tool was designed to be used to identify people with intellectual disabilities who may benefit from palliative care⁴¹. The tool was developed in the Netherlands using five-stage mixed methods design including retrospective survey, interviews, draft version, focus groups and finalisation for testing in practice. This tool, which consists of 39 questions, composed of eight main themes such as physical decline, changes in characteristic behaviour, and increases in symptom burden. The PALLI tool has been validated for use among patients with intellectual

disabilities in different settings, including primary care⁵⁰. PALLI tool shows promising construct validity and feasibility. There is, however, less and mixed evidence for the predictive validity of this tool⁵⁰.

9. Rainoe et al. used computerized electronic records to identify the most common factors associated with death within the next year among hospitalised patients²¹. A list of the identified factors (including age 75 and over and having diseases, such as heart failure and COPD) was used to identify people who may benefit from palliative care. The electronic tool has been validated against clinical assessment in an observational study in the USA, which found that 5.6% of primary care patients could benefit from palliative care²¹.
10. Electronic Frailty Index (eFI) was developed in the UK to identify elderly patients in primary care who may be living with frailty⁴⁸. The eFI uses a 'cumulative deficit' model to calculate a frailty score based on a range of deficits, which can be symptoms, signs, diseases and abnormal laboratory test values. The eFI has been used in two recent studies to identify people who are at increased risk of mortality and may need palliative care^{48, 49}. Initially, Stow et al. examined the ability of eFI to predict mortality by measuring it at a single time point, which found that 1.1% of individuals age 75 and over could benefit from palliative care⁴⁸. Stow et al. conducted another study using eFI to examine if changes in frailty index can be used to predict mortality and the need to palliative care⁴⁹. The study identified a distinct frailty trajectory which can be used to identify people who are at a higher risk of dying within 12 months. This study found that 0.49% of people age 75 and over were identified as potential candidates for palliative care. The predictive validity of eFI to identify patients with potential palliative care needs has been evaluated in both studies^{48, 49}.

Accuracy of screening tools

Eight studies reported accuracy data for five screening instruments (SPICT, SQ, NECPAL, eFI and early identification tool for palliative care patients "Rainoe tool")^{21, 35, 37, 45-49}. Reference standards (i.e. the comparator against which the tool was compared) varied across these studies, including 3-month mortality, 12-month mortality, 24-month mortality and clinical assessment. One study was excluded as data were available only on positively screened patients²⁸. Table 4 shows a summary of the sensitivity, specificity, positive predictive value, and negative predictive value for the screening tools.

Table 4: Summary of the sensitivity, specificity, positive predictive value, negative predictive value for the score

Reference	Length of Follow-Up	Comparison	Tool	Cut-off value	Reference standard	Final sample(n)	Age, median (range)
Mitchell, 2018	12 months	Intuition	SPICT (2012 version)	SQ+ with ≥ 2 general indicators or ≥ 1 clinical indicator)	h Mortality	1525	79.1, 81.3, 11.8)
			SQ	swer no to the 'surprise' question)	h Mortality		
Gómez-Batiste 2017	24 months	No	NECPAL	NECPAL+ (SQ+, and “ ≥ 1 general indicators or ≥ 1 specific indicators”)	12-months mortality	1059	81.3, 11.8)
					24-months mortality		
			SQ	SQ + (answer no to the 'surprise' question)	12-months mortality		
					24-months mortality		
Raincine, 2007	6-month (the length of the study)	No	Raincine tool	Patient is included if their electronic records contained at least one of the marker for high risk of death within the next year e.g. age >75 or a diagnosis of congestive heart failure.	Clinical assessment	18308	-
Barnes, 2008	12 months	No	SQ	SQ + (answer no to the 'surprise' question)	12-months mortality	231	77, m 71-82
Moroni, 2014	12-months	No	SQ	swer no to the 'surprise' question)	hs mortality	231	70.2
Lakin, 2016	12-months	No	SQ	swer no to the 'surprise' question)	hs mortality	1737	65, m
Stow, 2018,	12 months	No	eFI	with rapidly rising frailty (initial increase of 0.022 eFI per month before slowing from a baseline eFI of 0.21)	hs mortality	26,298	For ca mean For co mean
Stow, 2018,	3 months	No	eFI	alue > 0.19	s mortality	7890	For ca mean For co mean

SPICT= the Supportive & Palliative Care Indicators Tool; **NECPAL**= Necesidades Paliativas [Palliative Needs]; **SQ**= Surprise Question; **eFI** = electronic Frailty predictive value; **NPV**= negative predictive value; **+** = positive.

Across all screening tools and studies, only one study had a positive predictive value over 50% (83.8%). The negative predictive value was high for most tools and varied from 99% to 69%. The sensitivity and specificity values varied considerably and ranged from 3.2% to 94% and 26.4% to 99% respectively. Studies enrolling participants with advanced progressive diseases reported high sensitivity values; however, studies that targeted a general population of primary care (e.g. adults aged 70 and over) reported lower sensitivity values.

Methodological quality of studies which reported accuracy data for screening tools

The assessment of the risk of bias is summarized in Supplementary File S2. (RCTs), Supplementary File S3a. (cohort studies) and Supplementary File S3b. (case control studies). On the basis of the Newcastle–Ottawa scale, three of the five cohort studies were judged to bear a moderate risk of bias (fair quality)^{35, 46, 47} and one cohort was judged to have a high risk of bias (low quality) due to the lack of description of the follow-up and no adjustment for confounders²¹. Only one cohort study fulfilled most of the Newcastle-Ottawa scale criteria and had a low risk of bias (high quality)⁴⁵. Newcastle–Ottawa scale assessment revealed that the two case control study were all of a fair quality^{48, 49}. Based on the Cochrane risk of bias tool, the overall risk of bias for the included RCT was high because of unclear allocation concealment and differential drop-out rates between the two groups³⁷.

Discussion

Main findings

We identified 10 screening tools for identification of patients with advanced progressive diseases who are likely to have unmet palliative care needs in primary care which varied in content and accuracy, and in general, the validation studies were of low quality and with high risk of bias.

Most of the identified tools use either prediction of death or deterioration or both as proxies for the identification of people who are likely to have unmet palliative care needs. Patients with advanced progressive diseases experience different trajectories of decline and usually have varying needs at different phases in the illness trajectory^{51, 52}. Therefore, the identification process should not be based solely on predicting mortality or survival, but it should also focus on anticipating their needs whenever they occur, and predicting the rate and course of functional decline in order to trigger holistic assessment and make a proactive palliative care plan.

The proportion of patients identified with potential palliative care needs across studies ranged from 0.49% to 79%. The accuracy of five tools (of which data were available in 8 studies) showed sensitivity ranging from 3.2% to 94%, and specificity ranging from 26.4% to

99%. The wide variation in the accuracy of the screening tools may be caused by both variations in diagnostic groups and disease trajectory during the last year of life.

Strengths and weaknesses/limitations of the study

This is the first systematic review to assess the evidence on accuracy of screening tools for identification of patients with advanced progressive diseases who are likely to have unmet palliative care needs in primary care. We used a broad search strategy to identify all potentially relevant studies by searching Cochrane Library, MEDLINE, Embase and CINAHL, and we evaluated the quality of the validation studies using two reviewers independently with disagreements resolved by a third reviewer.

Our findings are limited by several issues. Firstly, our search strategy was designed to capture all of the relevant papers but given the nature of this topic, it is possible that some papers may have been missed. Although we conducted a comprehensive and broad search of the literature, we only included English language studies. We did not also include unpublished results or studies from the grey literature which may have introduced publication bias. However, the methodological quality of grey literature is usually lower than the quality of published studies literature^{53, 54}. Secondly, there is no current consensus about a reference standard against which the accuracy of a screening tool could be assessed. All studies used mortality as a reference standard, with the exception of one study that used clinical judgment to determine whether the identified patient could benefit from palliative care²¹. This is a major flaw in the evidence, in that we know palliative care needs do not relate particularly closely to time to death, especially for some illnesses such as organ failures. Data were universally missing on how many patients identified (or missed) by the screening tools actually had palliative care needs and so we cannot be certain of the true clinical value of these tools.

What this study adds

Improving identification of patients who are likely to have unmet palliative care needs is a crucial step to overcome inequity in access to palliative care and to ensure that patients receive the right care at the right time to meet their needs and preferences^{55 16}. Identification does not mean referral to specialist palliative care services is necessarily needed, but rather, it should trigger comprehensive and holistic assessment of palliative care needs of the identified patients and their families^{17, 56}.

Although some of the identified tools recommended some actions to be taken after the screening process, there is no clear or appropriate care pathway for people with advanced progressive diseases who have been identified as having potential palliative care needs. Based on the findings from this review, we created a conceptual graph to describe the

process of patient identification and assessment of palliative care needs (Figure 2). The first step in the process is using a screening tool to aid the identification of patients with advanced progressive diseases whose health is deteriorating and hence benefit from palliative care needs assessment. The screening tool should be based not solely on predicting mortality and deterioration but also on anticipating the needs whenever they occur and predicting the rate and course of functional decline. The identified patients who have potential palliative care needs could then be targeted for assessment to identify their unmet palliative care needs. The outcomes of the assessment can help to determine the level of care required and may prompt an introduction of a palliative care approach 'generalist palliative care' or referral to a specialist palliative care service.

Figure 2: The process of patient identification and assessment of palliative care needs

Primary care teams play a vital role in caring for people with advanced chronic diseases⁵⁷. One of the main challenges for them is to identify which of their patients might have unmet palliative care needs^{58, 59}. Implementing a systematic tool could help the primary care team to identify patients with advanced progressive diseases and potential palliative care needs. However, issues such as high workload and decreased resources and capacity in primary care can be barriers to implement such a screening tool⁴³. Therefore, we recommend the use of an electronic tool to systematically and automatically identify patients who might have unmet palliative care needs and trigger the use of a needs assessment tool. Although there are some electronic screening tools which have been used such as AnticiPal and Rainoe tools, their validity is unclear as they used the risk of deteriorating and dying as a proxy for the identification of people with potential palliative care needs^{21, 44, 57}.

The design of the future automated tools should be based on predicting functional decline and increasing needs as well as predicting mortality. Future studies of these tools should apply adequate reference standards such as palliative care interviews to examine whether the screening tools accurately identifies patients with potential palliative care needs^{56, 60}. The implementation and use of these tools within current clinical practice software required minimal resources and very little training and capacity which allow them to be used in busy primary care practices^{18, 61}. Implementation of validated and standardized screening tools would transform the identification process in primary care and improve timely access to palliative care for people with advanced progressive diseases and potential palliative care needs.

Conclusion

This systematic review identified 25 studies that reported the use or development of screening tools to identify patients who are likely to have unmet palliative care needs. The evaluation of these tools was limited because of a lack of a valid comparator and so their true clinical utility is unknown. Further research is needed to identify standardised screening

processes which are based not solely on predicting mortality and deterioration but also on anticipating a person's needs whenever they occur and predicting the rate and course of functional decline in order to trigger the use of a needs assessment tool to identify and address their unmet needs at the right time.

Author contributions

Y.E was involved in the conceptualisation, review of abstracts and articles, quality rating of studies and writing of the first draft. S.B. was involved in the review of abstracts and articles, data extraction and revising/drafting article. E.C. was involved quality rating of studies and revising/drafting article. L.Z. was involved in the review of abstracts and articles, data extraction and revising/drafting article. F.M, M.J. and M.B. were involved in conceptualization and revising/drafting article.

Acknowledgements

We thank Ms Natalie King, an Information Specialist in the Academic Unit of Health Economics of the University of Leeds, for her advice on designing the search strategy.

Declarations of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This work was supported by Yorkshire Cancer Research as part of the RESOLVE Programme Grant [grant number L412].

References

1. Brennan P, Perola M, van Ommen G-J, Riboli E and Consortium EC. Chronic disease research in Europe and the need for integrated population cohorts. *European journal of epidemiology*. 2017; 32: 741-9.
2. Divo MJ, Martinez CH and Mannino DM. **Ageing and the epidemiology of multimorbidity**. Eur Respiratory Soc, 2014.
3. Effiong A and Effiong AI. Palliative care for the management of chronic illness: a systematic review study protocol. *BMJ Open*. 2012; 2: e000899.

4. Moens K, Higginson IJ, Harding R, et al. Are there differences in the prevalence of palliative care-related problems in people living with advanced cancer and eight non-cancer conditions? A systematic review. *Journal of pain and symptom management*. 2014; 48: 660-77.
5. Chidiac C. The evidence of early specialist palliative care on patient and caregiver outcomes. *International Journal of Palliative Nursing*. 2018; 24: 230-7.
6. Currow DC, Allingham S, Bird S, et al. Referral patterns and proximity to palliative care inpatient services by level of socio-economic disadvantage. A national study using spatial analysis. *BMC Health Services Research*. 2012; 12: 424.
7. Temel JS, Greer JA, Muzikansky A, et al. Early Palliative Care for Patients with Metastatic Non–Small-Cell Lung Cancer. *New England Journal of Medicine*. 2010; 363: 733-42.
8. Vanbutsele G, Pardon K, Van Belle S, et al. Effect of early and systematic integration of palliative care in patients with advanced cancer: a randomised controlled trial. *The Lancet Oncology*. 2018; 19: 394-404.
9. Zimmermann C, Swami N, Krzyzanowska M, et al. Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial. *Lancet (London, England)*. 2014; 383: 1721-30.
10. Hui D, Hannon BL, Zimmermann C and Bruera E. Improving patient and caregiver outcomes in oncology: Team-based, timely, and targeted palliative care. *CA: A Cancer Journal for Clinicians*. 2018; 68: 356-76.
11. Gaertner J, Siemens W, Meerpohl JJ, et al. Effect of specialist palliative care services on quality of life in adults with advanced incurable illness in hospital, hospice, or community settings: systematic review and meta-analysis. *BMJ (Clinical research ed)*. 2017; 357: j2925.
12. Gomes B, Calanzani N, Curiale V, McCrone P and Higginson IJ. Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers. *Cochrane Database of Systematic Reviews*. 2013.
13. Kavalieratos D, Corbelli J, Zhang D, et al. Association between palliative care and patient and caregiver outcomes: a systematic review and meta-analysis. *Jama*. 2016; 316: 2104-14.
14. Allsop MJ, Ziegler LE and Mulvey MR. Duration and determinants of hospice-based specialist palliative care: A national retrospective cohort study. 2018; 32: 1322-33.
15. Funding the right care and support for everyone: creating a fair and transparent funding system. *Journal of pain & palliative care pharmacotherapy*. 2011; 25: 362-4.
16. Walsh RI, Mitchell G, Francis L and van Driel ML. What diagnostic tools exist for the early identification of palliative care patients in general practice? A systematic review. *Journal of Palliative care*. 2015; 31: 118-23.

17. George N, Phillips E, Zourova M, Song C, Lamba S and Grudzen C. Palliative care screening and assessment in the emergency department: a systematic review. *Journal of Pain and Symptom Management*. 2016; 51: 108-19. e2.
18. Downar J, Wegier P and Tanuseputro P. Early Identification of People Who Would Benefit From a Palliative Approach—Moving From Surprise to Routine. *JAMA network open*. 2019; 2: e1911146-e.
19. Maas EAT, Murray SA, Engels Y and Campbell C. What tools are available to identify patients with palliative care needs in primary care: a systematic literature review and survey of European practice. *BMJ supportive & palliative care*. 2013; 3: 444-51.
20. Hamano J, Oishi A and Kizawa Y. Identified palliative care approach needs with SPICT in family practice: A preliminary observational study. *Journal of Palliative Medicine*. 2018; 21: 992-8.
21. Rainone F, Blank A and Selwyn PA. The early identification of palliative care patients: preliminary processes and estimates from urban, family medicine practices. *American Journal of Hospice & Palliative Medicine*. 2007; 24: 137-40.
22. Murray SA, Boyd K, Sheikh A, Thomas K and Higginson IJ. Developing primary palliative care. British Medical Journal Publishing Group, 2004.
23. Maas EA, Murray SA, Engels Y and Campbell C. What tools are available to identify patients with palliative care needs in primary care: a systematic literature review and survey of European practice. *BMJ Support Palliat Care*. 2013; 3: 444-51.
24. R; R, consumers; C and group; cr. Communication Review Group. 'Cochrane Consumers and Communication Review Group: meta-analysis. 2013.
25. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. 2015; 4: 1.
26. Wells G. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non randomised studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. 2001.
27. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ (Clinical research ed)*. 2011; 343: d5928.
28. Scaccabarozzi G, Amodio E, Pellegrini G, et al. The "aRIANNA" Project: An Observational Study on a Model of Early Identification of Patients with Palliative Care Needs through the Integration between Primary Care and Italian Home Palliative Care Units. *Journal of Palliative Medicine*. 2018; 21: 631-7.
29. Clifford C, Thomas K and Armstrong Wilson J. Going for gold: the Gold Standards Framework programme and accreditation in primary care. *BMJ End Life J*. 2016; 6: e000028.
30. Afshar K, Feichtner A, Boyd K, et al. Systematic development and adjustment of the German version of the Supportive and Palliative Care Indicators Tool (SPICT-DE). *BMC Palliative Care*. 2018; 17: 27.

31. Hamano J, Oishi A and Kizawa Y. Prevalence and Characteristics of Patients Being at Risk of Deteriorating and Dying in Primary Care. *Journal of Pain and Symptom Management*. 2019; 57: 266-72.e1.
32. Fachado AA, Martínez NS, Roselló MM, et al. Spanish adaptation and validation of the supportive & palliative care indicators tool – SPICT-ES(TM). *Revista de Saúde Pública*. 2018; 52: 3.
33. Highet G, Crawford D, Murray SA and Boyd K. Development and evaluation of the Supportive and Palliative Care Indicators Tool (SPICT): a mixed-methods study. *BMJ supportive & palliative care*. 2014; 4: 285-90.
34. Gomez-Batiste X, Martinez-Munoz M, Blay C, et al. Prevalence and characteristics of patients with advanced chronic conditions in need of palliative care in the general population: a cross-sectional study. *Palliative Medicine*. 2014; 28: 302-11.
35. Gomez-Batiste X, Martinez-Munoz M, Blay C, et al. Utility of the NECPAL CCOMS-ICO tool and the Surprise Question as screening tools for early palliative care and to predict mortality in patients with advanced chronic conditions: A cohort study. *Palliative Medicine*. 2017; 31: 754-63.
36. Gomez-Batiste X, Martinez-Munoz M, Blay C, et al. Identifying patients with chronic conditions in need of palliative care in the general population: development of the NECPAL tool and preliminary prevalence rates in Catalonia. *BMJ Support Palliat Care*. 2013; 3: 300-8.
37. Mitchell GK, Senior HE, Rhee JJ, et al. Using intuition or a formal palliative care needs assessment screening process in general practice to predict death within 12 months: A randomised controlled trial. *Palliative Medicine*. 2018; 32: 384-94.
38. Thoonsen B, Engels Y, van Rijswijk E, et al. Early identification of palliative care patients in general practice: development of RADboud indicators for Palliative Care Needs (RADPAC). *British Journal of General Practice*. 2012; 62: e625-31.
39. Thoonsen B, Gerritzen SHM, Vissers KCP, et al. Training general practitioners contributes to the identification of palliative patients and to multidimensional care provision: secondary outcomes of an RCT. *BMJ Supportive & Palliative Care*. 2016: bmjpspcare-2015-001031.
40. Thoonsen B, Vissers K, Verhagen S, et al. Training general practitioners in early identification and anticipatory palliative care planning: a randomized controlled trial. *BMC Family Practice*. 2015; 16: 126.
41. Vrijmoeth C, Echteld MA, Assendelft P, et al. Development and applicability of a tool for identification of people with intellectual disabilities in need of palliative care (PALLI). *Journal of Applied Research in Intellectual Disabilities*. 2018; 31: 1122-32.
42. Vrijmoeth C, Groot CM, Christians MGM, et al. Feasibility and validity of a tool for identification of people with intellectual disabilities in need of palliative care (PALLI). *Research in developmental disabilities*. 2018; 72: 67-78.
43. Mason B, Boyd K, Murray SA, et al. Developing a computerised search to help UK General Practices identify more patients for palliative care planning: a feasibility study. *BMC Family Practice*. 2015; 16: 99.

44. Mason B, Boyd K, Steyn J, Kendall M, Macpherson S and Murray SA. Computer screening for palliative care needs in primary care: A mixed-methods study. *British Journal of General Practice*. 2018; 68: e360-e9.
45. Moroni M, Zocchi D, Bolognesi D, et al. The 'surprise' question in advanced cancer patients: A prospective study among general practitioners. *Palliative Medicine*. 2014; 28: 959-64.
46. Barnes S, Gott M, Payne S, et al. Predicting mortality among a general practice-based sample of older people with heart failure. *Chronic illness*. 2008; 4: 5-12.
47. Lakin JR, Robinson MG, Bernacki RE, et al. Estimating 1-Year Mortality for High-Risk Primary Care Patients Using the "Surprise" Question. *JAMA Intern Med*. 2016; 176: 1863-5.
48. Stow D, Matthews FE, Barclay S, et al. Evaluating frailty scores to predict mortality in older adults using data from population based electronic health records: case control study. *Age and Ageing*. 2018; 47: 564-9.
49. Stow D, Matthews FE and Hanratty B. Frailty trajectories to identify end of life: a longitudinal population-based study. *BMC medicine*. 2018; 16: 171.
50. Weijers F, Veldhoven C, Verhagen C, Vissers K and Engels Y. Adding a second surprise question triggers general practitioners to increase the thoroughness of palliative care planning: results of a pilot RCT with cage vignettes. *BMC palliative care*. 2018; 17: 64-.
51. Bausewein C, Booth S, Gysels M, Kühnbach R, Haberland B and Higginson IJ. Individual breathlessness trajectories do not match summary trajectories in advanced cancer and chronic obstructive pulmonary disease: results from a longitudinal study. *Palliative medicine*. 2010; 24: 777-86.
52. Campbell RT, Petrie MC, Jackson CE, et al. Which patients with heart failure should receive specialist palliative care? *European journal of heart failure*. 2018; 20: 1338-47.
53. Egger M, Juni P, Bartlett C, Holenstein F and Sterne J. How important are comprehensive literature searches and the assessment of trial quality in systematic reviews? Empirical study. *Health Technol Assess*. 2003; 7: 1-76.
54. Hopewell S, McDonald S, Clarke MJ and Egger M. Grey literature in meta-analyses of randomized trials of health care interventions. *Cochrane Database of Systematic Reviews*. 2007.
55. Primary Care Strategy and National Health Service (NHS) Contracts Group. 2019/20 General Medical Services (GMS) contract Quality and Outcomes Framework (QOF). National Health Service (NHS) England, 2019.
56. Ontario Palliative Care Network. Tools to Support Earlier Identification for Palliative Care. Canada: Ontario Palliative Care Network 2019.
57. Reynolds R, Dennis S, Hasan I, et al. A systematic review of chronic disease management interventions in primary care. *BMC family practice*. 2018; 19: 11.

58. Harrison N, Cavers D, Campbell C and Murray SA. Are UK primary care teams formally identifying patients for palliative care before they die? *Br J Gen Pract.* 2012; 62: e344-e52.
59. Marcucci FC, Cabrera MA, Perilla AB, et al. Identification and characteristics of patients with palliative care needs in Brazilian primary care. *BMC Palliative Care.* 2016; 15: 51.
60. Glare PA, Semple D, Stabler SM and Saltz LB. Palliative care in the outpatient oncology setting: evaluation of a practical set of referral criteria. *Journal of oncology practice.* 2011; 7: 366-70.
61. Ross J, Stevenson F, Dack C, et al. Developing an implementation strategy for a digital health intervention: an example in routine healthcare. *BMC health services research.* 2018; 18: 794.