

PHYSIOTHERAPY - AGE-RELATED CHANGES IN PHYSICAL FUNCTIONING: CORRELATES BETWEEN OBJECTIVE AND SELF-REPORTED OUTCOMES

Lisa Alcock^{1,2}, Thomas D. O'Brien^{3,4}, & Natalie Vanicek^{2,5}

¹ Institute of Neuroscience, Newcastle University Institute for Ageing, Newcastle University, UK

² Department of Sport, Health and Exercise Science, University of Hull, UK

³ Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, UK

⁴ School of Sport, Health and Exercise Science, Bangor University, UK

⁵ Discipline of Exercise and Sport Science, Faculty of Health Sciences, University of Sydney, AUSTRALIA

ABSTRACT

Objectives: Firstly, to quantify the variance attributable to age and estimate annual decline in physical function and self-reported health using a battery of outcome measures in healthy older females. Secondly, to determine whether self-reported functional losses are similar to those measured objectively and which best represent overall physical capacity.

Design: Experimental study, cross-sectional analysis.

Setting: Human Performance Laboratory, University setting.

Participants: Thirty-nine community-dwelling women (Mean[SD] age=71.5[7.3]years, range 60-83years) completed a battery of *objective* measures of function and a *self-reported* health status survey.

Intervention: None.

Main outcome measures: *Objective measures:* Gait speed; TUG test; sit-to-stand; concentric knee flexor and extensor moments; *self-reported:* the SF-36.

Results: Using a cross-sectional approach, annual declines were estimated for: TUG time (2.1%); gait speed (1.2%); knee extensor (2.2%) and flexor moments (3.0%); and *self-reported* Physical Functioning (0.9-1.2%) ($p \leq .001$). Linear regression indicated that age explained moderate variance in the *objective* ($R^2=21-34\%$) and *self-reported* ($R^2=14-28\%$) outcomes. TUG time and gait speed was significantly correlated with all *objective* outcomes except sit-to-stand ($r=0.46-0.83$) and most of the *self-reported* ($r=0.10-0.63$) outcomes ($p < .01$).

Conclusions: Age-related functional deterioration was estimated precisely across both objective and self-reported outcomes. Greater strength losses for the knee flexors compared to the extensors indicate an unequal strength loss of antagonistic muscle pairs which has implications for the safe completion of many functional tasks including obstacle negotiation, stair locomotion, postural transitions, and ultimately knee joint stability. Furthermore, walking speed and TUG time correlated most strongly with many of the outcomes highlighting their importance as global indicators of physical capacity.

Keywords: healthy ageing; objective and self-reported outcomes; physical capacity; functional performance

INTRODUCTION

The ability to perform common activities of daily living (ADL) autonomously is essential for independence and mobility. However, as we age, a combination of physical and psychological changes reduces our ability to complete these daily tasks(1-6). This may lead to increased sedentary behaviour, task avoidance and ultimately poorer quality of life(7). Government statistics have confirmed that adults ≥ 65 years have an increased life expectancy(8). Therefore, monitoring physical function within healthy older populations is critical to facilitate the early identification of decline and offer insight into the age-related loss of physical function and general health. Accordingly, the Chartered Society of Physiotherapy (CSP) has recommended that normative data for healthy older adults should be documented regularly as it provides a baseline from which age-related dysfunction may be identified(9). Furthermore, the Australian Physiotherapy Association have suggested that continued documentation is vital for monitoring, evaluating and justifying patient care(10).

An individual's ability to perform ADL can be monitored using a range of measured and self-reported outcomes(2, 3, 11). These assessments are often simple to perform, are appropriate for use with older adults(12) and individuals at risk of falling(13). Consequently, such "outcome

measures” are routinely used in physiotherapy, rehabilitation, occupational and geriatric settings(14) where they provide information about functional mobility(15), prospective falls risk(16), and offer moderate-to-good test-retest reliability(16, 17). Objectively measured functional losses (gait speed, chair-stand time, grip strength, and balance) correlate moderately with self-reported functioning (r ranged from $-.19$ to $.63$, $p < .05$)(18), and gait speed appears to be a global indicator of function over a broad range of capacities(18). Selecting the most appropriate battery of outcomes can be challenging for health professionals and is often dependent on the population of interest, the nature of the visit (i.e., routine check-up, hospital admission), and the equipment/expertise available.

Existing evidence originating from segmented age-group comparisons (ie. according to decade 60-69y, 70-79y etc) suggests that physical function worsens as older age ensues and that the decline is linear(2, 3, 5, 6), but the year-on-year progression of this deterioration is less defined. For example, the time taken to complete the ‘Timed get-Up and Go’ (TUG) test increases by ~40% (8.1–11.3s) between the ages of 60-99years(3) but the annual rate at which declines occur is less clear. Similar limitations in our knowledge exist with regards to the sit-to-stand (STS) task(4, 5). Comfortable gait speeds gradually slow with advancing age(19), and grouped means suggest that greater declines occur >79years(1). Lower limb strength decreases continuously across intervals of older age(20) with a 19% loss in knee extensor strength reportedly occurring across the 8th decade(21). Age-related changes in the SF-36 sub-scales indicate that, in general, the physical components deteriorate while the mental components remain constant or show small increases across age-grouped samples(22). However, these data do not include adults aged over 64 years, thus restricting generalisations to older populations. Very few studies have attempted to quantify annual losses in physical function, meaning subtle yet important changes may be overlooked when age-related decline is generalised from wide pre-defined sub-divisions of old age. Moreover, evaluating functional changes across broad age ranges makes it difficult to monitor functional decline across a shorter period of time and may mean that the optimal time to intervene could be missed inadvertently.

The few previous studies that have quantified annual changes have solely focused on the age-related change in functional performance in isolation such as muscle strength(21, 23), balance(24), measures of walking speed(25), and self-reported physical function only without substantiating objective measures(26). Moreover, TUG time has been used to estimate changes in a single(14) or a battery of outcomes (comfortable and fast gait speed, balance, TUG, STS, 6-minute walk time, and physical performance test)(1). Whilst the authors presented a range of measures, the study used both age and the use of an assistive device to understand age-related decline(1). Understanding the subtle nature of these losses that occur across the older age spectrum in a battery of outcomes will guide timely intervention to attenuate functional decline.

The aims of this study were to: (i) quantify annual changes in physical function in a convenience sample of healthy, older community-dwelling women across a battery of clinical outcomes; and (ii) to determine whether self-reported functional losses are similar to those measured objectively and which outcomes best represent overall physical capacity. It was hypothesised that linear relationships would exist between physical function and age with minimal annual (year-on-year) changes expected in self-reported well-being, throughout the age spectrum.

METHODS

PARTICIPANTS

Thirty-nine healthy, community-dwelling older women (Mean age[SD] 71.5[7.3] years, height 1.63[0.07]m, mass 70.6[12.4]kg) with no prior falls history gave written consent to participate in this study which was approved by the local NHS Ethics Committee (08/H1305/91). This work forms part of a larger set of studies that evaluated the influence of healthy ageing on the biomechanical profiles of ADL (27, 28). Due to technical failure during data acquisition, missing data were recorded for the STS test (n=37 remaining) and knee dynamometry (n=35 remaining).

OUTCOME MEASURES

The TUG and the STS tasks were assessed using the same standard chair with no arm rests (seat height:46cm, depth:38cm, and back height:74cm)(14, 15). A standard TUG test protocol was employed(29) whereby participants began seated, stood up, walked to and around a cone 3-metres away, and returned to a seated position. Participants were asked to complete the movement as quickly and safely as possible, refraining from using their arms for assistance during the chair rise, thus relying predominantly on the lower limbs for task completion. The movement was performed three times and the time to complete each trial was recorded using a stopwatch. To assess STS performance, participants began seated, and when ready stood up at their comfortable speed. Sagittal plane kinematics were measured at 100Hz (Qualisys, Sweden) to determine the time taken to complete one STS cycle. Movement initiation was defined as an increase in hip flexion of >1% of the maximum hip flexion during the movement (exhibited as increased forward trunk lean) that was shortly followed by knee extension. Maximum knee extension determined movement termination. Gait speed was derived from gait analysis data reported previously for the same sample(27). Briefly, 8-10 trials were collected while participants walked along a 10-metre walkway at a comfortable pace. Steady-state gait speed was obtained from the central part of the walkway and averaged across trials.

Knee flexor and extensor concentric strength were assessed bilaterally using dynamometry across the participants' full range of knee motion while the hip was flexed at 90° (Biodex System 3, Biodex Medical, Shirley, NY). Straps were secured around the trunk and hips for stability. The knee axis of rotation was aligned with the dynamometer axis of rotation and the dynamometer lever arm was secured to the distal end of the shank. Five practise trials were performed. Gravity corrected joint moments were recorded while participants performed maximal voluntary concentric contractions during five consecutive knee extension-flexion trials. Verbal encouragement was provided throughout. The angular speed of 180°/sec has been used previously with older adults(20) and concentric exercises have been shown to elicit significantly less cardiovascular stress than eccentric testing(30). Therefore, concentric testing at a high angular speed was chosen to

minimise cardiovascular stress and avoid potential injury(31). The SF-36 is regarded as a generic measure of health status(17). The survey is comprised of 36 questions covering both physical and mental health, each of which is composed of 4 sub-scales. Administration of each test was standardised to the following order: TUG test, STS, knee strength, gait speed and SF-36, as the first two assessments served as a whole body warm-up prior to strength testing.

DATA ANALYSIS

The fastest TUG time from the three trials obtained was selected for further analysis(32) permitting task familiarisation. The time taken to complete a single STS cycle comfortably was used. Knee moments were normalised to body weight (Nm/kg). The hamstrings-to-quadriceps (H:Q) ratio was calculated from the peak joint moments. Paired-samples t-tests indicated that no significant strength differences existed between the right and left limbs so were combined for all analyses.

The SF-36 was analysed according to the 8 sub-scales: Physical Functioning, Role limitations due to physical health problems (Role–Physical), Bodily Pain, General Health, Vitality, Social Functioning, Role limitations due to emotional problems (Role–Emotional), and Mental Health(33). In the event of missing data, scores were estimated per participant by averaging the answers given within the section of questions with missing data(33). Of the 1404 questions (36 questions, 39 participants) only 30 data points were missing for nine participants. For seven participants, missing data were estimated and two participants were excluded from further analysis (n=37 remaining). Linear transformations were computed to transform SF-36 scores into z-scores using the norm-based scoring procedure according to the currently available normative database of the 1998 general US population(34). Each of the sub-scales was aggregated using a T-score transformation to produce a Mental Component Summary (MCS) and a Physical Component Summary (PCS).

Each of the outcome measures presented were chosen for their reliability and suitability for use with older adults. Intra-class correlation coefficients(ICC) indicate how consistent or reproducible

a quantitative measurement is. For example, the TUG time offers high reproducibility with an ICC of 0.8 from a cohort of older people(35) and similar levels of consistency has been confirmed for STS(6). Mechanical reliability of the dynamometer is high(ICC=0.99) as is test re-test reliability of strength in older adults(ICC>0.92)(36). Respectable ICCs have also been presented for the gait speed of older adults(ICC=0.74)(5) and across the age continuum incorporating both young and old(ICC=~0.9)(2). Finally, the SF-36 is suitable for use with older populations and demonstrates good construct validity with Cronbach's alpha statistics ranging from 0.82 to 0.94 for each of the subscales(17).

STATISTICAL ANALYSIS

Aim 1 – Estimation of annual changes in health and physical function

Bivariate correlations (Pearson's r) expressed the strength of the relationship between each of the outcomes and age. Linear regression was computed for all outcomes using age as a single independent predictor. Dependent variables were plotted against age to check for linearity and statistical assumptions surrounding regressive procedures were confirmed. Outliers were determined from the standardised residuals and data $\pm 3SD$ from the mean were considered to be extreme and unlikely to have occurred by chance therefore they were removed(37, 38). For completeness, models excluding outliers are presented within the table and models with outliers included are presented within the footnotes of the table. R^2 (%) and the standardised regression coefficient (beta) are presented. Using the 75th percentile as a starting value, the annual change in function was calculated. 95% confidence intervals (95%CI) assessed the variation in estimated annual change.

Aim 2 – Changes in objective and self-reported physical function

Additional bivariate correlations were calculated between each of the objective and self-reported outcomes. Significance was accepted when $p \leq .05$.

RESULTS

Aim 1 - Estimation of annual changes in health and physical function

Five of the participants (aged 75, 78, 80 and two 83year olds) were unable to achieve the constant target velocity of 180°/s during the knee flexion dynamometry, and thus data for these individuals were removed from further analysis of flexion moments and H:Q ratio only. All of the objective assessments, with the exception of the STS, were significantly correlated with age ($p \leq .01$, Table 1, Figure 1). All significant relationships, except for TUG time, were negatively correlated with age ($r = -.46$ - $-.58$, $p \leq .05$). Strong negative correlations existed between age and knee strength measures (peak moments and H:Q ratio) and gait speed ($r = -.46$ - $-.58$, $p \leq .05$). The strongest negative correlations were found between age and knee moments ($r = -.58$, $p \leq .01$). Only the Physical Functioning sub-scale and PCS of the SF-36 were correlated with age ($r = -.53$ and $-.38$, respectively, Figure 2). Age contributed to explaining moderate-to-good levels of variance in all objective outcomes, except for STS time ($R^2 = 21-34\%$, Table 1). Significant point estimates (B) were found for all of the objective outcomes, except for STS, revealing annual changes of 1.2-3.0%/year. Age explained the greatest variance in the knee extensor ($R^2 = 33\%$) and flexor ($R^2 = 34\%$) moments.

Age contributed to explaining moderate-to-good levels of variance in the Physical Functioning sub-scale and PCS ($R^2 = 28\%$ and 14% , respectively) with self-reported losses of 1.2%/year and 0.9%/year, respectively. Knee flexor moments exhibited the greatest decline (3.0%/year), although 95%CI were wider for the knee strength measures compared to self-rated Physical Functioning ($\pm 1.3\%$) and the PCS ($\pm 1.5\%$). Regression analysis could not be computed with a number of SF-36 subscales, as the majority of the sample attained the maximum score (100) demonstrating that a ceiling effect had occurred: Role-Physical and Social Functioning (both $n = 29$, 74% of the sample) and Role-Emotional ($n = 32$, 82% of the sample).

1 **TABLE 1** Regression models developed to explain the variance in the outcome measures explained by advancing age

		MEAN	Correlation with Age (r)	R ² (%)	REGRESSION COEFFICIENT		95% CI	ANNUAL CHANGE (%/year)	95% CI ANNUAL CHANGE
		(SD)			beta	Sig.			
FUNCTIONAL MEASURES	TUG time (s) [1]	7.85 (2.9)	.53	28%	.181	p = .001	.084 : .277	+2.1%	1.0 : 3.3%
	STS time (s)	1.46 (0.26)	.15	0%	-.085	.386			
	Gait speed (m/s)	1.28 (0.20)	-.57	32%	-.017	p ≤ .001	-.026 : -.009	-1.2%	-1.9 : -0.6%
	HQ Ratio (%)	0.40 (0.15)	-.46	21%	-.010	p ≤ .001	-.017 : -.003	-2.0%	-3.4 : -0.6%
	Peak Knee Extensor Moment (Nm/kg)	0.48 (0.18)	-.58	33%	-.014	p ≤ .001	-.021 : -.007	-2.2%	-3.5 : -1.0%
	Peak Knee Flexor Moment (Nm/kg)	0.21 (0.11)	-.58	34%	-.009	p ≤ .001	-.014 : -.004	-3.0%	-4.7 : -1.3%
SELF-REPORTED MEASURES	SF36: Physical Functioning	81.7 (15.5)	-.53	28%	-1.143	p ≤ .001	-1.762 : -.525	-1.2%	-1.9 : -0.6%
	SF36: Role- Physical	84.2 (31.0)	-.19	^	^	^			
	SF36: Bodily Pain	77.4 (17.4)	-.30	9%	-.835	.065			
	SF36: General Health	73.4 (15.8)	-.21	10%	-.685	.059			
	SF36: Vitality	70.1 (16.9)	-.29	9%	-.770	.074			
	SF36: Social Functioning	93.8 (12.9)	-.03	^	^	^			
	SF36: Role – Emotional	92.1 (21.1)	-.11	^	^	^			
	SF36: Mental Health [2]	84.1 (10.3)	-.06	0%	-.105	.701			
	SF36: Mental Component Summary [3]	56.6 (5.1)	.01	0%	.008	.958			
	SF36: Physical Component Summary	50.3 (8.5)	-.38	14%	-.443	.022	-.801 : -.065	-0.9%	-1.6 : -0.1%

2 ^ indicates variables where regression could not be computed due to ceiling effects, CI denotes confidence interval, HQ: Hamstrings-to-Quadriceps Ratio.

3

4 Footnotes indicate the models with outliers included within the model


5 [1] Regression model: Age, TUG time - R² = 29%, Beta coefficient = 0.121, p = .121

6 [2] Regression model: Age, Mental health: - R² = 4%, Beta coefficient = 0.270, p = .267

7 [3] Regression model: Age, MCS: R² = 1%, B = -0.086, p = .524

8 *Aim 2 – Changes in objective and self-reported physical function*


9 Cross-correlations between the outcomes revealed that TUG time strongly correlated with knee
10 strength ($r=-.55$ - $-.68, p \leq .01$; Table 2) and was highly correlated with speed ($r= -.83, p \leq .01$). TUG
11 time was negatively correlated with many of the physical SF-36 sub-scales and the PCS ($r \geq -$
12 $.40, p \leq .05$). STS correlated well with TUG time ($r=.38, p \leq .05$) and gait speed ($r=-.35, p \leq .05$) but was
13 not correlated with knee strength. Like TUG time and gait speed, STS time was moderately
14 correlated with many of the SF-36 subscales ($r=-.41$ - $-.54, p \leq .01$). Gait speed was positively
15 correlated with many of the SF-36 variables ($r=.41-.63, p \leq .05$) including all of the physical
16 components.



17

18 **FIGURE 1** Relationships (r) between age and the functional outcome measures studied

19 For completeness graphs are presented with outliers, ** $p \leq .01$ (1-tailed), and *** $p \leq .001$ (1-tailed)



20

21 **FIGURE 2** Relationships (*r*) between age and the eight (physical and mental) sub-scales of the SF-

22 36

23 For completeness graphs are presented with outliers, * $p \leq .05$

24

25 **TABLE 2** Correlation matrix (r) between all outcome measures studied

		TUG time (s)	STS time (s)	Gait speed (m/s)	HQ ratio (%)	Peak Knee Extensor Moment (Nm/kg)	Peak Knee Flexor Moment (Nm/kg)	SF-36									
								PF	R-P	BP	GH	VT	SF	R-E	MH	MCS	PCS
FUNCTIONAL MEASURES	TUG time (s)		.38	-.83	-.55	-.57	-.68	-.58	-.27	-.40	-.43	-.49	-.30	-.40	-.29	-.31	-.47
	STS time (s)	-		-.35	-.07	-.18	-.12	-.46	-.34	-.43	-.28	-.46	-.54	-.45	-.50	-.50	-.41
	Gait speed (m/s)	-	-		.46	.67	.67	.63	.33	.44	.41	.49	.25	.41	.27	.27	.52
	HQ ratio (%)	-	-	-		.30	.82	.44	.26	.32	.46	.35	.03	-.08	.12	-.04	.45
	Peak Knee Extensor Moment (Nm/kg)	-	-	-	-		.76	.35	.07	.15	.09	.23	.07	.02	.03	.00	.21
	Peak Knee Flexor Moment (Nm/kg)	-	-	-	-	-		.53	.18	.35	.48	.31	.15	-.12	.01	-.14	.47
SELF-REPORTED MEASURES	SF36: Physical Functioning	-	-	-	-	-	-		.71	.74	.55	.55	.34	.60	.10	.12	.90
	SF36: Role – Physical	-	-	-	-	-	-	-		.69	.65	.68	.45	.63	.20	.29	.90
	SF36: Bodily Pain	-	-	-	-	-	-	-	-		.46	.55	.57	.64	.30	.34	.84
	SF36: General Health	-	-	-	-	-	-	-	-	-		.69	.28	.29	.33	.27	.71
	SF36: Vitality	-	-	-	-	-	-	-	-	-	-		.51	.40	.57	.59	.65
	SF36: Social Functioning	-	-	-	-	-	-	-	-	-	-	-		.74	.57	.84	.42
	SF36: Role – Emotional	-	-	-	-	-	-	-	-	-	-	-	-		.22	.59	.64
	SF36: Mental Health	-	-	-	-	-	-	-	-	-	-	-	-	-		.84	.11
	SF36: Mental Component Summary	-	-	-	-	-	-	-	-	-	-	-	-	-	-		.17
	SF36: Physical Component Summary	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

26

27 HQ: Hamstrings-to-Quadriceps Ratio, PF: Physical Functioning, R-P: Role-Physical, BP: Bodily Pain, GH: General Health, VT: Vitality, SF: Social Functioning, R-E: Role-Emotional, MH: Mental Health,

28 Light grey shading indicates moderate correlations $r = 0.4-0.6$ $p < .05$, and dark grey shading indicates strong correlations $r = 0.7-1.0$ $p < .05$.

30 **DISCUSSION**

31 This study used linear regression to quantify the changes in objective and self-reported function
32 that were attributable to age alone. This is a novel approach that furthers our understanding
33 beyond that of previous studies which have assessed functional loss using a single clinical
34 assessment (2, 3, 21, 24, 25, 39) or a battery of assessments according to broad pre-defined age
35 groupings (1, 3, 40-42). This study found that the pattern of age-related functional losses can be
36 estimated using linear regression, thereby enhancing our understanding of the rate of decline
37 throughout older age. Furthermore, large within-group variability (represented by the standard
38 deviation) was observed for many of the outcomes presented, indicating that categorising
39 individuals across large age ranges, as has been common in previous research(43, 44), leads to
40 considerable variation and limited information about age-related functional decline.

41
42 *Aim 1 - Estimation of annual changes in health and physical function*

43 Overlapping 95%CI for estimated annual changes in all of the objective outcomes (except for STS)
44 and the Physical Functioning and Physical Component Summary of the SF-36 (Table 1) indicate
45 that gradual losses due to biological ageing processes appear to occur at similar rates for which
46 age has been shown to explain significant variation ($R^2=14-34\%$). This is in agreement with
47 previous research demonstrating that age was a major contributor, explaining significant variation
48 in TUG time ($R^2=60\%$), gait speed ($R^2=63\%$) and timed STS ($R^2=37\%$)(1). While the convenience
49 sample in that study(1) showed gradual decline over a large age range (66-101years), both men
50 and women were tested which may have affected the point estimates due to gender differences in
51 musculoskeletal function/capacity(5, 6) and relative life expectancy(8). Our study has
52 demonstrated that significant annual losses in function occur within a cohort of healthy older
53 women and further analyses should be repeated with men for comparison. To our knowledge,
54 Lusardi et al.'s study(1) is the only study that has taken a similar approach to understanding age-
55 related decline in a battery of motor function indices. Other studies have quantified losses in a

56 similar fashion, for musculoskeletal strength (45) but have not investigated relative annual decline
57 for a range of functional characteristics.

58

59 It is noteworthy that greater age-induced losses were observed for the knee flexors compared with
60 the extensors, and this is in agreement with longitudinal observations of older women(45). Those
61 findings, along with our present data, indicate that the smaller flexor muscles experience greater
62 strength losses compared to the larger anti-gravity extensor group. Accordingly, the point
63 estimates presented in this study indicate that the hamstrings-to-quadriceps (H:Q) ratio reduced
64 with age, with an annual loss of 2.0%. This suggests that the strength difference between this
65 antagonistic pair became larger with age. Such declines may impact adversely on the ability to
66 negotiate obstacles and stairs safely; execute postural transitions; bend down to a lower level; and
67 ultimately knee joint stability. It should be noted that the knee flexion moments of five of the oldest
68 participants were excluded from all analyses because they were unable to achieve the constant
69 target velocity (isovelocity phase). This was despite multiple trials and verbal encouragement, and
70 was most likely because they lacked muscle power to accelerate the dynamometer lever arm
71 adequately. Consequently, the reported knee flexion moments for participants aged >75 years are
72 likely overestimated compared to the population and the annual decline for knee flexion moment
73 and H:Q ratio are likely underestimated. We consider this to further demonstrate the significant
74 age-related weakness in the oldest old (particularly for knee flexor tasks). However, even at the
75 magnitudes observed considerable losses were identified that in accumulation could restrict
76 mobility and independent functioning. The expansion of this work to other antagonistic muscle
77 pairs (hip and ankle) may help to define joint functionality and inform the nature of age-related
78 reciprocal muscle strength loss. While the accumulated annual changes in knee strength with age
79 are of clinical significance, it must be highlighted that in some cases, cross-sectional studies have
80 reportedly underestimated strength losses compared to longitudinal research(45).

81

82 Interestingly, age explained moderate levels of variance in Physical Functioning and the PCS,
83 which was of a similar magnitude to the measured physical decline observed in gait speed, TUG

84 time and knee strength. Age did not explain significant variance in any other SF-36 sub-scale
85 suggesting that some aspects of the SF-36 may reveal changes related to other factors such as
86 personality type, or demonstrate inconsistencies due to individual perception. A self-reporting
87 survey results in perceived ability and/or status rather than actual task performance, assessed
88 objectively. Although self-reporting general health scales have been suggested to predict functional
89 decline accurately(46), the under- or over-reporting of physical status has been linked to
90 personality traits and levels of self-awareness(47). This study has shown that, in a sample of
91 healthy older women with no prior falls history, self-reported function estimated measured
92 performance very accurately. We found the same annual loss for gait speed and for the Physical
93 Functioning sub-scale of the SF-36 (1.2%/year). 95%CI overlapped for these variables and were
94 small, indicating that within this cohort the changes were predicted very precisely. This was likely
95 indicative of the participants' good health and independent living status.

96

97 In summary, accumulative losses in function have been demonstrated in the current study. The
98 year-on-year changes in function presented may appear to be small and thus not clinically
99 significant. However, when considering the accumulation of these changes across several years
100 and in relation to critical thresholds proposed within the literature (For example, gait speeds of
101 <1.0m/s(48) and TUG times of >14s(13)) the optimal time to intervene may be planned to
102 attenuate losses before overall function is affected clinically. Larger population-based studies are
103 required to substantiate the declines presented in this study and then national databases may be
104 developed and used to monitor functional/ global health changes to estimate demographic health
105 and future costs.

106

107 *Aim 2 – Changes in objective and self-reported physical function*

108 Using gait speed as a clinical outcome is advantageous due to its high test-retest reliability(2, 5),
109 as well as the significant correlations other outcomes(1) and deterioration with age(2). This study

110 found that that in addition to speed, TUG time reflects the functional declines in many ADL. In fact,
111 utilising the TUG test in clinical practice may be advantageous due to the limited space/ equipment
112 and expertise required compared with the space required to attain, and record speed from, steady-
113 state gait. However, both of these objective outcomes should be included routinely as part of a
114 normal health check-up and physical assessment of older women. Reduced performance on these
115 outcomes may be caused by declining knee extensor strength which contributes heavily to both
116 measures (lower limb stability and forward continuance during level gait; and raising the body from
117 a seated position). Measured leg strength is not currently included on the list of validated measures
118 provided by the CSP. This is likely due to the dynamometry equipment and technical expertise
119 required for data acquisition. A viable compromise may include quantifying muscle strength
120 manually given its widespread use within a clinical setting(49, 50). Reliable, valid data may be
121 obtained from such measures(51, 52) providing that recommendations such as those proposed by
122 the International College of Applied Kinesiology are adhered to(53). However, extra care must be
123 taken when constructing controlled testing conditions for use with large age ranges given that
124 some of the oldest individuals appear unable to perform at higher joint angular velocities.
125 Furthermore, slower gait speeds may result from age-associated muscle weakness(54), and more
126 specifically knee extensor strength loss (as indicated by the findings in this study). Therefore,
127 reduced functioning when standing from sitting (TUG) combined with deteriorating gait speeds,
128 both relatively simple measures to obtain, may provide an early indication of knee extensor
129 weakness which consequently will adversely affect overall physical function. To this end, measures
130 of gait speed and TUG time proved to be informative markers indicative of functional decline,
131 related most strongly to performance across a range of ADLs, and thus should continue to be
132 collected as part of routine assessment in outpatients, clinics and physiotherapy/ rehabilitation.

133

134 Measuring STS performance did not reveal functional losses above and beyond that of comfortable
135 gait speed, TUG time and knee strength. Thus, performing the STS is unlikely to further enhance
136 our understanding of annual musculoskeletal changes in performance when used in combination
137 with other outcomes. Rarely, do we perform a single postural transition in isolation and STS time

138 may be criticised for its lack of ecological validity. Moderate correlations were observed between
139 STS time and some of the psychological subscales of the SF-36 and the role of psychological
140 status and STS time has been confirmed previously(55).

141

142 The results of this study emphasise that there was a continuous decline in function throughout
143 older age. These musculoskeletal losses occurred even in a sample of healthy older women, for
144 whom functional decline appeared to be linear. Gait speed and TUG time were reaffirmed as
145 markers indicative of functional decline and related most strongly to performance across a range of
146 ADLs. Considering their cost- and time-effectiveness and strong correlation with many other
147 outcomes, gait speed and TUG should be incorporated into routine clinical and geriatric
148 assessments starting from middle-age as declines occur in even healthy older women beyond 60
149 years of age. The significant correlations between H:Q ratio and TUG time ($r=-.55$) and gait speed
150 ($r=.46$) further substantiate their suitability for clinicians to obtain during routine assessments with
151 older adults. However, because knee flexor strength does not contribute to TUG time and gait
152 speed as much as extensor strength, additional assessments may be required to identify flexor
153 strength and H:Q ratio losses. The predicted yearly decline in knee flexor strength, which was
154 greater than for the knee extensors, has important implications for the safe execution of many
155 functional tasks which may not be otherwise highlighted by losses in gait speed and TUG time
156 alone. Consequently, this should be addressed in exercise interventions for older adults, as the
157 knee flexors may be 'neglected' in primarily anti-gravity based exercises.

* Ethical Approval: This study was approved by the local NHS Ethics Committee (Reference - 08/H1305/91).

* Funding: This study received no external funding.

* Conflict of Interest: The authors would like to disclose that there were no financial or personal motivations that may have influenced the research findings of this study.

REFERENCES

1. Lusardi MM, Pellecchia GL, Schulman M. Functional performance in community living older adults. *Journal of Geriatric Physical Therapy*. 2003;26(3):14.
2. Bohannon R. Comfortable and maximum walking speed of adults aged 20-79 years: reference values and determinants. *Age and Ageing*. 1997;26(1):15-9.
3. Bohannon RW. Reference values for the timed up and go test: a descriptive meta-analysis. *Journal of Geriatric Physical Therapy*. 2006;29(2):64-8.
4. Chen H, Lin C, Yu L. Normative Physical Fitness Scores for Community-Dwelling Older Adults. *Journal of Nursing Research*. 2009;17(1):30-41.
5. Butler AA, Menant JC, Tiedemann AC, Lord SR. Age and gender differences in seven tests of functional mobility. *Journal of NeuroEngineering and Rehabilitation*. 2009;30(6):31-9.
6. Jones C, Rikli R, Beam W. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Research quarterly for exercise and sport*. 1999;70(2):113-9.
7. Newsom JT, Schulz R. Social support as a mediator in the relation between functional status and quality of life in older adults. *Psychology and aging*. 1996;11(1):34-44.
8. Department for Work and Pensions. Opportunity Age Indicators: 2008 Update 2008 [9th October 2012]. Available from: www.dwp.gov.uk/docs/indicators-update-2008.pdf.
9. The Chartered Society of Physiotherapy. Priorities for Physiotherapy Research in the UK: Topics prioritised by the older people expert panel - Annex S. No. 5. 2002.
10. Australian Physiotherapy Association. Standards for Physiotherapy Practises. Melbourne 2011.
11. Garratt AM, Ruta DA, Abdalla MI, Buckingham JK, Russell IT. The SF36 health survey questionnaire: an outcome measure suitable for routine use within the NHS? *British Medical Journal*. 1993;306(6890):1440-4.
12. de Vries NM, Staal JB, van Ravensberg CD, Hobbelen JSM, Olde Rikkert MGM, Nijhuis-van der Sanden MWG. Outcome instruments to measure frailty: A systematic review. *Ageing Research Reviews*. 2011;10(1):104-14.

13. Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Physical Therapy*. 2000;80(9):896-903.
14. Freter SH, Fruchter N. Relationship between timed 'up and go' and gait time in an elderly orthopaedic rehabilitation population. *Clinical Rehabilitation*. 2000;14(1):96-101.
15. Bischoff HA, Stahelin HB, Monsch AU, Iversen MD, Weyh A, von Dechend M, et al. Identifying a cut-off point for normal mobility: a comparison of the timed 'up and go' test in community-dwelling and institutionalised elderly women. *Age and Ageing*. 2003;32(3):315-20.
16. Yelnik A, Bonan I. Clinical tools for assessing balance disorders. *Clinical Neurophysiology*. 2008;38(6):439-45.
17. Lyons RA, Perry HM, Littlepage BN. Evidence for the validity of the Short-form 36 Questionnaire (SF-36) in an elderly population. *Age and Ageing* 1994;23(3):182-4.
18. Cress ME, Schechtman KB, Mulrow CD, Fiatarone MA. Relationship between physical performance and self-perceived physical function. *Journal of the American Geriatrics Society*. 1995.
19. Dobbs RJ, Charlett A, Bowes SG, Oneill CJA, Weller C, Hughes J, et al. Is this walk normal? *Age and Ageing*. 1993;22(1):27-30.
20. Newman AB, Haggerty CL, Goodpaster B, Harris T, Kritchevsky S, Nevitt M, et al. Strength and muscle quality in a well-functioning cohort of older adults: the Health, Aging and Body Composition Study. *Journal of the American Geriatrics Society*. 2003;51(3):323-30.
21. Goodpaster BH, Carlson CL, Visser M, Kelley DE, Scherzinger A, Harris TB, et al. Attenuation of skeletal muscle and strength in the elderly: The Health ABC Study. *Journal of Applied Physiology*. 2001;90(6):2157-65.
22. Jenkinson C, Coulter A, Wright L. Short form 36 (SF36) health survey questionnaire: normative data for adults of working age. *British Medical Journal*. 1993;306(6890):1437-40.
23. Goodpaster BH, Park SW, Harris TB, Kritchevsky SB, Nevitt M, Schwartz AV, et al. The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2006;61(10):1059-64.

24. Vereeck L, Wuyts F, Truijen S, Van de Heyning P. Clinical assessment of balance: normative data, and gender and age effects. *International journal of audiology*. 2008;47(2):67-75.
25. Enright PL, Sherrill DL. Reference equations for the six-minute walk in healthy adults. *American Journal of Respiratory and Critical Care Medicine*. 1998;158(5):1384-7.
26. Beckett LA, Brock DB, Lemke JH, de Leon CFM, Guralnik JM, Fillenbaum GG, et al. Analysis of change in self-reported physical function among older persons in four population studies. *American Journal of Epidemiology*. 1996;143(8):766-78.
27. Alcock L, Vanicek N, O'Brien TD. Alterations in gait speed and age do not fully explain the changes in gait mechanics associated with healthy older women. *Gait Posture*. 2013;37(4):586-92.
28. Alcock L, O'Brien TD, Vanicek N. Biomechanical demands differentiate transitioning vs. continuous stair ascent gait in older women. *Clinical Biomechanics*. 2014;29(1):111-8.
29. Podsiadlo D, Richardson S. The timed" Up & Go": a test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society*. 1991;39(2):142-8.
30. Overend TJ, Versteegh TH, Thompson E, Birmingham TB, Vandervoort AA. Cardiovascular stress associated with concentric and eccentric isokinetic exercise in young and older adults. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2000;55(4):B177-B82.
31. Kirkendall DT, Garrett Jr WE. The effects of aging and training on skeletal muscle. *The American Journal of Sports Medicine*. 1998;26(4):598-602.
32. Samson MM, Meeuwssen I, Crowe A, Dessens J, Duursma SA, Verhaar H. Relationships between physical performance measures, age, height and body weight in healthy adults. *Age and Ageing*. 2000;29(3):235-42.
33. Ware JE, Snow KK, Kosinski M, Gandek B. SF-36 health survey: manual and interpretation guide: The Health Institute, New England Medical Center; 1993.
34. Ware JE, Kosinski M. SF-36 Physical and Mental Health Summary Scale: A manual for users of Version 1: QualityMetric; 2001.

35. Bohannon RW, Schaubert K. Long-term reliability of the timed up-and-go test among community-dwelling elders. *Journal of Physical Therapy Science*. 2005;17(2):93-6.
36. Schaubert KL, Bohannon RW. Reliability and validity of three strength measures obtained from community-dwelling elderly persons. *The Journal of Strength & Conditioning Research*. 2005;19(3):717-20.
37. Knorr EM. *Outliers and data mining: finding exceptions in data*: The University of British Columbia; 2002.
38. Freedman D, Pisani, R., Purves, R.,. *Statistics*. New York: W.W. Norton; 1978.
39. Lindle R, Metter E, Lynch N, Fleg J, Fozard J, Tobin J, et al. Age and gender comparisons of muscle strength in 654 women and men aged 20–93 yr. *Journal of Applied Physiology*. 1997;83(5):1581-7.
40. Oberg T, Karsznia A, Oberg K. Basic gait parameters: reference data for normal subjects, 10-79 years of age. *Journal of Rehabilitation Research and Development*. 1993;30(2):210-23.
41. Oberg T, Karsznia A, Oberg K. Joint angle parameters in gait: reference data for normal subjects, 10-79 years of age. *Journal of Rehabilitation Research and Development*. 1994;31(3):199-213.
42. Cohen H, Heaton LG, Congdon SL, Jenkins HA. Changes in sensory organization test scores with age. *Age and Ageing* 1996;25(1):39-44.
43. Brach JS, Studenski S, Perera S, VanSwearingen JM, Newman AB. Stance time and step width variability have unique contributing impairments in older persons. *Gait & Posture*. 2008;27(3):431-9.
44. Arnadottir SA, Mercer VS. Effects of footwear on measurements of balance and gait in women between the ages of 65 and 93 years. *Physical Therapy*. 2000;80(1):17-27.
45. Hughes VA, Frontera WR, Wood M, Evans WJ, Dallal GE, Roubenoff R, et al. Longitudinal muscle strength changes in older adults. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2001;56(5):B209-B17.
46. Lee Y. The predictive value of self assessed general, physical, and mental health on functional decline and mortality in older adults. *Journal of Epidemiology and Community Health*. 2000;54(2):123-9.

47. Suchy Y, Kraybill ML, Franchow E. Instrumental activities of daily living among community-dwelling older adults: Discrepancies between self-report and performance are mediated by cognitive reserve. *Journal of Clinical and Experimental Neuropsychology*. 2011;33(1):92-100.
48. Cesari M, Kritchevsky SB, Penninx BWHJ, Nicklas BJ, Simonsick EM, Newman AB, et al. Prognostic Value of Usual Gait Speed in Well - Functioning Older People—Results from the Health, Aging and Body Composition Study. *Journal of the American Geriatrics Society*. 2005;53(10):1675-80.
49. Conable KM, Rosner AL. A narrative review of manual muscle testing and implications for muscle testing research. *Journal of chiropractic medicine*. 2011;10(3):157-65.
50. Cuthbert SC, Goodheart GJ. On the reliability and validity of manual muscle testing: a literature review. *Chiropractic & Manual Therapies*. 2007;15(1):4.
51. Perry J, Weiss WB, Burnfield JM, Gronley JK. The supine hip extensor manual muscle test: a reliability and validity study. *Archives of physical medicine and rehabilitation*. 2004;85(8):1345-50.
52. Escolar DM, Henricson EK, Mayhew J, Florence J, Leshner R, Patel KM, et al. Clinical evaluator reliability for quantitative and manual muscle testing measures of strength in children. *Muscle & nerve*. 2001;24(6):787-93.
53. International College of Applied Kinesiology. Applied Kinesiology Status Statement: Executive Board of the International College of Applied Kinesiology-U.S.A; 1992 [25th February 2014]. Available from: http://www.icak.com/index.php?option=com_content&view=article&id=54&Itemid=65.
54. Frontera WR, Hughes VA, Lutz KJ, Evans WJ. A cross-sectional study of muscle strength and mass in 45- to 78-yr-old men and women. *Journal of Applied Physiology*. 1991;71(2):644-50.
55. Lord SR, Murray SM, Chapman K, Munro B, Tiedemann A. Sit-to-stand performance depends on sensation, speed, balance, and psychological status in addition to strength in older people. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2002;57(8):M539-M43.