

## ALUMINIUM-26 FROM MASSIVE BINARY STARS II. ROTATING SINGLE STARS UP TO CORE-COLLAPSE AND THEIR IMPACT ON THE EARLY SOLAR SYSTEM

HANNAH E. BRINKMAN,<sup>1,2</sup> J.W. DEN HARTOGH,<sup>1</sup> C. L. DOHERTY,<sup>1,3</sup> M. PIGNATARI,<sup>4,1,5,6</sup> AND M. LUGARO<sup>1,7,3</sup>

<sup>1</sup>*Konkoly Observatory, Research Centre for Astronomy and Earth Sciences (CSFK), Eötvös Loránd Research Network (ELKH), Konkoly Thege Miklós út 15-17, H-1121 Budapest, Hungary*

<sup>2</sup>*Graduate School of Physics, University of Szeged, Dom tér 9, Szeged, 6720 Hungary*

<sup>3</sup>*School of Physics and Astronomy, Monash University, VIC 3800, Australia*

<sup>4</sup>*E. A. Milne Centre for Astrophysics, Department of Physics and Mathematics, University of Hull, HU6 7RX, United Kingdom*

<sup>5</sup>*NuGrid Collaboration, <http://nugridstars.org>*

<sup>6</sup>*Joint Institute for Nuclear Astrophysics - Center for the Evolution of the Elements*

<sup>7</sup>*ELTE Eötvös Loránd University, Institute of Physics, Budapest 1117, Pázmány Péter sétány 1/A, Hungary*

(Received 30 April 2021; Revised 7 September 2021; Accepted 8 September 2021 for publication)

Submitted to ApJ

### ABSTRACT

Radioactive nuclei were present in the early Solar System, as inferred from analysis of meteorites. Many are produced in massive stars, either during their lives or their final explosions. In the first paper in this series (Brinkman et al. 2019), we focused on the production of <sup>26</sup>Al in massive binaries. Here, we focus on the production of another two short-lived radioactive nuclei, <sup>36</sup>Cl and <sup>41</sup>Ca, and the comparison to the early Solar System data. We used the MESA stellar evolution code with an extended nuclear network and computed massive (10–80 M<sub>⊙</sub>), rotating (with initial velocities of 150 and 300 km/s) and non-rotating single stars at solar metallicity (Z=0.014) up to the onset of core collapse. We present the wind yields for the radioactive isotopes <sup>26</sup>Al, <sup>36</sup>Cl, and <sup>41</sup>Ca, and the stable isotopes <sup>19</sup>F and <sup>22</sup>Ne. In relation to the stable isotopes, we find that only the most massive models, ≥ 60 M<sub>⊙</sub> and ≥ 40 M<sub>⊙</sub> give positive <sup>19</sup>F and <sup>22</sup>Ne yields, respectively, depending on the initial rotation rate. In relation to the radioactive isotopes, we find that the early Solar System abundances of <sup>26</sup>Al and <sup>41</sup>Ca can be matched with by models with initial masses ≥ 40 M<sub>⊙</sub>, while <sup>36</sup>Cl is matched only by our most massive models, ≥ 60 M<sub>⊙</sub>. <sup>60</sup>Fe is not significantly produced by any wind model, as required by the observations. Therefore, massive star winds are a favoured candidate for the origin of the very short-lived <sup>26</sup>Al, <sup>36</sup>Cl, and <sup>41</sup>Ca in the early Solar System.

*Keywords:* method: numerical - stars: evolution, mass-loss, winds, rotation

### 1. INTRODUCTION

Radioactive isotopes with short half-lives of less than a few Myr (hereafter short-lived radioactive isotopes, SLRs), and specifically the famous case of <sup>26</sup>Al with a half life of 0.72 Myr (Basunia & Hurst 2016), but also <sup>36</sup>Cl, <sup>41</sup>Ca, and <sup>60</sup>Fe, with half lives 0.301 Myr (Nica et al. 2012), 0.0994 Myr (Nesaraja & McCutchan 2016), and 2.62 Myr (Rugel et al. 2009), respectively, were present in the early Solar System (ESS). Their

abundances are inferred from meteoritic data reporting excesses in their daughter nuclei, for example, the ESS <sup>26</sup>Al/<sup>27</sup>Al ratio has been measured in calcium-aluminium-rich inclusions (CAIs) to be equal to  $(5.23 \pm 0.13) \times 10^{-5}$  (Jacobsen et al. 2008). These radioactive isotopes represent the fingerprint of the local nucleosynthesis that occurred nearby at the time and place of the birth of the Sun. Therefore, they give us clues about the environment and the circumstances of such birth (Adams 2010).

These four isotopes can be made in massive stars and expelled both by their winds, mainly during the Wolf-Rayet (WR) phase of stars with an initial mass

$\geq 35 M_{\odot}$  and/or due to binary interactions Brinkman et al. (2019), and in equal or larger amounts by their final core-collapse supernova (CCSN) (Meyer & Clayton (2000); Lugaro et al. (2018) and Lawson et al. in prep.). While the  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$  are ejected in significant amounts by both the wind and the CCSN, the amount of  $^{60}\text{Fe}$  in the stellar winds is negligible compared to that in the CCSN ejecta. This is because  $^{60}\text{Fe}$  is produced via neutron captures on the unstable  $^{59}\text{Fe}$ , and for this nucleus to capture a neutron instead of decaying, higher neutron densities ( $> 10^{10-11} \text{ cm}^{-3}$ ) are required than those produced during core He burning. Therefore,  $^{60}\text{Fe}$  is only produced in carbon shell-burning and in explosive He- and C-burning conditions (Limongi & Chieffi 2006; Tur et al. 2010; Jones et al. 2019) towards the end of the evolution. Instead, large amounts of  $^{26}\text{Al}$  are produced during H burning by proton captures on  $^{25}\text{Mg}$  and expelled by the winds. The majority of the  $^{26}\text{Al}$  is expelled together with  $^{35}\text{Cl}$  and  $^{40}\text{Ca}$ , which are produced during He burning by neutron captures on the preceding stable isotopes,  $^{35}\text{Cl}$  and  $^{40}\text{Ca}$ , respectively (e.g., Arnould et al. 1997, 2006; Gounelle & Meynet 2012; Brinkman et al. 2019).

Massive star winds have been suggested as a favoured site of the  $^{26}\text{Al}$  in the ESS (e.g., Arnould et al. 1997, 2006; Gounelle & Meynet 2012; Gaidos et al. 2009; Young 2014) also because they do not eject  $^{60}\text{Fe}$ . Candidate CCSNe sources of  $^{26}\text{Al}$  predict, instead, a more significant ejection of  $^{60}\text{Fe}$ , leading to  $^{60}\text{Fe}/^{56}\text{Fe}$  ratios orders of magnitude above the value observed in the ESS of  $\simeq 10^{-8}$  (Tang & Dauphas 2012; Trappitsch et al. 2018). Also, the ESS  $^{60}\text{Fe}/^{26}\text{Al}$  ratio is roughly two to three orders of magnitude lower than that observed via  $\gamma$ -ray, which sample the average galactic medium (Diehl 2013; Wang et al. 2020), and predicted by CCSN models (Austin et al. 2017; Sukhbold et al. 2016). This suggests that less  $^{60}\text{Fe}$  was present in the ESS as compared to the galactic average, and/or that extra source(s) of  $^{26}\text{Al}$  were present at the time of the birth of the Sun.

Arnould et al. (1997, 2006) considered production of several SLRs, including  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$ , by WR winds in both non-rotating and rotating models and concluded that these could have been the sources of these SLRs in the ESS. More recent studies have focused on  $^{26}\text{Al}$  only (Gounelle & Meynet 2012) and also concluded that WR winds are a possible source. In our first paper in this series, Brinkman et al. (2019) (hereafter Paper I), we also focused on  $^{26}\text{Al}$  and investigated how binary interactions between non-rotating massive stars can influence its wind yields. We showed that these interactions can lead to a significant increase in the  $^{26}\text{Al}$  wind yields in stars of masses 10-35  $M_{\odot}$ . For more massive stars, which

become WR stars, the effect of binary interaction is almost negligible.

In the present work we extend Paper I by computing the evolution of stellar models up to core collapse with a larger network of nuclear species and reactions, to calculate the wind yields of the SLRs  $^{36}\text{Cl}$ ,  $^{41}\text{Ca}$ , and  $^{60}\text{Fe}$ . We also include rotation in this study as this impacts stellar evolution and the winds (see, e.g., Maeder & Meynet 2012, for an overview). Our revision of the production of these SLRs is timely because updates in the ESS values of  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  have become recently available (Liu 2017; Tang et al. 2017). Moreover, the implementation of the mass-loss rates and of rotation represent some of the main uncertainties in the models of massive stars, and the differences obtained with different stellar evolution codes need to be considered carefully. We will therefore compare our results to those available in the literature.

With the extended nuclear network and calculations to beyond H burning, we also present new predictions for two stable isotopes that are produced during He burning and can be present in the stellar winds:  $^{19}\text{F}$  and  $^{22}\text{Ne}$ . These are of interest because Meynet & Arnould (2000) have shown that WR stars can contribute significantly to the galactic  $^{19}\text{F}$  abundance while Palacios et al. (2005) found that WRs are unlikely to be the source of galactic  $^{19}\text{F}$ , when including updated mass-loss prescriptions and reaction rates. Recently, the discussion around  $^{19}\text{F}$  was rekindled by Jönsson et al. (2014a,b, 2017) and Abia et al. (2019), who re-analysed observations of  $^{19}\text{F}$  and proposed that asymptotic giant branch stars are the most likely source of cosmic  $^{19}\text{F}$ . Still, due to the remaining uncertainty in both the mass-loss prescriptions and the reaction rates (see e.g. Stancliffe et al. 2005; Ugalde et al. 2008), WRs cannot be excluded as the sources of galactic  $^{19}\text{F}$  (for a recent overview, see e.g. Ryde et al. 2020). As for  $^{22}\text{Ne}$ , there are puzzling observations of an anomalous  $^{22}\text{Ne}/^{20}\text{Ne}$  ratio in cosmic rays, which is a factor of  $\sim 5$  higher than in the solar wind (Prantzos 2012). The comparison to model predictions may be a key to finding the source of cosmic rays in relation to OB associations of massive stars.

The paper structure is as follows; In Section 2 we describe the method and the physical input of our models. In Section 3 we discuss the stellar evolution results and all the relevant stellar evolution details for our models, and compare them to results from the literature. In Section 4 we present the nucleosynthetic yields of our models and compare these to various studies in the literature. In Section 5 we compare our findings to the abundances of the SLRs in the ESS and discuss which



(1995)<sup>1</sup> Below  $\log(T_{\text{eff}}) = 3.90$ , the alpha-parameter is undefined and we set it to 1, which makes the boost disappear. Above  $\log(T_{\text{eff}}) = 4.7$ , the alpha-parameter is again undefined. Here we have set it to 0.52, extrapolating the results of Lamers et al. (1995).

### 2.3. Yield calculations

Our focus is on the pre-supernova isotopic yields from the winds. To calculate these yields, we integrate over time, because wind mass-loss is a continuous process. For the stable isotopes, there are two yields to consider, the total yield and the net yield. The total yield is calculated as described above. The net yield is the total yield minus the initial abundance present in the star. For the SLRs the net yield is identical to the total yield, because there is no initial abundance present in the stars for these isotopes. The total yields and the relevant initial abundances are presented in Section 4.


## 3. STELLAR MODELS; RESULTS AND DISCUSSION

In this section we discuss the stellar evolution details of our models and the impact of rotation on them. In Table 1, selected relevant information of our models regarding the stellar evolution is presented: the total stellar mass and core mass at the end of the H, He, and C-core burning, the duration of these burning phases, the total lifetime, the total mass-loss, and the compactness parameter. The start and end points of the different burning phases are defined as in Götberg et al. (2018)<sup>2</sup>. By the end of carbon-burning the mass-loss phase has mostly ended (see Figure 3) and all our models have finished this stage. We then continued our models until a core-infall velocity of 300 km/s. In total, 31 out of our 36 models have reached this point, with numerical issues halting the calculations of the remaining 5 models slightly prior to this stage.

The duration of hydrogen burning,  $t_{\text{H}}$  (Column 3 of Table 1), is shown in Figure 1a. For all initial masses,  $t_{\text{H}}$  increases with the rotational velocity. The effect is the strongest at the lower mass end (17% for 10  $M_{\odot}$ ) and small for the three highest masses of our grid (9% for 80  $M_{\odot}$ ). The increasing duration of the main sequence is

<sup>1</sup> We implemented a step-function to connect their equation 2 and 3 at  $\log(T_{\text{eff}}) = 4.325$ , to match the data presented in their Figure 7.

<sup>2</sup> For completeness: the end of the core H burning stage is defined as when the central helium abundance is larger than 0.98 and the total luminosity produced by nuclear burning is larger than 0.5 the luminosity of the star, the end of He-core burning when the central carbon abundance is 0.4, the end of carbon-burning when the central carbon abundance is 0.01, and the end of oxygen burning when the central oxygen abundance is 0.04.




**Figure 1.** The duration of core hydrogen burning,  $t_{\text{H}}$  (solid lines in the top panel) and the duration of core helium burning,  $t_{\text{He}}$  (dashed lines in the bottom panel) for the three rotational velocities as a function of the initial mass.

due to rotational mixing. More hydrogen is mixed into the core from the envelope, adding more fuel to the core, and extending this burning phase. Together, this leads to larger hydrogen-depleted cores,  $M_{\text{c,He}}$ , at the end of the main sequence (dotted lines in Figure 2 and Column 4 in Table 1). The only exceptions to this trend are the two most massive models, 70 and 80  $M_{\odot}$ , which have a longer main-sequence yet smaller helium-core masses at the end of hydrogen burning when rotating.

This is especially noticeable for the highest initial rotational velocity. In these two most massive models, the increased mass loss limits the core growth. With more mass lost from the star, there is less fuel to add to the core and to increase its mass. This is shown by the solid lines in Figures 2 and 3, which give the final mass at the end of the main-sequence,  $M_{*,\text{H}}$  (Column 5 of Table 1) and the mass loss on the main sequence ( $M_{\text{ini}} - M_{*,\text{H}}$ ), respectively. For the models below  $\sim 50 M_{\odot}$ , there is little difference between the mass losses between the rotating and non-rotating models. The difference in final mass at the end of the main-sequences ranges between 1-4% for these stars. Above  $\sim 50 M_{\odot}$ , the extra mass loss for the rotating models becomes more significant, up to 50% more for our most massive model.

The mass-loss between the end of the main-sequence and the end of helium burning,  $M_{*,\text{H}} - M_{*,\text{He}}$ , represented by the dashed lines in Figure 3, is comparable between the rotating and non-rotating models with the same initial mass. The exceptions are again the two most massive models, for which the mass loss between the end of H- and the end of He-burning reduces with the





**Figure 2.** The total stellar mass  $M_{*,H}$ , (solid lines) and the the hydrogen depleted core mass  $M_{*,He}$ , (dashed lines) at the end of the main sequence for the three rotational velocities as a function of the initial mass.


increasing rotational velocity. This is because these models have already lost more mass on their main-sequence (solid lines in Figure 3), and as a consequence experience less mass loss afterwards.

As a result of the larger core masses at the end of hydrogen burning, the helium burning life-time,  $t_{He}$ , becomes shorter for most of the models with rotation (Figure 1b and Table 1). This is because the heavier the cores, the faster the burning. The three most massive models, however, the rotating models have smaller helium core masses at the end of hydrogen burning compared to their non-rotating counterparts. For these masses, the rotating models have longer  $t_{He}$  than their non-rotating counterparts. The turn-over point is around  $\sim 50 M_{\odot}$ , as can be seen in Figure 1b.

After helium burning barely any mass is lost from the stars (dotted lines in Figure 3). The helium-depleted core at the end of helium burning,  $M_{c,C}$  (Column 7 in Table 1) increase in mass for the models below  $\sim 50 M_{\odot}$ . For three highest masses however the mass of the helium-depleted cores decreases with the initial rotational velocity, which is a direct result of the smaller helium cores earlier.

We note that the mass of the carbon depleted core,  $M_{c,O}$  (reported in Column 10) is very sensitive to mixing in the final phases of the stellar evolution and a small fluctuation in the mixing can easily alter this value.

The mass loss during the main sequence always increases with higher rotational velocities (solid lines in Figure 3). The same applies for the total lifetime of the star,  $t_{tot}$  (Column 9 of Table 1). On the other hand, the total mass loss during the whole evolution,  $\Delta M$ ,



**Figure 3.** Mass loss for three different phases, the main sequence (solid lines,  $M_{ini}-M_{*,H}$ ), helium burning (dashed lines,  $M_{*,H}-M_{*,He}$ ) and carbon burning and beyond (dotted lines,  $M_{*,He}-M_{*,f}$ ), as a function of the initial mass. The final mass ( $M_{*,f}$ ) is the difference between the initial mass and the total mass loss,  $M_{ini}-\Delta M$ .

(see Table 1) is not always larger for the higher rotation rates.

The final column of Table 1 gives the compactness-parameter,  $\xi_{2.5}$ . This parameter, as defined by O'Connor & Ott (2011) in their Equation 10, determines how compact the core of the star is just before the collapse and therefore how difficult it is to explode the star. The compactness is sensitive to small changes in the structure, and is therefore strongly dependent on the model parameters and the codes used (see e.g., Sukhbold et al. 2016; Limongi & Chieffi 2018 and Schneider et al. 2021).

### 3.1. Comparison to other data sets

We compare our results primarily to those of Ekström et al. 2012, calculated using the GENEC stellar evolution code (hereafter E12), and Limongi & Chieffi 2018 calculated with the FRANEC stellar evolution code (hereafter LC18), for three reasons. The first is that both studies present rotating and non-rotating massive star models until the late phases of evolution. The second is that the implementation of rotation in these codes compared to what is used in MESA. Specifically, the treatment of rotation in MESA is based on a diffusive approximation, while in GENEC and FRANEC it is based on diffusion–advection approach (Maeder & Zahn 1998; Paxton et al. 2013), with it well established that the impact of rotation on stellar model computations varies depending on which rotational mixing approach is used (see, e.g, Maeder & Meynet 2000b). Fi-

nally, all the stellar evolution details from these two studies are publicly available, though for E12 we do not have yields for  $^{36}\text{Cl}$ ,  $^{41}\text{Ca}$ , and  $^{60}\text{Fe}$ . We compare our models to those from the literature at solar metallicity (taken to be either  $Z=0.014$ , or  $Z=0.02$ , depending on the source)

The E12 models have an initial rotational velocity of  $0.4\omega_{\text{crit}}$  (corresponding to  $\sim 260\text{-}350$  km/s). The LC18 models have the same initial rotational velocities of 150 and 300 km/s, as our models.

Our models show similar trends as in those works, such as the extended duration of the main-sequence and the increased mass-loss in the early phases of the evolution. Overall, all the different models show similar main-sequence lifetimes. However, the choice of mass loss prescription makes a big difference between the sets. The LC18 models lose much more mass in the early phases when rotation is included. When looking at their Figure 7, their lowest mass models including rotation move up almost vertically in the Hertzsprung-Russell diagram at the end of the main-sequence. This leads to a strong increase in the mass-loss, which is not seen in our models nor in those by E12.

#### 4. STELLAR YIELDS; RESULTS AND DISCUSSION

In this section we present and discuss the wind yields for  $^{26}\text{Al}$ ,  $^{19}\text{F}$ ,  $^{22}\text{Ne}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$  from our models and compare them to other studies in the literature. We do not present the  $^{60}\text{Fe}$  yields in quantitative detail because our models confirm all the previous results that this isotope is not ejected in the winds (the maximum yield we found is  $4.55 \times 10^{-10} M_{\odot}$  for the non-rotating  $70 M_{\odot}$  model). In Table 2 the yields of the five isotopes and the initial abundances of  $^{19}\text{F}$  and  $^{22}\text{Ne}$  are presented. The complete set of wind yields for all isotopes and models presented here, are available on Zenodo under a Creative Commons 4.0432 license: <https://doi.org/10.5281/zenodo.5497258>. As discussed in Section 3, for the low mass end,  $10\text{-}35 M_{\odot}$ , of the stars we investigate, most of the mass is lost between the main-sequence and the onset of helium-burning. At the high mass end,  $\sim 35\text{-}80 M_{\odot}$ , instead, the stars become WR stars and continue to lose mass even during and shortly after helium burning, stripping away not only the hydrogen-rich envelope, but also the top of the hydrogen-depleted core. This strongly impacts the yields of these stars, especially for the isotopes synthesized after hydrogen burning.

##### 4.1. Short lived radioactive isotopes


For  $^{26}\text{Al}$  (Figure 4a), the non-rotating yields are comparable to those from Paper I, with the exceptions of

the  $10$ ,  $40$ , and  $45 M_{\odot}$  models. The reason is that the  $10 M_{\odot}$  model loses more mass in our earlier study due to a longer main sequence lifetime, leading to a slightly higher yield. The  $40$  and  $45 M_{\odot}$  models are evolved until core-collapse here, while earlier they did not finish helium burning. This leads to a larger mass-loss and therefore a higher yield. The largest impact of the SNBs is at the lower mass-end,  $10\text{-}35 M_{\odot}$  (as shown in Paper I). However here we also find that, because both rotation and binary interaction lead to an increased mass-loss at similar points in the evolution, the impact of the SNBs becomes relatively smaller with increasing initial rotational velocity. The effect of binary interactions becomes negligible for models with initial masses of  $40\text{-}45 M_{\odot}$ , depending on the initial rotational velocity. Overall, our  $^{26}\text{Al}$  yields are only mildly sensitive to rotation and the wind prescription, since we changed the latter from Hamann et al. (1995) to Nugis & Lamers (2000) for the WR phase. The effect of the binary interactions is stronger than the effect of rotation or changing the wind.

Figures 4c and 4e show the  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  wind yields for our models. Unlike for  $^{26}\text{Al}$ , where the yield increases gradually with increasing mass, these yields show a sharp rise of almost 15 orders of magnitude at masses between  $20\text{-}30 M_{\odot}$ , depending on initial rotational velocity. This is easily understood when considering Figure 5, which shows the KHDs for non-rotating  $30 M_{\odot}$  and  $50 M_{\odot}$  models, with the  $^{41}\text{Ca}$  mass-fraction on the colour scale. Prior to core helium burning, there is no  $^{41}\text{Ca}$  present within these stars. The abundance of  $^{41}\text{Ca}$  (and  $^{36}\text{Cl}$ ) increases due to neutron captures during He burning on  $^{40}\text{Ca}$  (and  $^{35}\text{Cl}$ ), and it is also destroyed by neutron capture reactions (see, e.g., Lugaro et al. 2018, Figure 7). For the  $30 M_{\odot}$  model, this  $^{41}\text{Ca}$  (and  $^{36}\text{Cl}$ ) barely reaches the surface, leading to a very low yield. However, for the  $50 M_{\odot}$  model the top layers of the helium burning core are stripped away, leading to a more significant  $^{41}\text{Ca}$  ( $^{36}\text{Cl}$ ) yield. For the rotating models, the WR-phase is reached at a lower initial mass, and the increase in the yields also moves towards these lower initial masses. This means that these yields are highly sensitive to the wind prescription used, especially for the WR phase.

##### 4.2. Stable isotopes

Unlike the SLRs, the stable isotopes  $^{19}\text{F}$  and  $^{22}\text{Ne}$  are already present in the star at the time of its birth. Both these isotopes are typically destroyed by proton captures during hydrostatic H-burning as part of the CNO and NeNa-cycles through the  $^{19}\text{F}(p,\alpha)^{16}\text{O}$  and  $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$  reaction. During He-burning,  $^{19}\text{F}$  is pro-




**Figure 4.** The left panels show our SLR yields for the single star models (solid lines with symbols) for the three different initial rotational velocities. In panel (a),  $^{26}\text{Al}$ , the red line shows the yields from Paper I. The dashed lines give the “effective binary yields” defined as SNB yields averaged on the period range given in Paper I assuming a flat period distribution. These effective binary yields are presented to help visualise the effect of binary interactions. The shaded area between the single star models and the effective binary yields therefore represents the results from the potential parameter space covered by binary systems. The right panels show the single star models of this study together with the wind yields of various other studies, as listed in the legend.

**Table 1.** Selected details of the evolution of our stellar models.  $M_{\text{ini}}$  is the initial mass in  $M_{\odot}$ .  $V_{\text{ini}}$  is the initial rotational velocity in km/s.  $t_{\text{H}}$ ,  $t_{\text{He}}$ , and  $t_{\text{tot}}$  are the duration of hydrogen burning, helium burning, and the total evolution time in Myr, respectively.  $M_{*,\text{H}}$ ,  $M_{*,\text{He}}$ , and  $M_{*,\text{C}}$  are the masses of the stars at the end of their respective burning phases.  $M_{\text{c,He}}$ ,  $M_{\text{c,C}}$ , and  $M_{\text{c,O}}$  are the masses of the hydrogen-depleted core, the helium-depleted core, and the carbon depleted core at the end of the corresponding burning phases in  $M_{\odot}$ .  $\Delta M$  is the total mass lost in  $M_{\odot}$ .  $\xi_{2.5}$  is the compactness of the star at the final model.

| $M_{\text{ini}}$<br>( $M_{\odot}$ ) | $V_{\text{ini}}$<br>(km/s) | $t_{\text{H}}$<br>(Myr) | $M_{\text{c,He}}$<br>( $M_{\odot}$ ) | $M_{*,\text{H}}$<br>( $M_{\odot}$ ) | $t_{\text{He}}$<br>(Myr) | $M_{\text{c,C}}$<br>( $M_{\odot}$ ) | $M_{*,\text{He}}$<br>( $M_{\odot}$ ) | $t_{\text{tot}}$<br>(Myr) | $M_{\text{c,O}}$<br>( $M_{\odot}$ ) | $M_{*,\text{C}}$<br>( $M_{\odot}$ ) | $\Delta M$<br>( $M_{\odot}$ ) | $\xi_{2.5}$<br>- |
|-------------------------------------|----------------------------|-------------------------|--------------------------------------|-------------------------------------|--------------------------|-------------------------------------|--------------------------------------|---------------------------|-------------------------------------|-------------------------------------|-------------------------------|------------------|
| 10                                  | $0^1$                      | 23.00                   | 1.83                                 | 9.78                                | 1.98                     | 1.52                                | 9.11                                 | 25.37                     | 1.38                                | 9.04                                | 0.96                          | 6.09e-3          |
|                                     | $150^1$                    | 23.95                   | 1.92                                 | 9.75                                | 1.89                     | 1.58                                | 9.06                                 | 26.21                     | 1.34                                | 8.99                                | 1.01                          | 7.69e-3          |
|                                     | 300                        | 26.96                   | 2.04                                 | 9.64                                | 1.69                     | 1.84                                | 8.62                                 | 28.99                     | 1.91                                | 8.55                                | 1.45                          | 0.017            |
| 15                                  | 0                          | 12.15                   | 3.62                                 | 14.58                               | 0.96                     | 3.41                                | 11.49                                | 13.26                     | 1.71                                | 11.35                               | 3.65                          | 0.098            |
|                                     | 150                        | 12.49                   | 3.73                                 | 14.50                               | 0.92                     | 3.54                                | 11.22                                | 13.56                     | 1.60                                | 11.08                               | 3.91                          | 0.078            |
|                                     | 300                        | 14.05                   | 4.09                                 | 14.23                               | 0.86                     | 4.04                                | 9.96                                 | 15.03                     | 1.66                                | 9.80                                | 5.20                          | 0.086            |
| 20                                  | 0                          | 8.53                    | 5.73                                 | 19.12                               | 0.66                     | 5.66                                | 11.24                                | 9.29                      | 2.49                                | 11.02                               | 8.98                          | 0.23             |
|                                     | 150                        | 8.68                    | 5.82                                 | 18.96                               | 0.65                     | 5.83                                | 10.77                                | 9.43                      | 1.92                                | 10.55                               | 9.44                          | 0.16             |
|                                     | $300^1$                    | 9.87                    | 6.45                                 | 18.39                               | 0.60                     | 6.62                                | 9.39                                 | 10.55                     | 2.99                                | 9.28                                | 10.71                         | 0.40             |
| 25                                  | $0^1$                      | 6.80                    | 7.99                                 | 23.20                               | 0.53                     | 8.13                                | 11.04                                | 7.41                      | 1.71                                | 10.92                               | 14.07                         | 0.12             |
|                                     | 150                        | 6.96                    | 8.23                                 | 22.87                               | 0.53                     | 8.06                                | 11.27                                | 7.56                      | 1.77                                | 11.04                               | 13.95                         | 0.11             |
|                                     | 300                        | 7.68                    | 8.86                                 | 22.12                               | 0.50                     | 8.63                                | 11.06                                | 8.25                      | 1.90                                | 10.96                               | 14.03                         | 0.11             |
| 30                                  | $0^1$                      | 5.80                    | 10.35                                | 26.89                               | 0.47                     | 10.73                               | 13.56                                | 6.32                      | 2.44                                | 13.47                               | 16.52                         | 0.23             |
|                                     | 150                        | 6.00                    | 10.80                                | 26.13                               | 0.45                     | 10.70                               | 13.25                                | 6.50                      | 2.23                                | 13.12                               | 16.87                         | 0.22             |
|                                     | 300                        | 6.47                    | 11.29                                | 25.39                               | 0.44                     | 10.81                               | 13.24                                | 6.96                      | 2.16                                | 13.12                               | 16.86                         | 0.19             |
| 35                                  | 0                          | 5.15                    | 12.74                                | 30.38                               | 0.42                     | 12.95                               | 15.64                                | 5.62                      | 3.00                                | 15.46                               | 19.51                         | 0.18             |
|                                     | 150                        | 5.27                    | 13.05                                | 29.75                               | 0.41                     | 12.55                               | 15.15                                | 5.73                      | 2.79                                | 15.00                               | 19.98                         | 0.23             |
|                                     | 300                        | 5.70                    | 13.90                                | 29.09                               | 0.40                     | 12.12                               | 14.83                                | 6.15                      | 2.69                                | 14.65                               | 20.33                         | 0.29             |
| 40                                  | 0                          | 4.69                    | 15.17                                | 33.23                               | 0.39                     | 13.63                               | 16.40                                | 5.12                      | 3.21                                | 16.24                               | 23.74                         | 0.19             |
|                                     | 150                        | 4.78                    | 15.49                                | 32.55                               | 0.38                     | 13.21                               | 16.11                                | 5.20                      | 3.07                                | 16.04                               | 23.93                         | 0.23             |
|                                     | 300                        | 5.15                    | 16.39                                | 30.94                               | 0.37                     | 13.72                               | 16.70                                | 5.56                      | 3.09                                | 16.65                               | 23.32                         | 0.26             |
| 45                                  | 0                          | 4.35                    | 17.57                                | 35.74                               | 0.36                     | 14.85                               | 17.95                                | 4.76                      | 3.43                                | 17.90                               | 27.06                         | 0.26             |
|                                     | 150                        | 4.48                    | 18.28                                | 34.28                               | 0.35                     | 15.26                               | 18.20                                | 4.88                      | 3.28                                | 18.13                               | 26.82                         | 0.26             |
|                                     | 300                        | 4.81                    | 19.22                                | 34.03                               | 0.35                     | 15.57                               | 18.63                                | 5.20                      | 3.62                                | 18.55                               | 26.41                         | 0.31             |
| 50                                  | 0                          | 4.09                    | 20.05                                | 37.93                               | 0.35                     | 16.84                               | 19.92                                | 4.48                      | 3.77                                | 19.87                               | 30.07                         | 0.31             |
|                                     | 150                        | 4.21                    | 20.79                                | 35.93                               | 0.34                     | 16.68                               | 19.80                                | 4.59                      | 3.53                                | 19.71                               | 30.23                         | 0.30             |
|                                     | 300                        | 4.47                    | 21.77                                | 36.26                               | 0.38                     | 17.70                               | 20.81                                | 4.85                      | 3.99                                | 20.73                               | 29.21                         | 0.33             |
| 60                                  | 0                          | 3.71                    | 25.04                                | 41.39                               | 0.33                     | 19.49                               | 22.76                                | 4.07                      | 4.53                                | 22.67                               | 37.25                         | 0.31             |
|                                     | 150                        | 3.78                    | 25.84                                | 43.08                               | 0.32                     | 20.09                               | 23.37                                | 4.14                      | 4.60                                | 23.29                               | 36.63                         | 0.32             |
|                                     | 300                        | 4.02                    | 25.89                                | 34.26                               | 0.35                     | 13.19                               | 16.01                                | 4.40                      | 3.15                                | 15.93                               | 43.99                         | 0.23             |
| 70                                  | 0                          | 3.43                    | 30.29                                | 50.32                               | 0.31                     | 22.86                               | 26.36                                | 3.78                      | 15.14 <sup>2</sup>                  | 26.25                               | 43.65                         | 0.46             |
|                                     | 150                        | 3.53                    | 30.82                                | 43.31                               | 0.32                     | 16.89                               | 20.08                                | 3.89                      | 3.88                                | 19.99                               | 49.92                         | 0.32             |
|                                     | 300                        | 3.67                    | 28.80                                | 33.48                               | 0.36                     | 11.10                               | 13.68                                | 4.07                      | 2.74                                | 13.60                               | 56.30                         | 0.29             |
| 80                                  | 0                          | 3.24                    | 34.98                                | 50.72                               | 0.31                     | 20.68                               | 24.10                                | 3.58                      | 4.84                                | 23.99                               | 55.90                         | 0.36             |
|                                     | 150                        | 3.36                    | 30.33                                | 33.28                               | 0.36                     | 10.67                               | 13.24                                | 3.76                      | 2.45                                | 13.17                               | 66.72                         | 0.25             |
|                                     | 300                        | 3.54                    | 21.99                                | 25.20                               | 0.39                     | 9.10                                | 11.54                                | 3.97                      | 2.06                                | 11.46                               | 68.42                         | 0.16             |

<sup>1</sup> This run was terminated before core collapse was reached due to numerical difficulties, however, negligible mass loss is expected after this point. <sup>2</sup> This run experienced computational difficulties in the final phases, leading to a much larger  $M_{\text{c,O}}$  than for any of the other models.





**Figure 5.** Kippenhahn diagrams for a  $30 M_{\odot}$  star in the left panel and for a  $50 M_{\odot}$  the right panel, both non-rotating. The horizontal axis shows the time left until core collapse. The colour-scale shows the  $^{41}\text{Ca}$  mass-fraction. The green-hatched areas represent convective regions, the blue hatched areas, overshoot, and the red areas semi-convection. The red dotted line is the size of the hydrogen-depleted core.

duced by  $\alpha$ -captures on  $^{15}\text{N}$ , and also depleted by  $\alpha$ -captures, producing  $^{22}\text{Ne}$  (Meynet & Arnould 2000).  $^{22}\text{Ne}$  is produced in the helium burning, mainly by double  $\alpha$ -captures on  $^{14}\text{N}$ , while it is not significantly destroyed. To expel a significant amount  $^{19}\text{F}$  into the interstellar medium, the He-core needs to be exposed in an early stage of helium-burning, before  $^{19}\text{F}$  is destroyed.

For the majority of the models shown in Figure 6, the initial abundance (red lines) of the stable isotopes are higher than the amount expelled from the stars, and therefore the net yield is negative. Only the heaviest, rotating models in our set ( $\geq 60 M_{\odot}$ ) produce a positive net yield. This is in agreement with the earlier results for  $^{19}\text{F}$  by Meynet & Arnould (2000) and Palacios et al. (2005).


For  $^{22}\text{Ne}$  the stars have positive yields for masses  $40$ – $45 M_{\odot}$  and above, depending on initial rotational velocity. This is because, unlike  $^{19}\text{F}$ ,  $^{22}\text{Ne}$  is not completely destroyed during hydrogen burning and during helium burning,  $^{22}\text{Ne}$  is not completely destroyed while  $^{19}\text{F}$  does. Therefore, it is easier to obtain a positive yield. These yields are not only sensitive to the wind prescription, which determines how much material is ejected, but also to the internal mixing processes due to rotation, which leads to more or less destruction of the initial abundance.

#### 4.3. Comparison to other data sets

The right panels of Figures 4 and 6 show the results of our models along with several other studies from the literature. On top LC18 and E12, here we also consider the non-rotating models from Sukhbold et al. (2016),

computed with the KEPLER stellar evolution code, Pignatari et al. (2016), computed with GENEC for the for mass range of interest, and from Ritter et al. (2018), computed with MESA version 3709. We note that E12 and Pignatari et al. (2016) both use the GENEC code, however, the models of Pignatari et al. (2016) cover all burning stages, therefore all isotopes we are interested in are included, while for for E12 we do not have yields for  $^{36}\text{Cl}$ ,  $^{41}\text{Ca}$ , and  $^{60}\text{Fe}$ . Overall, the yield sets are in broad agreement within the uncertainties of the stellar evolution modelling. The models that stand out the most are the low-mass ( $10$ – $30 M_{\odot}$ ) LC18 models. These models show an increase in the SLR yields from non-rotating to rotating orders of magnitude higher than found by the other models. For  $^{26}\text{Al}$ , the LC18 increase is comparable to the increase in the  $^{26}\text{Al}$  yields we found in Paper I between single stars and binary models. This behaviour in the LC18 models is due to a very strong increase in the mass loss compared to their non-rotating models, leading to an increased yield. This difference in mass loss is due to the treatment of rotational mixing and the formation of a dust driven wind, as explained in Chieffi & Limongi (2013).

For the stable isotopes, also the highest mass models by Sukhbold et al. (2016) stand out. These models lose much more mass than our  $60$ – $80 M_{\odot}$  stars, especially their  $80 M_{\odot}$  model loses nearly  $74 M_{\odot}$ , compared to  $\simeq 56 M_{\odot}$  for our non-rotating model. The  $20$  and  $25 M_{\odot}$  E12 models have higher yields than the overall trend for these two masses, because these stars have a slight boost to their winds, independent of rotation.



**Figure 6.** The left panels show our  $^{19}\text{F}$  (top) and  $^{22}\text{Ne}$  (bottom) yields and the initial abundances. The red line indicates the initial abundance of the models. The right panels show the stellar models of this study together with the yields of various other studies.

## 5. EARLY SOLAR SYSTEM

The radioactive isotopes we have studied in the previous section were inferred to be present in the early Solar System (ESS) from observed excesses of their daughter nuclei in meteoritic inclusions. In this section, we consider a simple dilution model for  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$ , to investigate if their abundances in the ESS can be explained self-consistently with the models presented here. While other stellar objects in the galaxy can produce these isotopes, such as novae and asymptotic giant branch stars, much research has focused on massive stars as ESS polluters, because these stars live short enough to be able to eject material within star forming regions which have typical lifetimes of at most a few tens of Myr (Murray 2011). Furthermore, massive star winds are preferred by several authors (Gaidos et al. 2009; Gounelle & Meynet 2012; Young 2014; Dwarkadas et al. 2017) as a most favoured site of origin for  $^{26}\text{Al}$ , because

there are several difficulties for core-collapse supernovae to produce the abundances of SLRs in the proportion required to match the ESS values (see discussion in Lugaro et al. 2018). For example, they produce too high abundances of  $^{60}\text{Fe}$  and  $^{53}\text{Mn}$ . Both these isotopes have half lives 4 to 50 times higher than the SLRs that we consider in the calculation here, therefore, their abundances in the ESS can be explained by decaying their abundances in the interstellar medium as derived by galactic chemical evolution (see, e.g., Wasserburg et al. 2006; Tang & Dauphas 2012; Trappitsch et al. 2018; Côté et al. 2019a,b).

### 5.1. The dilution method

The comparison to the ESS values is performed in four steps:

Step 1 is to determine a “dilution factor”,  $f$  for each stellar model. This is defined as  $f = \frac{M_{\text{SLR}}^{\text{ESS}}}{M_{\text{SLR}}^*}$ , where  $M_{\text{SLR}}^{\text{ESS}}$

**Table 2.** Isotopic yields in  $M_{\odot}$ .  $M_{\text{ini}}$  is the initial mass in  $M_{\odot}$ .  $V_{\text{ini}}$  is the initial rotational velocity in km/s. For the stable isotopes, the initial amount present in the star is also given. The yields tabulated here are not corrected for radioactive decay that might take place in the ISM during the evolution of the star. Yields for all 209 isotopes in the nuclear network are available on Zenodo: <https://doi.org/10.5281/zenodo.5497258>.

| $M_{\text{ini}}$<br>( $M_{\odot}$ ) | $V_{\text{ini}}$<br>(km/s) | $^{19}\text{F}_{\text{ini}}$<br>( $M_{\odot}$ ) | $^{19}\text{F}$<br>( $M_{\odot}$ ) | $^{22}\text{Ne}_{\text{ini}}$<br>( $M_{\odot}$ ) | $^{22}\text{Ne}$<br>( $M_{\odot}$ ) | $^{26}\text{Al}$<br>( $M_{\odot}$ ) | $^{36}\text{Cl}$<br>( $M_{\odot}$ ) | $^{41}\text{Ca}$<br>( $M_{\odot}$ ) |
|-------------------------------------|----------------------------|---|------------------------------------|--|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| 10                                  | 0                          | 4.67e-06  | 4.61e-07                           | 8.90e-4  | 8.47e-05                            | 7.19e-11                            | 8.82e-23                            | 7.90e-23                            |
|                                     | 150                        | 4.67e-06  | 4.71e-07                           | 8.90e-4  | 8.68e-05                            | 2.78e-10                            | 1.11e-22                            | 1.29e-22                            |
|                                     | 300                        | 4.68e-06  | 4.20e-07                           | 8.92e-4  | 7.77e-05                            | 1.40e-10                            | 1.67e-22                            | 1.70e-22                            |
| 15                                  | 0                          | 6.42e-06  | 1.69e-06                           | 1.32e-3  | 3.12e-4                             | 8.85e-09                            | 3.00e-22                            | 1.41e-21                            |
|                                     | 150                        | 6.42e-06  | 1.74e-06                           | 1.32e-3  | 3.24e-4                             | 1.47e-08                            | 5.12e-22                            | 2.10e-21                            |
|                                     | 300                        | 6.45e-06  | 1.21e-06                           | 1.33e-3  | 2.26e-4                             | 2.34e-08                            | 5.32e-22                            | 3.06e-21                            |
| 20                                  | 0                          | 8.22e-06  | 4.01e-06                           | 1.79e-3  | 7.43e-4                             | 1.81e-07                            | 1.94e-21                            | 1.28e-20                            |
|                                     | 150                        | 8.24e-06  | 4.04e-06                           | 1.79e-3  | 7.57e-4                             | 1.92e-07                            | 9.48e-22                            | 1.33e-20                            |
|                                     | 300                        | 8.28e-06  | 2.04e-06                           | 1.79e-3  | 3.86e-4                             | 5.87e-07                            | 1.75e-21                            | 2.87e-20                            |
| 25                                  | 0                          | 9.67e-06  | 5.87e-06                           | 2.23e-3  | 1.09e-3                             | 1.17e-06                            | 2.80e-21                            | 6.27e-20                            |
|                                     | 150                        | 9.69e-06  | 5.23e-06                           | 2.23e-3  | 9.89e-4                             | 1.59e-06                            | 4.52e-21                            | 9.28e-20                            |
|                                     | 300                        | 9.75e-06  | 2.28e-06                           | 2.24e-3  | 4.41e-4                             | 3.93e-06                            | 4.98e-16                            | 2.06e-15                            |
| 30                                  | 0                          | 1.11e-05  | 6.38e-06                           | 2.69e-3  | 1.19e-3                             | 3.41e-06                            | 6.76e-21                            | 2.22e-19                            |
|                                     | 150                        | 1.12e-05  | 5.84e-06                           | 2.70e-3  | 1.12e-3                             | 4.68e-06                            | 3.30e-15                            | 1.42e-14                            |
|                                     | 300                        | 1.12e-05  | 2.39e-06                           | 2.70e-3  | 4.77e-4                             | 9.89e-06                            | 1.45e-13                            | 6.66e-13                            |
| 35                                  | 0                          | 1.33e-05  | 6.77e-06                           | 3.20e-3  | 1.27e-3                             | 8.44e-06                            | 4.70e-16                            | 1.96e-15                            |
|                                     | 150                        | 1.33e-05  | 5.84e-06                           | 3.20e-3  | 1.13e-3                             | 1.06e-05                            | 6.85e-13                            | 3.23e-12                            |
|                                     | 300                        | 1.34e-05  | 2.51e-06                           | 3.20e-3  | 5.27e-4                             | 2.11e-05                            | 1.85e-10                            | 9.29e-10                            |
| 40                                  | 0                          | 1.42e-05  | 7.06e-06                           | 3.64e-3  | 1.36e-3                             | 1.88e-05                            | 7.99e-11                            | 4.015e-10                           |
|                                     | 150                        | 1.43e-05  | 6.14e-06                           | 3.64e-3  | 2.13e-3                             | 2.05e-05                            | 3.56e-08                            | 6.07e-08                            |
|                                     | 300                        | 1.44e-05  | 2.82e-06                           | 3.65e-3  | 5.09e-3                             | 3.43e-05                            | 1.82e-07                            | 2.99e-07                            |
| 45                                  | 0                          | 1.61e-05  | 7.40e-06                           | 4.13e-3  | 5.18e-3                             | 2.94e-05                            | 1.45e-07                            | 2.44e-07                            |
|                                     | 150                        | 1.61e-05  | 6.31e-06                           | 4.13e-3  | 1.01e-2                             | 3.15e-05                            | 3.59e-07                            | 5.86e-07                            |
|                                     | 300                        | 1.63e-05  | 3.23e-06                           | 4.13e-3  | 1.54e-2                             | 5.32e-05                            | 5.99e-07                            | 9.78e-07                            |
| 50                                  | 0                          | 1.75e-05  | 7.82e-06                           | 4.60e-3  | 1.21e-2                             | 4.00e-05                            | 4.18e-07                            | 6.89e-07                            |
|                                     | 150                        | 1.76e-05  | 7.00e-06                           | 4.60e-3  | 2.30e-2                             | 4.30e-05                            | 8.78e-07                            | 1.43e-06                            |
|                                     | 300                        | 1.77e-05  | 3.42e-06                           | 4.61e-3  | 2.14e-2                             | 7.04e-05                            | 8.26e-07                            | 1.34e-06                            |
| 60                                  | 0                          | 1.91e-05  | 9.00e-06                           | 5.50e-3  | 3.84e-2                             | 6.65e-05                            | 1.48e-06                            | 2.42e-06                            |
|                                     | 150                        | 1.91e-05  | 7.28e-06                           | 5.50e-3  | 4.23e-2                             | 7.16e-05                            | 1.63e-06                            | 2.67e-06                            |
|                                     | 300                        | 1.93e-05  | 1.91e-05                           | 5.51e-3  | 0.12                                | 1.44e-4                             | 3.88e-06                            | 8.37e-06                            |
| 70                                  | 0                          | 2.09e-05  | 9.21e-06                           | 6.43e-3  | 5.58e-2                             | 9.70e-05                            | 2.10e-06                            | 3.46e-06                            |
|                                     | 150                        | 2.10e-05  | 1.78e-05                           | 6.43e-3  | 1.38e-1                             | 1.25e-4                             | 5.00e-06                            | 9.55e-06                            |
|                                     | 300                        | 2.11e-05  | 3.11e-05                           | 6.43e-3  | 0.17                                | 2.36e-4                             | 4.48e-06                            | 1.14e-05                            |
| 80                                  | 0                          | 2.43e-05  | 1.49e-05                           | 7.41e-3  | 0.14                                | 1.51e-4                             | 5.40e-06                            | 9.55e-06                            |
|                                     | 150                        | 2.44e-05  | 5.16e-05                           | 7.41e-3  | 0.18                                | 2.69e-3                             | 4.52e-06                            | 1.20e-05                            |
|                                     | 300                        | 2.46e-05  | 3.51e-05                           | 7.41e-3  | 0.11                                | 4.29e-4                             | 2.59e-06                            | 6.99e-06                            |

is the mass of a given SLR in the ESS, and  $M_{\text{SLR}}^*$  is the mass of the same SLR ejected by the stellar wind, i.e., the total yield. We use  $^{26}\text{Al}$  to determine  $f$  and then apply the same value to  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$ <sup>3</sup>, this is because the  $^{26}\text{Al}/^{27}\text{Al}$  ratio in the ESS is very well established to  $5.23 \times 10^{-5}$ , as reported by [Jacobsen et al. \(2008\)](#) and recently confirmed by [Luu et al. \(2019\)](#). We derive an initial amount of  $^{26}\text{Al}$  of  $3.1 \times 10^{-9} M_{\odot}$  assuming the solar abundance of  $^{27}\text{Al}$  ([Lodders 2003](#)) and a total mass of  $1 M_{\odot}$  to be polluted (see details in [Lugaro et al. 2018](#)).

Step 2 is to determine the “delay time” ( $\Delta t$ ), which is the time interval between wind ejection and the formation of the first solids (the calcium-aluminium rich inclusions (CAIs)) in the ESS, for which time the SLR values are given. After we have applied our determined  $f$  to the yields of  $^{41}\text{Ca}$ , to obtain the diluted amount of this isotope in Step 1, we decay the  $^{41}\text{Ca}$  until we reach the observed ESS  $^{41}\text{Ca}/^{40}\text{Ca}$  ratio of  $4.6 \times 10^{-9}$ . The time needed, is the delay time.

Step 3 is to calculate a new abundance of  $^{26}\text{Al}$  by reverse decaying the initial ESS amount of  $^{26}\text{Al}$  using the delay time from Step 2. With this we recalculate  $f$  and use it to repeat Step 2. We continue this iteration until we converge to a  $\Delta t$  value within a 10% difference from the previous value.

Step 4 apply the final  $f$  to calculate the diluted  $^{36}\text{Cl}$  abundance. We then repeat Step 2 for  $^{36}\text{Cl}$  also to determine a delay time for  $^{36}\text{Cl}$  ( $\Delta t_{\text{Cl}}$ ) using the ESS  $^{36}\text{Cl}/^{35}\text{Cl}$  ratio of  $2.44 \times 10^{-5}$  measured in the *Curious Marie* CAI ([Tang et al. 2017](#)).

A few remarks need to be made about this method. First, even though  $^{26}\text{Al}$  is not produced in the same evolutionary phase as  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$ , the bulk of these isotopes are expelled into the interstellar medium at the same time. Therefore, we do not need to take into account, for example, that some  $^{26}\text{Al}$  might have decayed before  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  were ejected the star. Second, we use the  $^{41}\text{Ca}/^{40}\text{Ca}$ -ratio to obtain  $\Delta t$  even though this ratio is not well constrained in the ESS ([Liu 2017](#)), because  $^{41}\text{Ca}$  is the shortest lived of the three isotopes considered here. Therefore, it is the most sensitive chronometer to short timescales (see also [Wasserburg et al. 2006](#)). Moreover, the abundances of both  $^{36}\text{Cl}$

and  $^{41}\text{Ca}$  can have a contribution from irradiation by solar cosmic rays in the disk. Higher values of  $^{36}\text{Cl}$  than the ESS value used here have been measured in other meteoritic inclusions and can be produced by irradiation within the ESS. The value measured in *Curious Marie* probably represents the primordial value derived from a stellar source because  $^{36}\text{Cl}$  coexists in this CAI with the canonical value of  $^{26}\text{Al}$  ([Tang et al. 2017](#)). We note also that the measurements of the ESS values of  $^{41}\text{Ca}$  and  $^{36}\text{Cl}$  might be affected by systematic uncertainties, because the abundance of  $^{41}\text{Ca}$  is very low and the abundance of  $^{36}\text{Cl}$  is based on the measurement of its roughly 2% decay channel into  $^{38}\text{S}$ . For  $^{41}\text{Ca}$  the latest data on a handful of CAIs ([Liu 2017](#)) demonstrate the presence of this very short-lived isotope in the ESS, however, the data precision is not high enough to be able to resolve possible heterogeneities.

## 5.2. Results and comparison to other studies

We apply the method described in Sec. 5.1 to our models and they are considered to be a solution for the ESS when the delay times for  $^{41}\text{Ca}$  and  $^{36}\text{Cl}$  are comparable with a factor of up to 5. For our non-rotating models, a solution can be found for  $^{26}\text{Al}$  and  $^{41}\text{Ca}$  in the mass range 45 to 80  $M_{\odot}$  and  $\Delta t$  between 0.7 and 1 Myr. In order to also match the  $^{36}\text{Cl}$  abundance, the mass-range needs to be restricted to stars with an initial mass of 60  $M_{\odot}$  and higher. There still is a small inconsistency, because  $\Delta t$  needs to be lower for  $^{36}\text{Cl}$  than for  $^{41}\text{Ca}$ , as the delay time for  $^{36}\text{Cl}$  is between 0.2 and 0.5 Myr. The results for our rotating models are similar, except that the mass range for which a solution is possible for  $^{26}\text{Al}$  and  $^{41}\text{Ca}$  may extend down to 40  $M_{\odot}$ . If we consider the three isotopes, it is possible to find a solution for initial rotational velocity of 150 and 300 km/s at masses 50-80  $M_{\odot}$  and 60 and 70  $M_{\odot}$ , respectively.

In Figure 7 we look more closely at a selected set of models. It shows the abundance ratios (R/S) for the three SLRs (R) over their stable reference isotope (S) versus delay time for the dilution factor  $f$ , for our stellar models with a mass of 40, 60, and 80  $M_{\odot}$ . The horizontal bars represent the ESS-ratios and their uncertainties. We compare our results with the results calculated from the LC18 models for the same initial masses.

For the 40  $M_{\odot}$  models (Figure 7a), the  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  yields from the non-rotating model (solid lines) are too low to match their ESS ratios. Only our rotating models (dashed and dotted lines in panel a) can match both the Al- and Ca-ratios (at  $10^{5.9}$  yrs), while the the Cl-ratio is between one and two orders of magnitude too low. This excludes this star as a potential solution for the ESS. For the LC18 models (Figure 7b), the non-rotating

<sup>3</sup> If the injection of the SLRs occurred in the form of dust grains, chemistry and dust formation could lead to different  $f$  values for the different isotopes because Al, Ca, and Cl are different elements. While Al and Ca should behave chemically in a very similar way, the situation for Cl may be different. We do not consider these uncertainties here.

model does not match Cl- or the Ca-ratio either. As for our models, the rotating models match both the Al- and Ca-ratios. The difference is that in the LC18 models the Cl-ratio is matched very early on in the calculation, at  $10^{4.9}$  yrs, while the other ratios are matched after 1 Myr, while in our models the Cl-ratio cannot be matched at all. However, due to this large difference in  $\Delta t$ , these models can still not be considered a potential solution.

For the 60  $M_{\odot}$  models (Figure 7c), all of our models can match the three isotopic ratios, the Cl-ratio at  $10^{5.5}$  yrs and the other two ratios at  $10^6$  yrs. For the LC18 models (Figure 7d), only their model with an initial rotational velocity of 150 km/s matches the Cl-ratio, but too early ( $10^{4-4.6}$  yrs) to be considered a solution together with the Al- and Ca-ratios.

Finally, for the 80  $M_{\odot}$  (Figure 7e), we can match the ESS ratios of the SLRs with our non-rotating model and the model rotating at an initial velocity of 150 km/s. There is still a difference in the delay times for  $^{41}\text{Ca}$  and  $^{36}\text{Cl}$  of a factor of 2. For the 80  $M_{\odot}$  LC18 models (Figure 7f), only the non-rotating model matches all three ratios, and the Cl-ratio is matched at 0.03 Myr, while the other two are matched at 1.14 Myr. This difference is again too large to consider this as a solution. The rotating models cannot match the Cl-ratio.

The difference between the sets can be explained by considering Figures 4d and 4f. While the  $^{41}\text{Ca}$  yields are very comparable between the two sets of both non-rotating and rotating models for 60 and 80  $M_{\odot}$ , the non-rotating and rotating  $^{36}\text{Cl}$  yields are lower for the LC18 models due to the differences in mass loss (see Section 4.1) compared to our yields. Therefore, there is less  $^{36}\text{Cl}$  compared to  $^{41}\text{Ca}$  in the LC18 models than in our models, making it harder to match both ratios within the uncertainties. For the 40  $M_{\odot}$  models, the rotating yields given by LC18 are higher than our rotating yields for  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$  which explains why their rotating models can match for the Cl-ratio, while our models can not.

Our results that the three radionuclides  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$  can be ejected by the winds of a variety of WR stars at relative levels compatible with the meteoritic observations are also in qualitative agreement with the results of Arnould et al. (1997, 2006). Because these authors used  $^{107}\text{Pd}$  to calculate  $f$ , our results are not directly comparable. However, if we consider the delay times shown in Figures 5-7 of Arnould et al. (2006), we find similar values for the delay time for the Ca-ratio, between  $10^{5.5-10^6}$  yrs. Their delay time for the Cl-ratio for their 40  $M_{\odot}$  model is much closer to their delay time for the Ca-ratio compared to our models and those by LC18. However, the delay times for the Cl-ratio

are worse for their 60 and 85  $M_{\odot}$  models than for our models or those by LC18. Our result also confirm the analysis of the production of  $^{26}\text{Al}$  in WR stars of mass between 32 and 120  $M_{\odot}$  by Gounelle & Meynet (2012). Our  $f$  factor may be compared to their  $\eta_{\text{wind}}/1000$  in their Equation (2), for which they find values ranging down to  $2 \times 10^{-5}$ . For our models it is possible to find a solution for  $^{26}\text{Al}$  and  $^{41}\text{Ca}$  for models with initial masses 60-80  $M_{\odot}$  with dilution factors in the range 0.00011 to  $7.2 \times 10^{-5}$  and  $\Delta t$  around 1 Myr.


Overall, we conform that WR stars are a robust candidate site for the production of  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$  in the ESS.

### 5.3. The impact of neutron-capture rates

From Figure 7, it becomes clear that to match the Al-, Ca-, and Cl-ratios using a self-consistent delay time within the scenario and the ESS ratios discussed in this work, more  $^{36}\text{Cl}$  and/or less  $^{41}\text{Ca}$  is required. To look more closely at this, we performed a sensitivity study for the destruction of these two isotopes. The dominant rate in the  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  destruction via neutron-captures are the  $^{36}\text{Cl}(n,p)^{36}\text{S}$  and  $^{41}\text{Ca}(n,\alpha)^{38}\text{Ar}$  rates, respectively. For the rate  $^{36}\text{Cl}(n,p)^{36}\text{S}$  we use the reaclib label “ths8” which corresponds to a theoretical determination by Rauscher (2008). For  $^{36}\text{Cl}(n,p)^{36}\text{S}$  reported experimental rates are both similar to the value used here (de Smet et al. 2007) and 30% lower, at the temperature of interest here around 200-300 MK. For the  $^{41}\text{Ca}(n,\alpha)^{38}\text{Ar}$  rate the JINA reaclib reference is Sevier et al. (1986), which is roughly a factor of two higher than the theoretical rate by Rauscher (2008). Given that there are significant differences between the current estimates, we computed three additional 60  $M_{\odot}$  non-rotating models for which we multiplied the neutron-capture reaction rates of interest by different constants as indicated in Table 3.

The table also gives the yields for  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$ . The  $^{36}\text{Cl}$  yield increases when the reaction rate is decreased, as was expected (models 2 and 4). For model 2, the  $^{41}\text{Ca}$  yield increases slightly. For models 3 and 4 the  $^{41}\text{Ca}$  yield decreases by about a factor 2, as is expected as well. The changes in the yields are not fully linear due to other (neutron-capture) reactions also playing a minor role in the destruction of the these two isotopes. If we apply the delay time calculation to these new models, we find that for model 1, the difference in the delay time between the Ca-ratio and the Cl-ratio is a factor of  $\sim 3$ . For model 2, the delay time difference is similar to that of model 1. This is visible in Figure 8, where the solid lines (model 1) and the dashed lines (model 2) are more or less overlapping. Models 3 (dotted lines)





**Figure 7.** Abundance ratios (R/S) for the three SLRs (R) over their stable reference isotope (S) for a certain dilution factor  $f$ . The solid lines represent the non-rotating models, the dashed lines the models with an initial rotational velocity of 150 km/s, and the dotted lines the models with an initial rotational velocity of 300 km/s. The horizontal bands represent the ESS ratios, with their respective errors. The vertical lines represent the delay time for the Ca-ratio. The left panels give the results for our models (40, 60, and 80  $M_{\odot}$ ), the right panels give the results for the models by LC18 for the same masses. The range of values of  $f$  for each panel (in units of  $10^{-4}$ ) are: a) 1.32-2.64, b) 2.44-3.03, c) 0.60-1.13, d) 0.54-1.37, e) 0.16-0.5, and f) 0.08-0.24.

**Table 3.** Factors used to multiply the indicated reaction rates from their standard values in the 60  $M_{\odot}$  non-rotating models considered in this section and the yields in  $M_{\odot}$  resulting from these changes.


|         | $^{36}\text{Cl}(n,p)^{36}\text{S}$ | $^{41}\text{Ca}(n,\alpha)^{38}\text{Ar}$ | $^{36}\text{Cl} (M_{\odot})$ | $^{41}\text{Ca} (M_{\odot})$ |
|---------|------------------------------------|--|------------------------------|------------------------------|
| Model 1 | 1                                  | 1  | 1.48e-06                     | 2.42e-06                     |
| Model 2 | 0.5                                | 1  | 2.47e-06                     | 2.55e-06                     |
| Model 3 | 1                                  | 2  | 1.51e-06                     | 1.29e-06                     |
| Model 4 | 0.5                                | 2  | 2.41e-06                     | 1.29e-06                     |

with  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  ratios at most roughly 8 and 20 times higher than solar. Therefore, when we add the O isotopic wind yields of the selected models diluted by  $f$  to the O solar system abundances, we obtained a decrease of the order of 0.1 to 1.5% in both the  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  ratios, increasing the absolute value with increasing the stellar mass from 40-50  $M_{\odot}$  to 80  $M_{\odot}$ . While variations at the lower values of this range would not be detectable, those at the upper values would be. Therefore, if we assume the specific scenario where the  $^{26}\text{Al}$ -poor and  $^{26}\text{Al}$ -rich grains formed before and after injection, respectively, then the lower masses would be favoured to avoid changes in the O ratios. However, the higher masses reproduce the  $^{36}\text{Cl}$  abundance. This discrepancy needs to be further investigated.

6. CONCLUSION

We have investigated the production of selected stable and radioactive isotopes in the winds of massive rotating stars. The selected SLRs of interest for the ESS are:  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$ , while the selected stable isotopes are:  $^{19}\text{F}$  and  $^{22}\text{Ne}$ . These results have been compared to various studies in the literature. We also determined which of these models could self-consistently explain the ESS abundances of  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{41}\text{Ca}$ . We have found that:

- For the SLRs, it is mostly the WRs in the mass-range 40-80  $M_{\odot}$  that give significant yields. However, the  $^{60}\text{Fe}$  yields are insignificant compare to the supernova yields.
- Only our most massive rotating stars produce a net positive  $^{19}\text{F}$  yield ( $\geq 60 M_{\odot}$  for an initial rotation rate of 300 km/s). For  $^{22}\text{Ne}$ , more stars give a net positive yield, from  $\geq 40$ -45  $M_{\odot}$ , depending on the initial rotational velocity.
- The main effect of rotation is that it lowers the initial mass for which the stars become WRs. For  $^{26}\text{Al}$  the effect of rotation on the yields is minimal, and only noticeable around the WR limit. For  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  a higher rotation rate leads to an increase in the yields at lower masses, shifting from  $\sim 30 M_{\odot}$  to  $\sim 20 M_{\odot}$ . From  $\sim 45 M_{\odot}$  the yields become again comparable for all models. For the stable isotopes, the rotational mixing leads to lower yields below 50  $M_{\odot}$  and 35  $M_{\odot}$  for  $^{19}\text{F}$  and  $^{22}\text{Ne}$ , respectively.
- Overall, the yields from our models compare well to those from the literature. There are some differences caused by a different prescription of the mass loss and/or a different approach to rotational



**Figure 8.** Results for the reaction rate tests. The colours and bands are the same as in Figure 7.

and 4 (dashed-dotted lines), provide a better solution as the difference in the delay time is reduced to a factor 1.8. There is little difference between these two models. From this, we can conclude that to obtain a better match for all three SLR-ratios, a decrease in the amount of  $^{41}\text{Ca}$  has more impact than the increase in the amount of  $^{36}\text{Cl}$ .

5.4. Oxygen isotopic ratios

For sake of completeness, we also checked if the models that reproduce  $^{26}\text{Al}$  and  $^{41}\text{Ca}$  affect the O isotopic ratios. This is because a successful pollution model should avoid predicting a correlation between the presence of  $^{26}\text{Al}$  and modification of the O isotopes (Gounelle & Meibom 2007) to match the observational evidence that some CAIs (the Fractionated and Unknown Nuclear anomalies, FUN, class) and some corundum grains are poor in  $^{26}\text{Al}$ , but they have virtually the same O isotopic composition as those that are rich in  $^{26}\text{Al}$  (see, e.g., Makide et al. 2011). The composition of the O isotopes in the winds is dominated by H and He burning and its main features are production of  $^{16}\text{O}$  and depletion of  $^{17}\text{O}$  and  $^{18}\text{O}$ , relative to their initial amounts in the star,

mixing. This clearly shows that the treatment of stellar winds and the increase of mass-loss due to rotation, as well as the treatment of rotation and rotational mixing, have still a large impact on the yields.

- In Section 5, we have investigated which of the stellar models described in this paper could explain the ESS abundances. Depending on the initial rotational velocity, stars with an initial mass of 40-45  $M_{\odot}$  and higher could explain the  $^{26}\text{Al}$  and  $^{41}\text{Ca}$  abundances. However, only the most massive models ( $\geq 60 M_{\odot}$ ) can also explain the  $^{36}\text{Cl}$  abundances. We remind that also the following CCSNe of massive star models will expel a significant amount of these isotopes, however, they produce an overabundance of  $^{60}\text{Fe}$  relatively to its ESS value.
- From our tests of the neutron-capture rates of  $^{36}\text{Cl}$  and  $^{41}\text{Ca}$  we conclude that to obtain a better match for all three SLR-ratios, a decrease in the amount of  $^{41}\text{Ca}$ , derived from increasing its (n, $\alpha$ ) rate, has more impact than the increase in the amount of  $^{36}\text{Cl}$ , derived from decreasing its (n,p) rate.
- When comparing our models with the oxygen-ratios in the Solar System, which are known to high precision, however, we find that the high mass models decrease the oxygen isotopic ratios,  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$ , too much, while the lower mass models stay within the error-margins of the measurements.

For all isotopes discussed here, except for  $^{26}\text{Al}$  which we presented in Paper I, the influence of binary interactions still needs to be investigated. We will examine this in our upcoming paper, where we will also calculate whether these binary yields could match the ESS abundances. Future work includes the formation of dust from

the WR binary stars (see, e.g., Lau et al. 2020), since dust may be needed to incorporate SLRs into the ESS (Dwarkadas et al. 2017). Furthermore, a more detailed analysis of several uncertainties should be performed, which includes different prescriptions for the winds and for the rotational boost on wind loss, as well as investigations of the effect of reaction rate uncertainties specifically on the destruction of  $^{19}\text{F}$ ,  $^{41}\text{Ca}$ , and  $^{36}\text{Cl}$ , and the neutron source  $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$  reaction (Adsley et al. 2021). Finally, to present a complete view of the isotopes discussed here, the explosive nucleosynthetic yields will need to be calculated using our models as the progenitors of the explosion.

#### ACKNOWLEDGEMENTS

HEB thanks the MESA team for making their code publicly available, Frank Timmes for his support with the pre-supernova models, and Bill Paxton for other clarifications of the code. We thank Zsolt Keszthelyi for his help with implementing the rotational boost into MESA. We thank Alessandro Chieffi and Marco Limongi for clarifications of their results in their 2018 paper. We also thank Alan Boss for discussion on the ESS, and Andrés Yague for his assistance with Python. We also thank the referee for all their comments and help with improving this paper. This work is supported by the ERC via CoG-2016 RADIOSTAR [11] (Grant Agreement 724560). We also acknowledge support from “ChETEC” COST Action(CA16117), supported by COST (European Cooperation in Science and Technology). MP acknowledges significant support to NuGrid from STFC (through the University of Hull’s Consolidated Grant ST/R000840/1), from the National Science Foundation (NSF, USA) under grant No. PHY-1430152 (JINA Center for the Evolution of the Elements) and from the “Lendulet-2014” Program of the Hungarian Academy of Sciences (Hungary). MP thanks the support from the US IRENA Accelnet network.

*Software:* MESA (Paxton et al. 2011, 2013, 2015, 2018)

#### REFERENCES

- Abia, C., Cristallo, S., Cunha, K., de Laverny, P., & Smith, V. V. 2019, A&A, 625, A40
- Adams, F. C. 2010, ARA&A, 48, 47
- Adsley, P., Battino, U., Best, A., et al. 2021, PhRvC, 103, 015805
- Aerts, C., Mathis, S., & Rogers, T. M. 2019, ARA&A, 57, 35
- Arnould, M., Goriely, S., & Meynet, G. 2006, A&A, 453, 653
- Arnould, M., Paulus, G., & Meynet, G. 1997, A&A, 321, 452
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481
- Austin, S. M., West, C., & Heger, A. 2017, ApJL, 839, L9
- Basunia, M., & Hurst, A. 2016, Nuclear Data Sheets, 134, 75
- Belczynski, K., Klencki, J., Fields, C. E., et al. 2020, A&A, 636, A104

- Brinkman, H. E., Doherty, C. L., Pols, O. R., et al. 2019, *ApJ*, 884, 38
- Chieffi, A., & Limongi, M. 2013, *ApJ*, 764, 21
- Côté, B., Lugaro, M., Reifarth, R., et al. 2019a, *ApJ*, 878, 156
- Côté, B., Yagüe, A., Világos, B., & Lugaro, M. 2019b, *ApJ*, 887, 213
- Cyburt, R. H., Amthor, A. M., Ferguson, R., et al. 2010, *ApJS*, 189, 240
- de Smet, L., Wagemans, C., Goeminne, G., Heyse, J., & van Gils, J. 2007, *PhRvC*, 75, 034617
- Diehl, R. 2013, *Reports on Progress in Physics*, 76, 026301
- Dwarkadas, V. V., Dauphas, N., Meyer, B., Boyajian, P., & Bojazi, M. 2017, *ApJ*, 851, 147
- Ekström, S., Georgy, C., Eggenberger, P., et al. 2012, *A&A*, 537, A146
- Farmer, R., Fields, C. E., Petermann, I., et al. 2016, *ApJS*, 227, 22
- Gaidos, E., Krot, A. N., Williams, J. P., & Raymond, S. N. 2009, *ApJ*, 696, 1854
- Götberg, Y., de Mink, S. E., Groh, J. H., et al. 2018, *A&A*, 615, A78
- Gounelle, M., & Meibom, A. 2007, *ApJL*, 664, L123
- Gounelle, M., & Meynet, G. 2012, *A&A*, 545, A4
- Hamann, W.-R., Koesterke, L., & Wessolowski, U. 1995, *A&A*, 299, 151
- Heger, A., Langer, N., & Woosley, S. E. 2000, *ApJ*, 528, 368
- Heger, A., Woosley, S. E., & Spruit, H. C. 2005, *ApJ*, 626, 350
- Imbriani, G., Costantini, H., Formicola, A., et al. 2005, *European Physical Journal A*, 25, 455
- Jacobsen, B., Yin, Q.-z., Moynier, F., et al. 2008, *Earth and Planetary Science Letters*, 272, 353
- Jones, S. W., Möller, H., Fryer, C. L., et al. 2019, *MNRAS*, 485, 4287
- Jönsson, H., Ryde, N., Harper, G. M., Richter, M. J., & Hinkle, K. H. 2014a, *ApJL*, 789, L41
- Jönsson, H., Ryde, N., Spitoni, E., et al. 2017, *ApJ*, 835, 50
- Jönsson, H., Ryde, N., Harper, G. M., et al. 2014b, *A&A*, 564, A122
- Keszthelyi, Z., Meynet, G., Shultz, M. E., et al. 2020, *MNRAS*, 493, 518
- Lamers, H. J. G. L. M., Snow, T. P., & Lindholm, D. M. 1995, *ApJ*, 455, 269
- Langer, N. 1998, *A&A*, 329, 551
- Lau, R. M., Eldridge, J. J., Hankins, M. J., et al. 2020, *ApJ*, 898, 74
- Limongi, M., & Chieffi, A. 2006, *ApJ*, 647, 483
- . 2018, *ApJS*, 237, 13
- Liu, M.-C. 2017, *GeoCoA*, 201, 123
- Lodders, K. 2003, *ApJ*, 591, 1220
- Lugaro, M., Ott, U., & Kereszturi, Á. 2018, *Progress in Particle and Nuclear Physics*, 102, 1
- Luu, T.-H., Hin, R. C., Coath, C. D., & Elliott, T. 2019, *Earth and Planetary Science Letters*, 522, 166
- Maeder, A., & Meynet, G. 2000a, *A&A*, 361, 159
- . 2000b, *ARA&A*, 38, 143
- . 2012, *Reviews of Modern Physics*, 84, 25
- Maeder, A., & Zahn, J.-P. 1998, *A&A*, 334, 1000
- Makide, K., Nagashima, K., Krot, A. N., et al. 2011, *ApJL*, 733, L31
- Meyer, B. S., & Clayton, D. D. 2000, *SSRv*, 92, 133
- Meynet, G., & Arnould, M. 2000, *A&A*, 355, 176
- Murray, N. 2011, *ApJ*, 729, 133
- Nesaraja, C., & McCutchan, E. 2016, *Nuclear Data Sheets*, 133, 120
- Nica, N., Cameron, J., & Singh, J. 2012, *Nuclear Data Sheets*, 113, 25
- Nieuwenhuijzen, H., & de Jager, C. 1990, *A&A*, 231, 134
- Nugis, T., & Lamers, H. J. G. L. M. 2000, *A&A*, 360, 227
- O'Connor, E., & Ott, C. D. 2011, *ApJ*, 730, 70
- Palacios, A., Arnould, M., & Meynet, G. 2005, *A&A*, 443, 243
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, *ApJS*, 208, 4
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, *ApJS*, 220, 15
- Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, *ApJS*, 234, 34
- Pignatari, M., Herwig, F., Hirschi, R., et al. 2016, *VizieR Online Data Catalog*, J/ApJS/225/24
- Prantzos, N. 2012, *A&A*, 538, A80
- Rauscher, T. 2008, *Online code NON-SMOKERWEB, version 5.0w and higher*
- Ritter, C., Herwig, F., Jones, S., et al. 2018, *MNRAS*, 480, 538
- Rugel, G., Faestermann, T., Knie, K., et al. 2009, *PhRvL*, 103, 072502
- Ryde, N., Jönsson, H., Mace, G., et al. 2020, *ApJ*, 893, 37
- Schneider, F. R. N., Podsiadlowski, P., & Müller, B. 2021, *A&A*, 645, A5
- Sevier, M., Mitchell, L., Tingwell, C., & Sargood, D. 1986, *Nuclear Physics A*, 454, 128
- Stancliffe, R. J., Lugaro, M., Ugalde, C., et al. 2005, *MNRAS*, 360, 375
- Sukhbold, T., Ertl, T., Woosley, S. E., Brown, J. M., & Janka, H.-T. 2016, *ApJ*, 821, 38

- Tang, H., & Dauphas, N. 2012, *Earth and Planetary Science Letters*, 359, 248
- Tang, H., Liu, M.-C., McKeegan, K. D., Tissot, F. L. H., & Dauphas, N. 2017, *GeoCoA*, 207, 1
- Trappitsch, R., Boehnke, P., Stephan, T., et al. 2018, *ApJL*, 857, L15
- Tur, C., Heger, A., & Austin, S. M. 2010, *ApJ*, 718, 357
- Ugalde, C., Azuma, R. E., Couture, A., et al. 2008, *PhRvC*, 77, 035801
- Vink, J. S., & de Koter, A. 2005, *A&A*, 442, 587
- Vink, J. S., de Koter, A., & Lamers, H. J. G. L. M. 2000, *A&A*, 362, 295
- . 2001, *A&A*, 369, 574
- Wang, W., Siegert, T., Dai, Z. G., et al. 2020, *ApJ*, 889, 169
- Wasserburg, G. J., Busso, M., Gallino, R., & Nollett, K. M. 2006, *NuPhA*, 777, 5
- Young, E. D. 2014, *Earth and Planetary Science Letters*, 392, 16