

2D coordination polymers based on the interaction between cucurbit[5]uril and light lanthanide metal ions in the presence of copper salts, and use for the detection of nitrogenous bases

Lin Huang ^a, Jie Li ^a, Ming-Hui Zhang ^a, Zhao-Qiang Zhu ^a, Kai Chen, ^{a,*} and Carl Redshaw ^b

^a Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, P. R. China. Email: kaichen85@nuist.edu.cn or catQchen@163.com (K. Chen)

^b Chemistry, School of Natural Sciences, University of Hull, Hull HU6 7RX, UK.




Fig. S1 The fit curve of A and U.







Fig. S2 Fluorescence quenching and recyclability test for 1.



(a)



(b)



(c)

Fig. S3. Fluorescence enhancement of 1 dispersed in water upon addition of T(a), C(b), and G(c).

Table S1. Standard deviation and detection limit calculation for A and U.

	A	U
	874.0613	858.1628
	872.4453	852.3751
NO.	871.1987	860.4659
	873.6598	853.9072
	875.4375	859.0945
Standard deviation(σ)	1.442606	3.114764
Slope (m)	14.2	12.7
Detection limit ($3\sigma/m$, mM)	0.304776	0.340773