

Short report

# DNA methylation changes in CD4<sup>+</sup> T cells isolated from multiple sclerosis patients on dimethyl fumarate

Vicki E Maltby , Rodney A Lea, Karen A Ribbons, Katherine A Sanders, Daniel Kennedy, Myintzu Min, Rodney J Scott and Jeannette Lechner-Scott

Multiple Sclerosis Journal—  
Experimental, Translational  
and Clinical

July–September 2018, 1–6

DOI: 10.1177/  
2055217318787826

© The Author(s), 2018.  
Reprints and permissions:  
[http://www.sagepub.co.uk/  
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

## Abstract

**Background:** Dimethyl fumarate is an oral treatment for multiple sclerosis, whose mechanism of action is not fully understood.

**Objective:** To investigate the effects of dimethyl fumarate on DNA methylation in the CD4<sup>+</sup> T cells of multiple sclerosis patients.

**Methods:** We performed Illumina EPIC arrays to investigate the DNA methylation profiles of CD4<sup>+</sup> T cells derived from multiple sclerosis patients before and after dimethyl fumarate treatment.

**Results:** Treatment with dimethyl fumarate resulted in 97% of differentially methylated positions showing hypermethylation. Four genes, *SNORD1A*, *SHTN1*, *MZB1* and *TNF* had a differentially methylated region located within the transcriptional start site.

**Conclusion:** This study investigates the effect of dimethyl fumarate on DNA methylation in multiple sclerosis patients.

**Keywords:** Multiple sclerosis, dimethyl fumarate, immunology, DNA methylation, relapsing–remitting, CD4<sup>+</sup> T cells, tumour necrosis factor

Date received: 6 February 2018; revised 23 April 2018; accepted: 24 May 2018

## Introduction

Although increasing numbers of treatments are available for multiple sclerosis (MS), the exact mechanism of action of these therapies is often unclear. Patients are frequently required to trial several treatments to identify which is most suitable for their disease activity. Dimethyl fumarate (DMF; Tecfidera, Biogen Idec, Cambridge MA, USA) is approved in Europe and Australia as a first-line oral drug for the treatment of relapsing–remitting multiple sclerosis, and its use is associated with a reduction in disease activity and a variable effect on progression.<sup>1,2</sup>

Although the exact mode of action is not fully elucidated, DMF has been shown to have both anti-inflammatory and anti-oxidative properties. Decreased absolute lymphocyte counts and a shift

in T lymphocyte polarisation from T helper (Th)1 and Th17 (pro-inflammatory) to Th2 phenotype (anti-inflammatory) has been reported after DMF treatment in MS patients.<sup>3</sup> DMF also promotes translocation of nuclear factor erythroid 2-related factor 2 into the nucleus, which upregulates the transcription of anti-oxidative enzymes.<sup>3</sup>

DNA methylation refers to the epigenetic modification whereby the addition/removal of methyl groups to CpG dinucleotides regulates gene transcription. We, and others, have assessed global methylation profiles in CD4<sup>+</sup> and CD8<sup>+</sup> T cells from MS patients compared to healthy controls.<sup>4–6</sup> Our studies have demonstrated altered methylation profiles in the CD4<sup>+</sup> T cells of treatment-naïve patients or in the absence of treatment. However, the effect of disease-modifying therapies (DMTs) on methylation

Correspondence to:  
**Jeannette Lechner-Scott**,  
Department of Neurology,  
John Hunter Hospital,  
Australia.  
[jeannette.lechner-scott@  
hnehealth.sw.gov.au](mailto:jeannette.lechner-scott@hnehealth.sw.gov.au)

**Vicki E Maltby**,  
School of Medicine and  
Public Health, University of  
Newcastle, Australia  
Brain and Mental Health,  
Hunter Medical Research  
Institute, Australia

**Rodney A Lea**,  
Brain and Mental Health,  
Hunter Medical Research  
Institute, Australia  
Institute of Health and  
Biomedical Innovation,  
Queensland University of  
Technology, Australia

**Karen A Ribbons**,  
Brain and Mental Health,  
Hunter Medical Research



Institute, Australia  
Department of Neurology,  
John Hunter  
Hospital, Australia

**Katherine A Sanders,**  
Faculty of Health Sciences  
and Medicine, Bond  
University, Australia

**Daniel Kennedy,**  
ARC Centre of Excellence  
for the Mathematical and  
Statistical Frontiers,  
Queensland University of  
Technology,  
Brisbane, Australia

**Myintzu Min,**  
Brain and Mental Health,  
Hunter Medical Research  
Institute, Australia  
Department of Neurology,  
John Hunter  
Hospital, Australia

**Rodney J Scott,**  
Brain and Mental Health,  
Hunter Medical Research  
Institute, Australia  
School of Biomedical  
Sciences and Pharmacy,  
University of Newcastle,  
Australia  
Department of Medical  
Genetics, John Hunter  
Hospital, Australia

**Jeannette Lechner-Scott,**  
School of Medicine and  
Public Health, University of  
Newcastle, Australia  
Brain and Mental Health,  
Hunter Medical Research  
Institute, Australia  
Department of Neurology,  
John Hunter  
Hospital, Australia

profiles remains unclear. Neither group found significant changes in CD8<sup>+</sup> T cells.<sup>5,6</sup>

Here we performed a longitudinal study of the genome-wide methylation profiles of CD4<sup>+</sup> T cells in MS patients before and after DMF treatment.

## Methods

We recruited seven MS patients (three men and four women) who were either treatment naive or had been off DMT for at least 3 months and were planning to start DMF therapy (Table 1). The majority of patients had not had steroid use for at least 2 months prior to entry into this study (Table 1). Samples were collected and processed as previously described.<sup>7</sup> Blood was collected prior to the first dose of DMF and 6 months following treatment initiation. At 6 months, all patients remained on therapy and had no change in Expanded Disability Status Score (EDSS). Two patients had evidence of disease activity as assessed by the appearance of new lesions on magnetic resonance imaging (MRI). However, both of these patients showed no new disease activity at their next MRI and remain on therapy.

CD4<sup>+</sup> T cells were extracted using magnetic isolation kits (Stem Cell Technologies, Canada) and purity (minimum threshold  $\geq 90\%$ ) was assessed using the FACS CantoII (BD Biosciences) system. Purified DNA was bisulphite converted and hybridised to Illumina EPIC arrays. Raw fluorescence data were processed using a combination of R/Bioconductor and custom scripts. Differences in mean methylation before and after the 6-month treatment period were tested using a paired samples *t*-test for each CpG. A CpG was considered a differentially methylated position (DMP) if the *P* value was less

than 0.0005 and the absolute difference in mean methylation between groups was greater than 5%. A differentially methylated region (DMR) was defined as two or more contiguous DMPs located within 500 bp of each other, whose methylation changes were in the same direction. If a DMP was located outside of the 500 bp region but was less than 500 bp from the last DMP it was also included in the DMR.

## Results and discussion

In total, 945 DMPs were identified when comparing the 6-month time point to baseline, the majority of which were hypermethylated after treatment (912; 97%) (see Supplementary Table 1). The most altered DMP between baseline and treatment was 17.5% hypermethylated (cg14048158); however, this site maps to an area with no known gene association. To identify sites of potential functional consequence, we filtered DMPs to include only those with a DMR, gene name and position annotation. Table 2 shows the DMPs with the largest percentage change for each of the resulting 64 genes.

Four genes had at least two adjacent DMPs located in the transcriptional start site (TSS) (Table 3). *SNORD1A* (small nucleolar RNA, C/D box 1A) encodes for an uncharacterised small nucleolar RNA. *SHTN1* encodes shootin1, a protein involved in neuronal polarisation of axons.<sup>8</sup> *MZBI* (marginal zone B and B1 cell-specific protein) codes for an endoplasmic reticulum calcium regulator. While it has not previously been linked to MS, a study by Belkaya et al. (2013) found that overexpression of miR-185 resulted in a nearly five-fold decrease of *MZBI* in mice.<sup>9</sup> This decrease corresponded with lymphopenia and a reduced proliferative response in CD4<sup>+</sup> T

**Table 1.** MS cohort demographic/clinical features at baseline (prior to DMF treatment).

Sex	Age (years)	Prior DMT	Prior steroids (days prior to collection)	EDSS at baseline	EDSS at 6 months
M	23	Naive	312	1	1
M	40	Naive	7	3	3
M	35	Naive	54	1.5	1.5
F	32	Naive	3400	2.5	2.5
F	42	(Fingolimod)	1224	3	3
F	31	(Glatiramer acetate)	72	1.5	1.5
F	43	(Peginterferon beta-1a, interferon beta-1a)	95	2	2

MS: multiple sclerosis; DMF: dimethyl fumarate; DMT: disease-modifying therapy; EDSS: Expanded Disability Scale Status.

**Table 2.** DMRs with gene name and annotation.

Chr.	CpG ID	Position	Gene name	Element	Mean methylation			T stat.	P value
					Baseline	6 Months	% Change		
1	cg16144718	23115066	<i>EPHB2</i>	Body	0.50	0.62	11.47	7.06	$4.05 \times 10^{-4}$
1	cg06808725	32264502	<i>SPOCD1</i>	Body	0.44	0.56	11.74	6.64	$5.64 \times 10^{-4}$
1	cg24533227	42145514	<i>HIVEP3</i>	5'UTR	0.65	0.76	10.63	6.71	$5.32 \times 10^{-4}$
1	cg02410801	55046065	<i>ACOT11</i>	Body	0.56	0.66	9.02	8.87	$1.15 \times 10^{-4}$
1	cg25130912	201982886	<i>ELF3</i>	Body	0.66	0.76	9.67	7.11	$3.89 \times 10^{-4}$
2	cg05333614	1168186	<i>SNTG2</i>	Body	0.72	0.77	5.36	7.77	$2.40 \times 10^{-4}$
2	cg03771015	15831147	<i>LOC101926966</i>	Body	0.62	0.70	7.78	6.35	$7.16 \times 10^{-4}$
2	cg14501323	31279457	<i>GALNT14</i>	Body	0.81	0.87	6.37	6.96	$4.38 \times 10^{-4}$
2	cg10796691	65135899	<i>LOC400958</i>	Body	0.60	0.69	8.36	9.16	$9.53 \times 10^{-5}$
2	cg16603943	121614683	<i>GLI2</i>	Body	0.51	0.63	11.66	6.35	$7.13 \times 10^{-4}$
2	cg20772458	158983130	<i>UPP2</i>	Body	0.74	0.80	5.89	7.26	$3.46 \times 10^{-4}$
2	cg18707238	218688237	<i>TNS1</i>	Body	0.73	0.78	5.52	6.47	$6.49 \times 10^{-4}$
3	cg15756415	14932169	<i>FGD5</i>	Body	0.42	0.54	12.05	6.48	$6.40 \times 10^{-4}$
3	cg02790932	23373256	<i>UBE2E2</i>	Body	0.65	0.72	7.55	8.65	$1.31 \times 10^{-4}$
3	cg00049674	123058535	<i>ADCY5</i>	Body	0.61	0.68	7.51	7.01	$4.20 \times 10^{-4}$
5	cg27073488	14262157	<i>TRIO</i>	Body	0.70	0.75	5.39	8.81	$1.19 \times 10^{-4}$
5	cg16375820	55289001	<i>IL6ST</i>	5'UTR	0.31	0.23	-7.30	-9.25	$9.03 \times 10^{-5}$
5	cg27346756	90431802	<i>ADGRV1</i>	Body	0.58	0.64	6.52	7.38	$3.18 \times 10^{-4}$
5	cg16558774	132579360	<i>FSTL4</i>	Body	0.68	0.75	6.66	9.33	$8.58 \times 10^{-5}$
5	cg11988321	138725622	<i>MZB1</i>	TSS200	0.42	0.54	12.21	16.30	$3.39 \times 10^{-6}$
6	cg04095776	31106941	<i>PSORS1C1</i>	Body	0.66	0.72	6.25	7.34	$3.28 \times 10^{-4}$
6	cg19978379	31542671	<i>TNF</i>	TSS1500	0.54	0.67	13.00	7.09	$3.95 \times 10^{-4}$
6	cg15496866	40491590	<i>LRFN2</i>	5'UTR	0.61	0.72	11.04	7.47	$2.97 \times 10^{-4}$
6	cg01473948	148823785	<i>SASH1</i>	Body	0.59	0.66	7.47	6.91	$4.54 \times 10^{-4}$
7	cg13800949	47343103	<i>TNS3</i>	Body	0.79	0.85	5.90	8.14	$1.85 \times 10^{-4}$
7	cg14797899	69882555	<i>AUTS2</i>	Body	0.68	0.78	9.85	7.70	$2.51 \times 10^{-4}$
7	cg02170577	104939331	<i>SRPK2</i>	Body	0.72	0.77	5.03	6.94	$4.44 \times 10^{-4}$
7	cg05476934	133859100	<i>LRGUK</i>	Body	0.52	0.60	8.64	6.74	$5.18 \times 10^{-4}$
7	cg09891341	138619424	<i>KIAA1549</i>	Body	0.78	0.84	6.24	6.81	$4.90 \times 10^{-4}$
7	cg06679384	158049077	<i>PTPRN2</i>	Body	0.60	0.66	6.23	7.57	$2.76 \times 10^{-4}$
9	cg08290373	8633541	<i>PTPRD</i>	Body	0.68	0.78	10.02	6.66	$5.52 \times 10^{-4}$
9	cg17557530	90193634	<i>DAPK1</i>	Body	0.61	0.73	12.22	6.62	$5.74 \times 10^{-4}$
9	cg06749278	97662692	<i>C9orf3</i>	Body	0.75	0.83	7.50	6.91	$4.55 \times 10^{-4}$
10	cg16203213	45398814	<i>TMEM72-AS1</i>	Body	0.66	0.74	7.24	9.01	$1.05 \times 10^{-4}$
10	cg26754789	49857879	<i>ARHGAP22</i>	Body	0.74	0.79	5.36	9.19	$9.34 \times 10^{-5}$
10	cg13312268	50019744	<i>WDFY4</i>	ExonBnd	0.72	0.78	6.84	6.68	$5.44 \times 10^{-4}$
10	cg12552633	71573337	<i>COL13A1</i>	Body	0.45	0.55	10.40	6.72	$5.29 \times 10^{-4}$
10	cg24587741	79313774	<i>KCNMA1</i>	Body	0.68	0.75	7.07	6.89	$4.62 \times 10^{-4}$
10	cg17753789	81026766	<i>ZMIZ1</i>	Body	0.69	0.76	7.74	6.52	$6.21 \times 10^{-4}$
10	cg16035098	118886914	<i>SHTN1</i>	TSS1500	0.46	0.55	9.16	9.66	$7.04 \times 10^{-5}$
10	cg01613414	126693304	<i>CTBP2</i>	Body	0.52	0.62	10.31	7.40	$3.13 \times 10^{-4}$
11	cg09731767	503628	<i>RNH1</i>	5'UTR	0.53	0.61	7.79	9.85	$6.32 \times 10^{-5}$
11	cg11922498	4936427	<i>OR51G2</i>	1stExon	0.65	0.71	6.09	9.45	$7.99 \times 10^{-5}$
11	cg00842359	10686144	<i>MRVII</i>	5'UTR	0.67	0.77	9.37	9.28	$8.85 \times 10^{-5}$
11	cg14595291	35993855	<i>LDLRAD3</i>	5'UTR	0.50	0.65	14.89	6.42	$6.76 \times 10^{-4}$
11	cg00964019	117593395	<i>DSCAML1</i>	Body	0.76	0.82	5.81	7.08	$3.97 \times 10^{-4}$
12	cg11439695	2561024	<i>CACNA1C</i>	Body	0.46	0.56	9.93	8.25	$1.71 \times 10^{-4}$
12	cg17451712	122293122	<i>HPD</i>	Body	0.68	0.75	7.13	6.57	$5.95 \times 10^{-4}$

(continued)

**Table 2.** Continued

Chr.	CpG ID	Position	Gene name	Element	Mean methylation			T stat.	P value
					Baseline	6 Months	% Change		
14	cg03725784	61992305	<i>PRKCH</i>	Body	0.41	0.54	12.42	7.14	$3.81 \times 10^{-4}$
14	cg11198334	75040680	<i>LTBP2</i>	Body	0.63	0.74	10.81	6.90	$4.56 \times 10^{-4}$
14	cg07399096	91050031	<i>TTC7B</i>	Body	0.69	0.75	6.38	6.58	$5.91 \times 10^{-4}$
14	cg15325186	102562217	<i>HSP90AA1</i>	Body	0.51	0.60	8.52	7.79	$2.36 \times 10^{-4}$
15	cg25814224	51572976	<i>CYP19A1</i>	5'UTR	0.65	0.71	6.39	10.93	$3.49 \times 10^{-5}$
16	cg02260059	78262124	<i>WWOX</i>	Body	0.72	0.78	5.78	6.62	$5.74 \times 10^{-4}$
17	cg04456720	54250143	<i>ANKFN1</i>	Body	0.67	0.77	10.60	8.93	$1.10 \times 10^{-4}$
17	cg19439071	74557625	<i>SNORD1A</i>	TSS200	0.56	0.67	10.87	8.68	$1.29 \times 10^{-4}$
17	cg11476241	78866235	<i>RPTOR</i>	Body	0.56	0.66	10.10	6.45	$6.58 \times 10^{-4}$
18	cg13297582	13288627	<i>LDLRAD4</i>	5'UTR	0.51	0.62	10.73	7.79	$2.36 \times 10^{-4}$
18	cg03385871	46311648	<i>CTIF</i>	Body	0.45	0.56	11.75	6.56	$6.00 \times 10^{-4}$
19	cg07345937	1175444	<i>SBNO2</i>	TSS1500	0.53	0.63	9.74	10.58	$4.19 \times 10^{-5}$
20	cg10453816	37499530	<i>PPP1R16B</i>	Body	0.51	0.64	13.41	6.85	$4.74 \times 10^{-4}$
20	cg04991444	50057438	<i>NFATC2</i>	Body	0.41	0.53	12.25	9.96	$5.94 \times 10^{-5}$
21	cg10919441	44143035	<i>PDE9A</i>	5'UTR	0.67	0.73	6.59	7.18	$3.68 \times 10^{-4}$

DMT: disease-modifying therapy; Chr.: chromosome.

**Table 3.** Genes with DMRs in the transcriptional start site.

Chr.	CpG ID	Position	Gene name	Element	Mean methylation			T stat.	P value
					Baseline	6 months	% Change		
5	cg11988321	138725622	<i>MZB1</i>	TSS200	0.421426	0.543513	12.20872	16.29927	$3.39 \times 10^{-6}$
5	cg04359635	138725975	<i>MZB1</i>	TSS1500	0.513568	0.633291	11.97221	6.619687	$5.72 \times 10^{-4}$
6	cg19978379	31542671	<i>TNF</i>	TSS1500	0.537903	0.667883	12.99794	7.088953	$3.95 \times 10^{-4}$
6	cg24452282	31542740	<i>TNF</i>	TSS1500	0.470376	0.584539	11.41626	6.396426	$6.88 \times 10^{-4}$
10	cg16035098	118886914	<i>SHTN1</i>	TSS1500	0.458233	0.549798	9.156483	9.664102	$7.04 \times 10^{-5}$
10	cg23251794	118886883	<i>SHTN1</i>	TSS1500	0.641729	0.707866	6.613676	9.553839	$7.51 \times 10^{-5}$
17	cg19439071	74557625	<i>SNORD1A</i>	TSS200	0.562719	0.671405	10.86864	8.67833	$1.29 \times 10^{-4}$
17	cg07180212	74557703	<i>SNORD1A</i>	TSS200	0.62678	0.726408	9.962796	6.422776	$6.73 \times 10^{-4}$
17	cg13664588	74557494	<i>SNORD1A</i>	TSS1500	0.527718	0.624507	9.678917	6.578915	$5.92 \times 10^{-4}$

DMR: differentially methylated region; Chr.: chromosome.

cells.<sup>9</sup> The observed increase in DNA methylation identified in the *MZB1* TSS in our dataset may result in a similar decrease in *MZB1* transcription. A resulting decrease in CD4<sup>+</sup> T cells would be consistent with the known anti-inflammatory action of DMF.

The fourth DMR identified is located at the TSS of tumour necrosis factor (*TNF*). *TNF* is a pro-inflammatory cytokine that is produced by many cell types, including lymphocytes (reviewed in Wajant *et al.*)<sup>10</sup> TNF binding to its receptor activates the nuclear factor kappa B (NF-κB) pathway, which

activates the transcription of genes involved in cell survival and proliferation, inflammatory response and anti-apoptotic factors. Hypermethylation at the *TNF* TSS may result in decreased TNF production, and a decrease in activation of the NF-κB pathway. One known mechanism of action for DMF is preventing translocation of NF-κB to the nucleus, resulting in a decrease of pro-inflammatory cytokines and an increase of anti-inflammatory cytokines (reviewed in Pistono *et al.*)<sup>3</sup> It is possible that altered DNA methylation profiles at the *TNF* TSS may contribute to this mechanism.

DMF has previously been linked to other epigenetic mechanisms in a study by Kalinin *et al.* (2013), in which they reported that DMF increased expression of histone deacetylases in cultured rat astrocytes.<sup>11</sup> Both DNA methylation and histone deacetylation are associated with gene repression.<sup>12</sup> Taken together there is now evidence that DMF may act as an epigenetic modifier with the function of shutting down transcription associated with pro-inflammatory activity.

One limitation of this study is that we only assessed patients who started DMF treatment. Also, although the majority of patients were stable at the time of baseline collection, two patients had recently had a relapse, only one of whom was treated with steroids. We are therefore unable to determine for certain if the changes in methylation profiles are due to treatment effects or stabilisation of disease. Future studies comparing changes following different therapies and different disease severities are required. A further limitation is the small sample size and lack of transcriptional data. Future studies characterising treatment responses in larger populations that also investigate the functional changes at the transcriptional level are warranted.

This is the first longitudinal study to investigate the effect of DMF on the DNA methylation of CD4<sup>+</sup> T cells of MS patients. Of the most interest, the DMRs identified at *TNF* and *MZB1* provide a potential novel mechanism of action for DMF. Treatment with DMF resulted in overall hypermethylation suggesting that DMF may act to promote DNA methylation. Larger studies are warranted to elucidate further the functional link between DMF and epigenetic mechanisms.

### Acknowledgements

The authors would like to thank the MS patients and clinical team at the John Hunter Hospital MS clinic who participated in this study. They also acknowledge the analytical biomolecular research facility at the University of Newcastle for flow cytometry support.

### Author contribution

VEM, KAS, RAL, JLS, RJS and KAR initiated and designed the original study. VEM and KAS performed all laboratory experiments. VEM wrote the final manuscript and revised all versions of the manuscript. RAL and DK performed the statistical analysis. RAL, KAS, JLS, KAR, MM and RJS helped interpret the data and critically reviewed the manuscript.

### Availability of data and material

The datasets generated or analysed during the current study are included in this published article (Supplementary Table 1). Raw data files are available from Rodney A Lea.

### Conflict on Interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: JLS's institution receives non-directed funding as well as honoraria for presentations and membership on advisory boards from Sanofi Aventis, Biogen Idec, Bayer Health Care, Merck Serono, Teva and Novartis Australia.

### Ethics approval and consent to participate

The Hunter New England health research ethics committee and University of Newcastle ethics committee approved this study (05/04/13.09 and H-505-0607, respectively), and methods were carried out in accordance with institutional guidelines on human subject experiments. Written and informed consent was obtained from all patient and control subjects.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by philanthropic contributions to the Hunter Medical Research Institute. VEM is supported by fellowships from the Canadian Institutes of Health Research and Multiple Sclerosis Research Australia (MSRA). RAL is partially funded by a fellowship from the MSRA. KAS was funded by a fellowship from the MSRA and the Trish MS research Foundation. KAR, DK, MM and JLS have no funding to declare.

### ORCID iD

Vicki E Maltby  <http://orcid.org/0000-0002-3785-4742>

### Supplemental Material

Supplementary material is available for this article online.

### References

1. Fox RJ, Miller DH, Phillips JT, *et al.* Placebo-controlled phase 3 study of oral BG-12 or glatiramer in multiple sclerosis. *N Engl J Med* 2012; 367: 1087–1097.
2. Gold R, Kappos L, Arnold DL, *et al.* Placebo-controlled phase 3 study of oral BG-12 for relapsing multiple sclerosis. *N Engl J Med* 2012; 367: 1098–1107.
3. Pistono C, Osera C, Boiocchi C, *et al.* What's new about oral treatments in multiple sclerosis? Immunogenetics still under question. *Pharmacol Res* 2017; 120: 279–293.

4. Maltby VE, Lea RA, Sanders KA, et al. Differential methylation at MHC in CD4<sup>+</sup> T cells is associated with multiple sclerosis independently of HLA-DRB1. *Clin Epigenetics* 2017; 9: 71.
5. Bos SD, Page CM, Andreassen BK, et al. Genome-wide DNA methylation profiles indicate CD8<sup>+</sup> T cell hypermethylation in multiple sclerosis. *PLoS One* 2015; 10: e0117403.
6. Maltby VE, Graves MC, Lea RA, et al. Genome-wide DNA methylation profiling of CD8<sup>+</sup> T cells shows a distinct epigenetic signature to CD4<sup>+</sup> T cells in multiple sclerosis patients. *Clin Epigenetics* 2015; 7: 118.
7. Graves M, Benton M, Lea R, et al. Methylation differences at the HLA-DRB1 locus in CD4<sup>+</sup> T-cells are associated with multiple sclerosis. *Mult Scler* 2013; 20: 1033–1041.
8. Toriyama M, Shimada T, Kim KB, et al. Shootin1: a protein involved in the organization of an asymmetric signal for neuronal polarization. *J Cell Biol* 2006; 175: 147–157.
9. Belkaya S, Murray SE, Eitson JL, et al. Transgenic expression of microRNA-185 causes a developmental arrest of T cells by targeting multiple genes including Mzb1. *J Biol Chem* 2013; 288: 30752–30762.
10. Wajant H, Pfizenmaier K and Scheurich P. Tumor necrosis factor signaling. *Cell Death Differ* 2003; 10: 45–65.
11. Kalinin S, Polak PE, Lin SX, et al. Dimethyl fumarate regulates histone deacetylase expression in astrocytes. *J Neuroimmunol* 2013; 263: 13–19.
12. Allis CD and Jenuwein T. The molecular hallmarks of epigenetic control. *Nat Rev Genet* 2016; 17: 487–500.