

Opportunistic screening for alcohol use problems in adolescents attending emergency departments: an evaluation of screening tools

Simon Coulton¹, M. Fasihul Alam², Sadie Boniface³, Paolo Deluca³, Kim Donoghue³, Eilish Gilvarry⁴, Eileen Kaner⁵, Ellen Lynch⁵, Ian Maconochie⁶, Paul McArdle⁴, Ruth McGovern⁵, Dorothy Newbury-Birch⁷, Robert Patton⁸, Ceri J. Phillips⁹, Thomas Phillips^{3,10}, Hannah Rose³, Ian Russell¹¹, John Strang³, Colin Drummond³

¹Centre for Health Services Studies, School of Social Science and Social Policy Research, George Allen Wing, University of Kent, Canterbury, Kent CT2 7NZ, UK

²College of Health Sciences, Qatar University, Doha, Qatar

³Addictions Department, National Addiction Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

⁴Northumberland Tyne and Wear NHS Foundation Trust, Newcastle, UK

⁵Institute of Health and Society, Newcastle University, Newcastle, UK

⁶Paediatric Emergency Medicine, Imperial College London, London, UK

⁷School of Health and Social Care, Teesside University, Middlesbrough, UK

⁸School of Psychology, University of Surrey, Guildford, UK

⁹Swansea Centre for Health Economics, College of Human and Health Sciences, Swansea University, Swansea, UK

¹⁰University of Hull, Hull, UK

¹¹Swansea University Medical School, Swansea, UK

Address correspondence to Simon Coulton, E-mail: s.coulton@kent.ac.uk

ABSTRACT

Objective To estimate and compare the optimal cut-off score of Alcohol Use Disorders Identification Test (AUDIT) and AUDIT-C in identifying at-risk alcohol consumption, heavy episodic alcohol use, ICD-10 alcohol abuse and alcohol dependence in adolescents attending ED in England.

Design Opportunistic cross-sectional survey.

Setting 10 emergency departments across England.

Participants Adolescents ($n = 5377$) aged between their 10th and 18th birthday who attended emergency departments between December 2012 and May 2013.

Measures Scores on the AUDIT and AUDIT-C. At-risk alcohol consumption and monthly episodic alcohol consumption in the past 3 months were derived using the time-line follow back method. Alcohol abuse and alcohol dependence was assessed in accordance with ICD-10 criteria using the MINI-KID.

Findings AUDIT-C with a score of 3 was more effective for at-risk alcohol use (AUC 0.81; sensitivity 87%, specificity 97%), heavy episodic use (0.84; 76%, 98%) and alcohol abuse (0.98; 91%, 90%). AUDIT with a score of 7 was more effective in identifying alcohol dependence (0.92; 96%, 94%).

Conclusions The 3-item AUDIT-C is more effective than AUDIT in screening adolescents for at-risk alcohol use, heavy episodic alcohol use and alcohol abuse. AUDIT is more effective than AUDIT-C for the identification of alcohol dependence.

Keywords adolescent, alcohol, diagnosis, screening

Simon Coulton, Professor of Health Service Research

M. Fasihul Alam, Assistant Professor in Health Economics

Sadie Boniface, Trial Manager

Paolo Deluca, Reader in Addictions Research

Kim Donoghue, Trial Manager

Eilish Gilvarry, Hon Professor of Addictions and Consultant Psychiatrist

Eileen Kaner, Professor of Public Health and Primary Care Research

Ellen Lynch, Research Associate

Ian Maconochie, Honorary Senior Lecturer and Consultant in Paediatric Emergency Medicine

Paul McArdle, Consultant Child and Adolescent Psychiatrist

Ruth McGovern, Senior Research Associate

Dorothy Newbury-Birch, Professor of Alcohol & Public Health Research

Robert Patton, Lecturer in Clinical Psychology

Ceri J. Phillips, Professor of Health Economics

Thomas Phillips, Professor of Nursing (Addictions)

Hannah Rose, Research Assistant

Ian Russell, Professor Emeritus (Medicine)

John Strang, Professor of Addiction Psychiatry

Colin Drummond, Professor of Addiction Psychiatry

Introduction

The excessive consumption of alcohol is a major global public health issue^{1,2} and places a significant burden on international health systems. While the majority of this burden lies with adult populations, for many the roots of problematic alcohol use lies in adolescence.³ Adolescence is a critical developmental stage when young people make behavioural and lifestyle choices that have the potential to impact on their health and wellbeing into adulthood. Inappropriate risk-taking is significantly associated with health and social harm during adolescence.⁴ Young people are much more vulnerable than adults to the adverse effects of alcohol use due to a range of physical and psychological factors that often interact. Adolescence is also a unique period whereby neural proliferation and subsequent 'pruning' processes may leave brain structures particularly vulnerable to the effects of alcohol.^{5,6}

A recent survey of alcohol consumed by 14–15 years old across 36 European countries reported that in the United Kingdom (UK) 87% had consumed alcohol at least once in their lifetime and 57% had consumed alcohol at least once in the past month.⁷ The prevalence of consuming alcohol increases with age, with data from 2016 indicating that 9% of boys aged 11–15 years, and 11% of girls had consumed alcohol in the past 7 days. Of these, 1% of 11 years old consumed alcohol in the past 7 days, increasing to 24% at age 15. In terms of quantity of alcohol consumed in the past 7 days mean consumption was 10.3 units for boys and 8.9 units for girls aged 11–15 years.⁸

An evidence based review of the risks and harms of alcohol consumption in young people⁹ provided a basis for the Chief Medical Officer for England recommendations for alcohol consumption in young people—that young people up to the age of 15 abstain completely from drinking and those aged 15–17 are advised not to drink, but if they do drink, they should not exceed 2–3 standard drinks in any day and no more than once per week.¹⁰

While there is a body of evidence addressing the effects of school based interventions for delaying the onset of drinking in adolescents,¹¹ and some evidence for interventions to delay the age of onset or reduce alcohol consumption for adolescents in other settings,^{12,13} there exists a paucity of evidence of the effectiveness of interventions to reduce adolescent alcohol use in primary care settings. Recommendations from the World Health Organization, US Surgeon General and American Academy of Paediatrics advocate that more evidence is needed on the effectiveness of opportunistic screening and interventions for adolescents who consume alcohol^{14,15} and this population has been identified as a key target group for the reduction of alcohol

use and related harm^{16,17} in both English and Scottish alcohol strategies.

The identification of adolescents who consume alcohol at problematic levels is a key element in any screening and intervention strategy. To offer such interventions practitioners need access to screening tools that are high in both sensitivity and specificity and are quick and easy to apply at minimal cost. Biochemical markers of alcohol use such as γ -glutamyl-transferase, aspartate aminotransferase, erythrocyte mean cell volume and percent carbohydrate deficit transferrin are impractical and of little use in this population and have been found to be inferior to short paper instruments in adult populations.¹⁸ The Alcohol Use Disorders Identification Test (AUDIT)¹⁹ is a 10-item self-completion instrument with established diagnostic properties for problematic alcohol use in adults that addresses three domains of alcohol-related problems; consumption, negative consequences and symptoms of dependence. AUDIT is one of the few screening instruments that specifically incorporates consumption into the scoring algorithm and may be particularly suitable for adolescents who are more likely to experience a range of alcohol-related problems as a result of consumption rather than psychological consequences of alcohol use. Further, it may be the case that the three specific alcohol consumption questions, AUDIT-C, may be equally efficient as a brief screening instrument as the full AUDIT. Previous studies suggest that the AUDIT may be more useful than other brief screening instruments in adolescent populations, but there is less consensus regarding appropriate cut-off points for different severities of alcohol use^{20–25} and no previous research has compared the relative effectiveness of AUDIT versus AUDIT-C as opportunistic screening approaches for adolescent populations. Much of the prior research has aimed to compare the performance of a variety of different screening instruments^{21,26–28} against more severe clinical alcohol use disorder criteria whereas adolescents are more likely to experience alcohol-related difficulties at lower levels of consumption and this is in part due to the pattern of consumption in the form of heavy episodic alcohol use.²⁹ In addition, the majority of studies have been conducted in older adolescent populations^{20,22} and often involve college students, primary care or hospitalized participants, rather than an opportunistic sample and are limited in their generalizability to the wider adolescent population and particularly limited in their generalizability to the UK.

Our aim was to estimate and compare the sensitivity, specificity, and diagnostic odd ratio of the AUDIT and AUDIT-C in identifying at-risk alcohol use, monthly heavy episodic alcohol use, alcohol abuse and alcohol dependence in the context of an opportunistic screening programme for adolescents, aged between 10 and 17 years, attending emergency

departments (ED) in England. To be acceptable as a screening test in clinical practice we expected the sensitivity and specificity at a selected cut-point would exceed 0.70.

Methods

The study was conducted in accordance with ethical approval from the National Health Service Multi-Centre Research Ethics Committee (ref: 12/L0/0799) and was registered in an appropriate trial registry (ref: ISRCTN 45300218).

Design

An opportunistic cross-sectional survey conducted between December 2012 and May 2013 across 10 ED's in England, encompassing a mix of metropolitan urban and rural centres across the North East, Yorkshire and Humber, London and the South. Consecutive attendees, between the hours of 8 am and midnight were approached by trained researchers after the initial triage assessment.

Researcher assessment was conducted blind to the results of the screening measure and the order of presentation of all measures was randomized using random permuted blocks of random length and embodied within the electronic data collection tool, stratified by age and centre. All assessment instruments used a 3-month assessment time-frame.

Measures

Gold standard measures

To elicit the gold-standard measures of at-risk drinking and monthly heavy episodic alcohol use we used the Time-Line Follow Back –90 days (TLFB90). This is a reliable and valid method to ascertain the frequency and quantity of alcohol consumed in clinical and non-clinical populations for periods ranging from 1 to 365 days.³⁰ The method has established psychometric properties for adolescent populations³¹ and is conducted by a trained researcher and the 90-day version takes ~30 min to complete. The responses to the interview are converted to UK standard drinks and can be used as either continuous or categorical outcomes. At-risk drinking was defined as consuming three or more standard drinks, where a standard drink equates to 8 g of pure ethanol, in a single day in the past 90 days. Monthly heavy episodic alcohol consumption was defined as consuming six or more standard drinks in a single drinking episode in each month over the past 3 months.

MINI-KID has established validity and reliability in the identification of psychiatric diagnoses for children and adolescents.³² The alcohol use module consists of seven detailed

questions that diagnose both alcohol abuse and alcohol dependence in accordance with ICD-10 criteria.

Screening tools

The AUDIT¹⁹ is a 10-item self-completion questionnaire that measures the quantity and frequency of alcohol consumption, drinking behaviour, alcohol-related problems and the symptoms of alcohol dependence. Each item is scored 0–4 and summed to create an overall score with a maximum of 40. The instrument is widely used in adult populations and a cut-off score of 8 or more has high levels of sensitivity (92%) and specificity (94%) for at-risk drinking in adult populations.¹⁹ The AUDIT-C³³ consists of the three consumption items of AUDIT and has been validated as a short-screen in adults, AUDIT-C scores range from 0 to 12, with five or more being indicative of at-risk alcohol use.

Participant recruitment

To be included in the survey, participants had to be aged between their 10th and 18th birthday, alert and orientated and able to communicate in English sufficiently to complete the survey. Participants were excluded if they had a severe injury requiring immediate intervention, were grossly intoxicated, had a serious mental health presentation or if they, or their parent or guardian, refused to provide consent.

Participants were provided with the study information sheet and allowed to ask any questions prior to providing consent. Where a child was aged 16 years or less Gillick competency was assessed³⁴ by a member of the clinical staff in the ED, and where a participant was not found competent consent was sought from the parent or carer. If a parent or carer was present with the child, parent consent was sought in addition to child consent. The survey was conducted in a private area of the ED with a trained researcher who was available to answer any questions and provide appropriate assistance. The survey was anonymous and self-completed using an electronic tablet device with the exception of the time-line follow back interview (TLFB)³⁰ that was conducted by the researcher. At the end of the survey participants were thanked for their time and returned to the care of the ED, were provided with an age-appropriate alcohol awareness leaflet and given a £5 gift voucher for participating.

Statistical methods

We compiled and analysed the results using STATA14. The influence of potential covariates of age and gender, and clustering by ED, were incorporated into the analysis using the ROCREG function. We constructed receiver operator characteristic curves on the basis of all continuous values of the

test results for AUDIT and AUDIT-C compared with each of the gold-standards; at-risk drinking, monthly heavy episodic alcohol use, alcohol abuse and alcohol dependence. We estimated the sensitivity and specificity of each cut-off point and generated the diagnostic odds ratio and associated 95% confidence interval. The diagnostic odds ratio was used to estimate optimal cut-points and is a measure of effectiveness of a dichotomous classification that is the ratio of the odds of being positive if truly positive relative to the odds of being positive if truly negative. It has advantages over other methods of diagnostic test effectiveness in that it is less susceptible to statistical artefacts, a criticism of the Youden Index, and does not rely on the sample prevalence, making it more useful for comparison across different study samples.³⁵

Results

Overall 5781 participants were asked to participate in the survey of whom 5377 (93%) consented to participate across the 10 ED's. The mean age was 13.3 (SD 2.1) years with

similar proportions of male (53.7%) and female (46.3%) participants and the majority White (72.6%). Overall 2112 (39.3%) had consumed alcohol at some time in the past and 1378 (25.6%) had consumed alcohol in the past 3 months. Those who had consumed alcohol tended to be older (14.8 versus 12.3 years) and were more likely to be white (83.4 versus 65.6%) (Table 1).

Using the sample to estimate the prevalence of drinking behaviours in adolescents attending ED, the prevalence of at-risk drinking was 14.8% (95% CI: 13.9–15.8%). The prevalence of monthly heavy episodic alcohol use was 10.6% (9.8–11.4%), alcohol abuse 2.4% (2.0–2.8%) and alcohol dependence 1.2% (0.9–1.5%). In the sample of those who had consumed alcohol in the past 3 months the prevalence of these behaviours was significantly higher (Table 2).

A significant positive correlation was identified for AUDIT score with the total number of standard drinks consumed in the past 3 months (Spearman rho, $r = 0.72$, 95% CI: 0.71–0.73; $P < 0.001$) and a similar correlation identified for AUDIT-C score ($r = 0.69$, 95% CI: 0.68–0.70; $P < 0.001$). Screening properties of the questionnaire were tested

Table 1 Demographic variables in 5377 adolescent attendees overall and by drinking status

Variable	All attendees (n = 5377)	Drinkers (n = 2112)	Non-drinkers (n = 3265)
Mean age (SD)	13.28 (2.07)	14.77 (1.64)	12.33 (1.74)
Age 10, n (%)	570 (10.6)	24 (1.1)	543 (16.8)
Age 11, n (%)	701 (13.0)	50 (2.4)	647 (20.0)
Age 12, n (%)	809 (15.0)	133 (6.3)	668 (20.6)
Age 13, n (%)	845 (15.7)	248 (11.7)	595 (18.4)
Age 14, n (%)	751 (14.0)	387 (18.3)	363 (11.2)
Age 15, n (%)	784 (14.6)	502 (23.8)	276 (8.5)
Age 16, n (%)	534 (9.9)	428 (20.3)	105 (3.2)
Age 17, n (%)	382 (7.1)	340 (16.1)	40 (1.2)
Male, n (%)	2886 (53.7)	1093 (51.8)	1793 (54.9)
Ethnicity, n (%)			
White	3726 (72.6)	1687 (83.4)	2039 (65.6)
Black	698 (13.6)	150 (7.4)	548 (17.6)
Chinese	4 (0.1)	1	3 (0.1)
Mixed	289 (5.6)	97 (4.8)	192 (6.2)
Asian	255 (5.0)	35 (1.7)	220 (7.1)
Other	144 (2.8)	45 (2.2)	99 (3.2)
Mode of arrival, n (%)			
Own means	3953 (74.0)	1667 (79.1)	2286 (70.6)
Ambulance	331 (6.2)	143 (6.8)	188 (5.8)
Police	2 (0.05)	2 (0.1)	0
Other	1059 (19.8)	295 (14.0)	764 (23.6)
Smoker, n (%)	481 (9.0)	455 (21.6)	26 (0.8)
Consumed alcohol in the past 3 months, n (%)	1378 (25.6)	1378 (64.9)	0

Table 2 Alcohol-related variables for all participants and those who consumed alcohol in the past 3 months

Variable	All participants (n = 5377)	Those who consumed alcohol in past 3 months (n = 1378)
Consumed alcohol in past 24 h, n (%)	115 (2.1)	115 (8.5)
Mean age in years (SD)	13.28 (2.07)	15.12 (1.51)
Mean age of first drink in years (SD)	12.74 (2.24)	12.90 (2.17)
Total alcohol consumed in past 3 months in standard units ^a (SD)	7.19 (39.47)	33.09 (79.28)
Hazardous alcohol consumption in past 3 months ^b , n (%)	796 (14.8)	796 (67.9)
Heavy episodic alcohol consumption in past 3 months ^c , n (%)	572 (10.6)	572 (48.8)
Alcohol abuse ^d , n (%)	127 (2.4)	127 (9.2)
Alcohol dependent ^d , n (%)	67 (1.2)	67 (5.0)
Mean AUDIT score (SD) (values can range from 0 to 40 with higher scores indicative of greater problems)	1.18 (1.78)	4.83 (5.03)
Mean AUDIT-C score (SD) (values can range from 0 to 12 with higher scores indicative of greater problems)	0.75 (3.23)	2.98 (2.46)

^aStandard unit equivalent to 8 g of ethanol.

^bHazardous consumption defined as drinking three or more standard units in a single day.

^cHeavy episodic consumption defined as drinking six or more standard units in a single drinking episode.

^dUsing ICD-10 criteria using MINI-KID.

against the gold standard criteria for at-risk drinking, heavy episodic alcohol consumption, alcohol abuse and alcohol dependence. Screening results for all cut-points were assessed and the results of those around the optimal cut-point are reported in Table 3.

The optimum cut-off point for AUDIT in identifying either at-risk drinking, monthly heavy episodic drinking or alcohol abuse was 4 or more, which provided the optimal cut-point to provide acceptable sensitivity, specificity and diagnostic odds. An AUDIT-C score of 3 or more demonstrated almost identical diagnostic properties but with a significantly better sensitivity for at-risk drinking.

An AUDIT score of 7 or more provided a significantly more effective cut-point for alcohol dependence than any other cut-point and demonstrated significantly better diagnostic properties than an AUDIT-C score of 5 or more.

We assessed the potential influence of age, gender and ED on our findings and found these effects to minimal and not statistically significant from our main findings. The results without incorporation of these variables is therefore reported.

Discussion

Main findings of this study

A simple short three item self-completed screening instrument, the AUDIT-C, is overall more effective than the longer 10-item AUDIT in identifying adolescents who engage in

at-risk of alcohol consumption, monthly heavy episodic alcohol use and fulfil ICD-10 criteria for alcohol abuse. Further the AUDIT with a cut-off score of 7 is more efficient than AUDIT-C in identifying adolescents with alcohol dependence. In addition, AUDIT-C and AUDIT are widely employed as screening tools for adults in clinical and non-clinical settings and these can be applied equally to adolescent populations with these lower cut-off scores. We conclude that AUDIT-C should be employed with this population with a cut-off score of 3 as a positive screen for at-risk drinking, monthly heavy episodic alcohol use and alcohol abuse. For those who score 5 or more on AUDIT-C we recommend the use of the additional 7 questions constituting the full AUDIT be administered. With those scoring 7 or more being clinically assessed for alcohol dependence.

What is already known on this topic

There is a body of evidence suggesting that interventions for alcohol using adolescents are effective and that they are more effective when targeted as secondary prevention strategies, i.e. at those already engaged in consuming alcohol.^{12,13}

A critical first step in the delivery of interventions is employing opportunistic screening tools and the combination of effective screening tools and intervention strategies offers significant potential to reduce the burden of alcohol use on adolescents, health systems and wider society and further consideration should be given to the routine opportunistic

Table 3 Area under the receiver operator curve (AUC), sensitivity, specificity and diagnostic odd ratio of AUDIT and AUDIT-C cut-points for hazardous drinking, monthly episodic alcohol use, alcohol abuse and alcohol dependence for 5377 adolescent attendees at ED

<i>Outcome</i>	<i>Prevalence % (95% CI)</i>	<i>AUC (95% CI)</i>	<i>Sensitivity % (95% CI)</i>	<i>Specificity % (95% CI)</i>	<i>Diagnostic odd ratio (95% CI)</i>
At-risk/hazardous drinking	15 (14; 16)				
AUDIT					
≥3		0.81 (0.79; 0.94)	78 (75; 82)	94 (94; 95)	55 (47; 87)
≥4		0.81 (0.79; 0.94)	75 (72; 78)	98 (98; 99)	147 (126; 351)
≥5		0.84 (0.82; 0.87)	65 (61; 69)	98 (98; 99)	91 (77; 220)
AUDIT-C					
≥2		0.84 (0.82; 0.87)	91 (88; 93)	89 (87; 91)	81 (49; 134)
≥3		0.98 (0.97; 0.99)	89 (86; 91)	97 (96; 97)	261 (147; 242)
≥4		0.98 (0.97; 0.98)	72 (68; 77)	97 (96; 97)	83 (51; 108)
Monthly episodic use	10 (10; 11)				
AUDIT					
≥3		0.92 (0.90; 0.95)	80 (77; 82)	92 (89; 95)	46 (27; 86)
≥4		0.87 (0.84; 0.91)	78 (74; 81)	97 (97; 98)	114 (92; 109)
≥5			58 (54; 63)	98 (94; 99)	67 (18; 168)
AUDIT-C					
≥2			82 (79; 85)	89 (87; 90)	37 (25; 51)
≥3			76 (73; 80)	98 (97; 98)	155 (87; 196)
≥4			61 (57; 66)	99 (96; 99)	77 (32; 192)
Alcohol abuse	2 (2; 3)				
AUDIT					
≥3			94 (88; 97)	85 (82; 88)	88 (33; 237)
≥4			93 (87; 96)	88 (87; 89)	97 (44; 194)
≥5			83 (75; 88)	92 (91; 93)	56 (30; 97)
AUDIT-C					
≥2			91 (85; 95)	85 (84; 86)	57 (30; 116)
≥3			91 (85; 95)	90 (88; 91)	91 (42; 192)
≥4			65 (56; 73)	93 (92; 93)	25 (15; 36)
Alcohol dependent	1 (1;2)				
AUDIT					
≥6			96 (89; 99)	92 (90; 94)	276 (73; 1551)
≥7			96 (89; 99)	94 (95; 95)	376 (154; 1881)
≥8			91 (81; 96)	95 (95; 96)	192 (81; 576)
AUDIT-C					
≥4			85 (79; 88)	92 (91; 93)	65 (43; 97)
≥5			80 (67; 89)	95 (95; 95)	76 (39; 154)
≥6			67 (55; 77)	97 (96; 97)	65 (39; 108)

implementation of screening strategies for adolescent populations.

What this study adds

Routine alcohol screening of adolescents should be considered across the UK National Health Service. This study

demonstrates that the process can be simplified by using short screening tools already in use for adult populations. This requires appropriate training, resources and incentives for staff. Identifying those adolescents that may benefit from interventions to address alcohol use and associated multiple risk behaviours will help to reduce the burden of alcohol use

across the health service and society. This has the potential to enhance the future health of the adolescent population well into adulthood.

Limitations of this study

Our study was conducted in ED and this could be seen as compromising the generalizability of the findings to other health settings. Yet adolescents are far less frequent attenders at primary care and the ED provides an opportunity to access this population and in turn provides the ‘teachable moment’, that is hypothesized to play a crucial role in effective behaviour change.³⁶ Further, we aimed to ensure generalizability of our sample to other ED’s in the UK by including centres covering rural and urban areas and areas with the lowest and highest population prevalence of adolescent alcohol use and areas of high and low socio-economic status. In addition, our estimates of alcohol use problems compare well with national epidemiological surveys, that suggest 27% of adolescents consume alcohol versus 26% in our study, 9% have been drunk three or more times in the past 4 weeks compared with 11% of episodic drinkers in the past 3 months in our study.³⁷ We also recognize that those who scored negative on the screening tool and outcome assessments may have misreported their alcohol consumption and we took a variety of steps to ameliorate this by ensuring anonymity and confidentiality. Previous evidence would suggest this form of social desirability bias is limited.³⁸ This study was the first study of the screening instruments in a real-life health setting in the UK, one where the burden of alcohol use is a real concern.

Conflicts of interest

The authors have no conflict of interest to declare.

Acknowledgement

The views expressed are those of the authors and not necessarily those of the NHS, NIHR or Department of Health. We thank all young people who helped us design the survey and guided the research team on how to best present the survey to young people to maximize engagement. We also thank all those young people who took the time to complete the survey and all the staff in participating ED’s.

Funding

This work was funded by the NIHR Programme Grants for Applied Research (Rp-PG-0609-10162). Colin Drummond is partly funded by the NIHR Biomedical Research Centre for

Mental Health at South London and Maudsley NHS Foundation Trust and King’s College London and partly funded by the NIHR Collaborations for Leadership in Applied Health Research and Care South London at King’s College Hospital NHS Foundation Trust.

Authors contributions

SC, PD, KD, e.g. EK, IM, PM, RM, DNB, RP, TP, IR, JS and CD contributed to the design of the programme of research. FA, SB, EL, CP, HR contribute to the ongoing data collection analysis and interpretation of the research. SC conducted the analysis reported in the article and wrote the initial draft. All authors have read and commented on subsequent drafts of the article.

References

- 1 Rehm J, Room R, Graham K *et al.* The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction* 2003;**98**:1209–28.
- 2 Rehm J, Room R, Monteiro M *et al.* Alcohol as a risk factor for global burden of disease. *Eur Addict Res* 2003;**9**:157–64.
- 3 Bellis M, Phillips-Howard P, Hughes K *et al.* Teenage drinking, alcohol availability and pricing: a cross-sectional study of risk and protective factors for alcohol-related harms in school children. *BMC Public Health* 2009;**9**:380.
- 4 Odgers CL, Caspi A, Nagin DS *et al.* Is it important to prevent early exposure to drugs and alcohol among adolescents? *Psychol Sci* 2008;**19**:1037–44.
- 5 Guerri C, Pascual M. Mechanisms involved in the neurotoxic, cognitive, and neurobehavioral effects of alcohol consumption during adolescence. *Alcohol* 2010;**44**:15–26.
- 6 Squeglia LM, Jacobus J, Tapert SF. The influence of substance use on adolescent brain development. *Clin EEG Neurosci* 2009;**40**:31–8.
- 7 Danielsson AK, Wennberg P, Hibell B *et al.* Alcohol use, heavy episodic drinking and subsequent problems among adolescents in 23 European countries: does the prevention paradox apply? *Addiction* 2012;**107**:71–80.
- 8 NHS Digital. *Smoking, Drinking and Drug Use Among Young People*. London: HMSO, 2016.
- 9 Newbury-Birch D, Gilvarry E, McArdle P *et al.* The impact of alcohol consumption on young people: a review of reviews. Department of Children Schools and Families, 2009.
- 10 Donaldson L. *Guidance on the Consumption of Alcohol by Children and Young People*. London: Department of Health, 2009.
- 11 Conrod PJ, Castellanos N, Mackie C. Personality-targeted interventions delay the growth of adolescent drinking and binge drinking. *J Child Psychol Psychiatry* 2008;**49**:181–90.
- 12 Tanner-Smith EE, Lipsey MW. Brief alcohol interventions for adolescents and young adults: a systematic review and meta-analysis. *J Subst Abuse Treat* 2015;**51**:1–18.

- 13 Patton R, Deluca P, Kaner E *et al.* Alcohol screening and brief intervention for adolescents: the how, what and where of reducing alcohol consumption and related harm among young people. *Alcohol Alcohol* 2014;**49**:207–12.
- 14 American Academy of Pediatrics Committee on Substance Abuse. *Alcohol Use and Abuse*. Washington: Pediatrics AAo, 2001.
- 15 World Health Organisation. *Orientation Programme on Adolescent Health for Healthcare Providers*. Geneva: Organization WH, 2006.
- 16 UK Home Office. *The Government's Alcohol Strategy*. London: Office H, 2012.
- 17 Scottish Government. Changing Scotland's relationship with alcohol: a discussion paper on our strategic approach. Edinburgh, June 2008.
- 18 Coulton S, Drummond C, James D *et al.* Opportunistic screening for alcohol use disorders in primary care: comparative study. *Br Med J* 2006;**332**:511–7.
- 19 Saunders JB, Aasland OG, Babor TF *et al.* Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on early detection of persons with harmful alcohol consumption. *Addiction* 1993;**88**:791–804.
- 20 Aertgeerts B, Buntinx F, Bande-Knops J *et al.* The value of CAGE, CUGE, and AUDIT in screening for alcohol abuse and dependence among college freshmen. *Alcohol Clin Exp Res* 2000;**24**:53–7.
- 21 Chung T, Colby S, Barnett N *et al.* Screening adolescents for problem drinking: performance of brief screens against DSM-IV alcohol diagnoses. *Centre Alcohol Addict Stud* 2000;**61**:579–87.
- 22 Demartini KS, Carey KB. Optimizing the use of the AUDIT for alcohol screening in college students. *Psychol Assess* 2012;**24**:954–63.
- 23 Kelly T, Donovan J, Kinnane J *et al.* A comparison of alcohol screening instruments among under-aged drinkers treated in emergency departments. *Alcohol Alcohol* 2002;**37**:444–50.
- 24 Knight J, Sherritt L, Harris S *et al.* Validity of brief alcohol screening tests among adolescents: a comparison of the AUDIT, POSIT, CAGE and CRAFFT. *Alcoholism* 2003;**27**:67–73.
- 25 Santis R, Garmendia ML, Acuna G *et al.* The Alcohol Use Disorders Identification Test (AUDIT) as a screening instrument for adolescents. *Drug Alcohol Depend* 2009;**103**:155–8.
- 26 Knight JR, Sherritt L, Harris SK *et al.* Validity of brief alcohol screening tests among adolescents: a comparison of the AUDIT, POSIT, CAGE, and CRAFFT. *Alcohol Clin Exp Res* 2003;**27**:67–73.
- 27 Kelly TM, Donovan JE, Kinnane JM *et al.* A comparison of alcohol screening instruments among under-aged drinkers treated in emergency departments. *Alcohol Alcohol* 2002;**37**:444–50.
- 28 Kelly TM, Donovan J, Chung T *et al.* Alcohol use disorders among emergency department-treated older adolescents: a new brief screen (RUFT-Cut) using the AUDIT, CAGE, CRAFFT, and RAPS-QF. *Alcoholism* 2004;**28**:746–53.
- 29 Donoghue K, Rose H, Boniface S *et al.* Alcohol consumption, early-onset drinking, and health-related consequences in adolescents presenting at emergency departments in England. *J Adolesc Health* 2017;**60**(4):438–46.
- 30 Sobell LC, Sobell B. Time-Line Follow Back: a technique for assessing self-reported alcohol consumption. In: Litten RZ, Allen P (eds). *Measuring Alcohol Consumption: Psychosocial and Biological Methods*. Totowa, NJ: Human Press, 1992,41–72.
- 31 Donohue B, Azrin N, Strada M *et al.* Psychometric evaluation of self- and collateral timeline follow-back reports of drug and alcohol use in a sample of drug-abusing and conduct-disordered adolescents and their parents. *Psychol Addict Behav* 2004;**18**:184–9.
- 32 Sheehan DV, Sheehan KH, Shytle RD *et al.* Reliability and validity of the Mini International Neuropsychiatric Interview for Children and Adolescents (MINI-KID). *J Clin Psychiatry* 2010;**71**:313–26.
- 33 Bradley K, Debenedetti A, Volk R *et al.* AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcoholism* 2007;**31**:1208–17 (10).
- 34 Hunter D, Pierscionek BK. Children, Gillick competency and consent for involvement in research. *J Med Ethics* 2007;**33**:659–62.
- 35 Glas AS, Lijmer JG, Prins MH *et al.* The diagnostic odds ratio: a single indicator of test performance. *J Clin Epidemiol* 2003;**56**:1129–35.
- 36 Touquet R, Brown A. Alcohol misuse: positive response. Alcohol Health Work for every acute hospital saves money and reduces repeat attendances. *Emerg Med Aust* 2006;**18**:103–7.
- 37 Fuller E. *Smoking, Drinking and Drug Use Among Young People in England*. London: NHS Information Centre 2012, 2011.
- 38 Kypri K, Wilson A, Attia J *et al.* Social desirability bias in the reporting of alcohol consumption: a randomized trial. *J Stud Alcohol Drugs* 2016;**77**:526–31.