A pyrene-based approach to tune emission color from blue to yellow

Chuan-Zeng Wang,^a Hisashi Ichiyanagi,^a Koya Sakaguchi,^a Xing Feng,^{*b} Mark R.J. Elsegood,^c Carl Redshaw,^d and Takehiko Yamato^{*a}

a Department of Applied Chemistry, Faculty of Science and Engineering, Saga University, Honjo-machi 1, Saga 840-8502 Japan, E-mail: yamatot@cc.saga-u.ac.jp.

b School of Printing and Packaging Engineering, Beijing Institute of Graphic Communication, 1 Xinghua Avenue (Band Two), Daxing, Beijing, 102600, P. R. China, E-mail: hyxhn@sina.com.

c Chemistry Department, Loughborough University, Loughborough, LE11 3TU, UK.

d School of Mathematics and Physical Sciences, The University of Hull, Cottingham Road, Hull, Yorkshire HU6 7RX, UK.



ABSTRACT: The development of functionalized, luminescent, pyrene-based monomers has been and continues to be an area of great interest in terms of the design and fabrication of optical and electronic devices. Herein, a facile strategy to tune the emission color of pyrene-based chromophores has been established by simple functional group modification at the para position to the diphenylamino on the donor building block. Intriguing photophysical properties were obtained and are described both in different solutions and in the solid state. The results obtained could be explained by the Hammett method and by density functional theory (DFT) calculations. A good correlation was observed between the Hammett σ_{para} constants of the functional groups para to the phenyl and the wavenumber (cm⁻¹) of the emission profile. This positive correlation, namely between the σ constants of the functional groups and the emission properties of the monomers, can be used to develop a predictive method for these types of systems.

INTRODUCTION

Chromophores are a class of functional π -electron systems that have progressively established themselves as useful tuneable molecules in modern organic electronics and organic electronic devices.¹ Indeed, it is important to have the ability to tune the emission property of materials in a facile approach when designing light-emitting diodes,² bio-imaging probes,³ and other photoe-lectric emitting devices,⁴ particularly in the visible region. Various strategies have been established to achieve efficient full-color emitting materials which are based on underlying mechanisms, including intramolecular charge-transfer (ICT),⁵ twisted intramolecular charge-transfer (TICT),⁶ excimer,⁷ excited-state proton transfer (ESIPT),⁸ and photo-induced electron transfer (PET).⁹ Such strategies provide significant guidance when designing novel luminescent molecules. In particular, it is worth noting that in recent years the introduction of donors and acceptors has played a crucial role in organic electronics based on ICT.^{5, 10} Furthermore, much effort has been devoted to developing highly efficient host materials.

Among the many promising unmodified host candidates, polyaromatic pyrene have attracted attention because of their high thermal and photochemical stability, pure blue fluorescence, planar geometry, and natural high charge carrier mobility.¹¹ It is also possible to append electronaccepting/donor groups at the suitable positions on the pyrene core, which can result in enhanced intramolecular charge-transfer. Generally speaking, acceptors and donors provide energetically high lying occupied molecular orbitals (HOMOs) and low lying unoccupied molecular orbitals (LUMOs), respectively. The D–A unit is conducive to the fine-tuning of the electronic interactions and charge transfer efficiencies.¹² A wide variety of D–A type pyrene derivatives have been investigated by comparing experimental and theoretical results in recent years.¹³ For instance, many pyrene-based chromophores, based on both covalent and acceptor-donor structures, have been found to possess differing yet distinctive emission properties by functionalization at the different positions of the pyrene core.^{4a, 14} On the basis of these experimental observations, in case of a donor/acceptor substituted molecule, the band-gap of the new material can be significantly reduced when compared to the pure components.¹⁵

Herein, we present a facile strategy to investigate the influence of covalent donor groups. Specifically, a series of classical donor groups, which differ only by the functional group *para* to an N,N-diphenylamine core, have been studied. The groups were introduced at the 1-, and 3positions of 7-*tert*-butylpyrene *via* a Buchwald–Hartwig amination reaction. As anticipated, we observed distinct/different emission properties both in solution and in the solid state, simply by fine-tuning of the *para* substituents of the *N*,*N*-diphenylamine core. Broad and tuneable emission, from blue to yellow wavelengths, were achieved in dilute CH₂Cl₂ solutions. Moreover, one of the compounds was found to emit orange light in the solid state. To decipher the underlying mechanisms responsible for these attractive properties, we further investigated the solvato-chromism of the five compounds and report their absorption and emission spectra. The pronounced positive solvatochromism combined with theoretical calculations indicate that these systems have potential for the design of full-color organic electrochromic devices.¹



Scheme 1. Synthesis of pyrene-based light-emitting monomers 3.

RESULTS AND DISCUSSION

Shown in Scheme 1 is the synthesis of **3a-e** starting from 7-tert-butylpyrene. This synthetic strategy was adapted from the well-known Buchwald-Hartwig amination reaction. The compounds were characterized by ¹H and ¹³C NMR spectroscopy, high resolution mass spectrometry. The detailed synthetic procedures and characterization data are given in the Experimental Section and the Supporting Information (Figure S1-10). All compounds exhibited good solubility in common organic solvents with excellent thermal stability. The thermal properties of these monomers were determined by differential scanning calorimetry (DSC) and thermogravimetric analysis (TGA) under a nitrogen atmosphere and the results are shown in Table 1. Compounds of type 3 showed very high thermal stability with decomposition temperatures (T_d) of 348 to 392 °C and melting temperatures (T_m) of 198 to 262 °C. The high thermal values endow high morphologic stability of the amorphous phase in the solid state. The molecular structure of 7-tert-butyl-1,3bis[N,N-bis(4'-methoxyphenyl)amino]pyrene (3e) was unambiguously confirmed by X-ray crystallography. Shown in Figure 1 is the crystal structure of **3e**, which shows the dihedral angles between the C₆ aromatic rings and the pyrene unit. Unsurprisingly, the *p*-OMe-phenyl groups point in different directions (Figure 1b). The dihedral angles between the pyrene core (C1 > C16) and the C₆ aromatic rings are as follows: $75.81(3)^{\circ}$ (C21 > C26), $79.34(3)^{\circ}$ (C28 > C33),

60.79(11)° (C35 > C40) for the major disorder component, and 81.25(3)° (C42 > C47), respectively. The aromatic rings of the diphenylamino unit exhibit a twisted geometry, which, when combined with the bulky *tert*-butyl groups, reduces the π - π stacking interactions. By comparison, both the phenyl and pyrene units of **3e** are involved in two kinds of π interactions between adjacent molecules, which results in the formation of a three-dimensional framework (Figure 2a).¹⁷



Figure 1. The crystal structure of compound 3e: (a) top view and (b) side view.



Figure 2. (a) The molecular packing of compound 3e as viewed parallel to the *b* axis of the unit cell; (b) the principal intermolecular packing interactions (Most hydrogen atoms are omitted for clarity).

A detailed investigation indicated that two types of π interactions were present, namely 1) a MeC-H… π interaction of the methoxy group with the phenyl unit at a distance of 2.86 Å (yellow dashed lines); and 2) an edge-to-face ArC-H… π interaction between the edge of the phenyl unit and the face of the pyrene core at a distance in the range 3.56–4.03 Å (green dashed lines). This interestingly results in a regular arrangement of alternating, anti-parallel stacks (Figure 2b). To be more exact, a compact supramolecular assembly is observed from alternating donor of phenyl groups (orange color) and the pyrene core (blue color) due to weak intermolecular interactions.

R	$\frac{\lambda_{abs}\left(nm\right)}{sol^{a}}$	λ _{PL} (nm) sol ^a / <mark>solid</mark> ^b	$\begin{array}{c} \log \epsilon \\ M^{-1} \cdot cm^{-1} \end{array}$	$T_{\rm m}$ (°C) ^c	$T_{\rm d}$ (°C) ^d	$arPhi_{ ext{PL}}(\%) \ ext{sol}^a/ rac{ ext{solid}^b}{ ext{solid}^b}$	HOMO (eV) ^e	LUMO (eV) ^f	$E_{\rm g}({\rm eV})^{\rm g}$
3 a	294, 414	465/525	4.59	198	348	79/33	-4.95	-1.95	3.00
3b	299, 414	467/514	4.62	238	372	82/46	-4.79	-1.79	3.00
3c	302, 424	489/494	4.65	214	383	78/29	-4.65	-1.72	2.93
3d	302, 425	491/472	4.61	229	391	57/10	-4.68	-1.76	2.92
3e	301, 434	525/573	4.62	262	392	60/24	-4.43	-1.57	2.86
^a Measured in dichloromethane at room temperature. ^b Measure in the solid state as powder. ^c Melting point: Obtained from DSC									
measurements. ^d Thermal decomposition temperature: Obtained from TGA measurements. ^e DFT/B3LYP/6-31G* using Gaussian.									
f LUMO = Eg + HOMO, ^g Estimated from the absorption edge of UV-Vis spectra.									

Table 1. The physical properties of compounds of type 3a-e.

These molecular designs were guided by density functional theory (DFT) calculations at the B3LYP/6-31g-(d)) level to examine their lowest unoccupied molecular orbital (LUMO), and highest occupied molecular orbital (HOMO). As depicted in Figure 3, the HOMOs of **3a-e** are mainly spread over the diphenylamino moiety and the pyrene core, while the LUMOs are mostly distributed on the pyrene core and slightly extended to the phenyl ring. It is noteworthy that the separated HOMOs and LUMOs are important for high luminescence efficiency of the molecule in certain circumstances, which could be increases the potential of Thermally Activated Delayed

Fluorescence emitters by adjusting the energy gap.¹⁸



Figure 3. Frontier-molecular-orbital distributions and energy levels diagram of **3a-e** and **1** by DFT calculations.

More and more classical donors and acceptors are attracting interest in the field of organic electronics. Herein, donor functionalized diphenylamines were selected as the hole-transport materials.¹⁹ The contours of the LUMO and HOMO of compounds **3a-e** revealed that the incorporation of a strong electron-donating group on the diphenylamine decreases the contribution of the pyrene core in the HOMOs. The distribution of LUMO orbitals of **3a-e** indicated the acceptor nature of the pyrene unit. It is very likely that intramolecular charge-transfer (ICT) states would be formed for these monomers. In this case of the donor appended molecules, the energy gaps were further evaluated. As shown in Table 1 and the energy level diagram Figure 3, the energetic proximity of the low lying LUMO and high lying HOMO results in a reduced energy gap compared with compound **1**.²⁰ The most interesting finding from the DFT calculations is that there is a positive correlation trend between the electronegativity of the functional groups occupying the *para* position at the phenyl moiety and the HOMO–LUMO gaps, which makes it possible to predict that the monomers **3a-e** would have red-shifted absorption spectra. Based on our preliminary theoretical guidance, the steady-state absorption and emission properties were investigated both in solution and in the solid state.



Figure 4. Absorption (top) and emission (bottom) profiles for compounds 3 in CH₂Cl₂ solution. The absorption and emission spectra of compounds 3a-e both in solution and in the solid state are shown in Figure 4, S11 and summarized in Table 1. As shown in Figure 4 (top), type 3 compounds have a strong, high-energy band centered at 294–302 nm, which is mainly due to the π - π * transitions of the substituents and pyrene core. In this type of molecule, with classical donor groups at the terminal aromatic rings of pyrene, all of the monomers exhibited a broad absorption band with low-energy absorption (414–434 nm), indicating that their excited states possess some charge transfer (CT) absorption associated with the ICT from the para substituents of the N,N-diphenylamine terminal to the pyrene unit, which is attributed to the separated HOMO and LUMO distributions. It was found that the low-energy band is relatively more affected and sensitive than the highenergy band by the nature of the functional groups *para* to the phenyl moiety according to the molar absorption coefficient (Figure 4 top). Compounds 3a-e have a high-energy band which corresponds to a molar absorption coefficient centered at 48046-50002 cm⁻¹ M⁻¹, and a lowenergy band which corresponds to a molar absorption coefficient ranging from 22135 to 31614 cm⁻¹ M⁻¹. The band gap energy (E_g) of compounds **3a-e** calculated from the absorption edge of the UV/Vis spectra are also summarized in Table 1. The LUMO levels of 3a-e were estimated from the E_g and HOMO levels.



Figure 5. Hammett σ_{para} constants *versus* wavenumber (cm⁻¹) for **3a** (black), **3b** (red), **3c** (blue), **3d** (green), and **3e** (orange).

The emission maxima of this set of monomers **3** are in the range 465–525 nm in dilute CH₂Cl₂ solutions with a systematic bathochromic shift in the order **3a** < **3b** < **3c** < **3d** < **3e** (Figure 4 bottom), consistent with the electronegativity of the substituents at the *para* position of the phenyl moiety. These results, combined with the theoretical calculations, indicate that the energy gap (E_g) could be tuned between the group and the excited states by introducing different substituent groups. More interestingly, when the $\lambda_{em max}$ are converted to wavenumbers (cm⁻¹) and plotted *versus* the Hammett σ_{para} constants for the functional groups *para* on the phenyl, a positive correlation was clearly observed obviously (Figure 5). This result establishes a promising strategy for the design of pyrene-based luminophores by choosing functional groups on the basis of their Hammett σ_{para} value.

The emission spectra in the solid state were also recorded, and all the compounds were found to exhibit a red shift trend compared with their corresponding emission spectra in solution (Table 1, and Figure S1). However, the emission band maxima of **3d** was blue-shifted (19 nm) relative to the results in solution, because of the introduction of the *para*-position of bulky electron donor *tert*-butyl group in phenyl ring that not only plays a role in restricting aggregation in the solid state but also affects the conformation of the electronic structures. It was observed that the quantum yields showed no trend in solution. Additionally, a decrease in the photoluminescence quantum yield of these compounds was observed in the solid state. In order to gain further insight into the pronounced difference of luminescence quantum efficiency in both these two states, additional analysis of compound **3e** using powder X-ray diffraction (PXRD) was investigated. And the PXRD pattern of **3e** shows an obvious crystalline phase (Figure S12).

that the powder is mainly crystalline, so it can be better understandable through the guidance of stacking of X-ray structure. The *H*- and *J*-aggregates²¹ were elucidated since it is generally agreed that these are the best known theoretical predictors (Figure 6). According to the optical results, we assumed that these molecules may aggregate in a parallel way (face-to-face stacking) to form a sandwich-type arrangement (*H*-dimer) with the aid of the ArC–H··· π interaction between the pyrene core combine the crystallization results. Interestingly, the donor moiety with bulky substituent can induces displacement of the monomers into a slipped stacking arrangement in the solid state, which reveals typically self-assemble into *H*-aggregates due to the slipping angles resulting from the equilibration of electrostatic and dispersion interactions clearly favors sandwich face-to-face contacts with markedly overlapping π surfaces rather than the more slipped arrangements required for the *J*-aggregates.²² Further evidence to be confirmed, the low quantum yield of compounds **3** in the solid state are also attributed to the *H*-aggregate of the dimer, as the lowest excited energy level is forbidden.



Figure 6. Schematic representation of the relationship between chromophore arrangement based on the molecular exciton theory.



Figure 7. (left) Emission spectra of **3e** in cyclohexane, 1,4-dioxane, THF, DCM, and DMF at room temperature; (right) Lippert-Mataga plots for compound **3e**.

The fluorescence spectra of **3a-e** are highly sensitive toward the solvent polarity, while the absorption spectra exhibit slightly shift (to see figure 13). The spectral properties of the solvatochromism were investigated in different solvents with varying polarity (cyclohexane, 1,4-dioxane, tetrahydrofuran, dichloromethane, and dimethyl formamide), the results are shown in the Supporting Information (Figure S 3). Compound **3e** is a representative example, for which Figure 7 (left) shows a significant red-shift on increasing the solvent polarity from cyclohexane to DMF (47 nm for **3a**, 43 nm for **3b**, 54 nm for **3c**, 54 nm for **3d**, 71 nm for **3e**, respectively). This phenomenon was further evaluated by the relationship between the Stokes shifts in various solvents and the Lippert equation,²³ which shows a linear correlation between these two factors (Figure 7 right). The trend in the slope of the Lippert-Mataga plots follow the order **3e** > **3d** > **3c** > **3b** \approx **3a**, and we can find that the value of the slope of the fitting line for **3e** (5021.5) is higher than that for **3b** (3502.4), indicating the intramolecular excited state with a larger dipolar moment for the former than the latter. And those results revealing extraordinary sensitivity to solvent polarity. Moreover, the distinct charge separation and higher dipole moment were further confirmed in the excited state.²⁴

CONCLUSIONS

In summary, a series of tuneable pyrene-based monomers **3a-e** were designed and synthesized by a facile strategy, and their optical properties were investigated both experimentally and computationally. It was determined that these systems exhibited predictable photophysical properties by introducing different substituents at the *para* position of the phenyl moieties. Moreover, a positive correlation between the wavelength of the $\lambda_{em max}$ and the Hammett σ_{para} constants for the functional groups provided evidence that these systems may be used to tune the emission color from blue to yellow or orange, and further investigations aimed at developing these optoelectronic materials for practical applications are ongoing in our group.

EXPERIMENTAL SECTION

All melting points (Yanagimoto MP-S1) are uncorrected. ¹H NMR spectra (300 MHz) were recorded on a Nippon Denshi JEOL FT-300 NMR spectrometer with SiMe₄ as an internal reference: *J*-values are given in Hz. IR spectra were measured for samples as KBr pellets in a Nippon Denshi JIR-AQ2OM spectrophotometer. UV-vis spectra were recorded on a Perkin Elmer Lambda 19 UV/VIS/NIR spectrometer. Mass spectra were obtained on a Nippon Denshi JMS-01SA-2 spectrometer at 75 eV using a direct-inlet system.

Materials

Unless otherwise stated, all other reagents used were purchased from commercial sources and were used without further purification. The preparations of 2-*tert*-butylpyrene (1) and 7-*tert*-butyl-1,3-dibromopyrene (2) were described previously.²⁵

Synthetic Procedures

Synthesis of 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-fluorophenyl)amino)pyrene (3a)

A mixture of 7-*tert*-butyl-1,3-dibromopyrene **2** (300 mg, 0.72 mmol), bis(4-fluorophenyl)amine (443 mg, 2.16 mmol), palladium(II) acetate (8.1 mg, 36 µmol), sodium *tert*-butoxide (416 mg, 4.33 mmol), and tri-*tert*-butylphosphine (1M in toluene, 0.07 mL, 70 µmol) in toluene (15 mL) was heated to reflux for 12 h under argon. The mixture was chilled to room temperature and quenched with water (100 mL). The mixture was extracted with dichloromethane (3 × 50 mL), and the combined extracts were washed with water and brine, dried with MgSO4 and concentrated. The residue was chromatographed over silica gel (Wako C-300, 200 g) with chloroform:hexane (2:8) as eluent to give an orange solid. Recrystallization from methanol afforded 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-fluorophenyl)amino)pyrene **3a** (368 mg, 77%) as orange prisms. M.p. 198–200°C; ¹H NMR (300 MHz, CDCl₃): $\delta_{\rm H} = 1.53$ (s, 9H, *t*Bu), 6.85–6.97 (m, 16H, Ar-*H*), 7.47 (s, 1H, pyrene-*H*), 7.88 (d, *J* = 9.2 Hz, 2H, pyrene-*H*), 8.00 (d, *J* = 9.2 Hz, 2H, pyrene-*H*), 8.13 ppm (s, 2H, pyrene-*H*); ¹³C NMR (100 MHz, CDCl₃): $\delta_{\rm C} =$ **31.8**, 35.20, 115.9, 116.1, 122.8, 123.2, 123.2, 126.0, 127.7, 127.8, 128.0, 131.0, 141.6, 144.7, 144.7, 150.0, 156.9, 159.3 ppm; **FAB-MS (MALDI-TOF)**: *m/z* calcd for C44H₃₂F4N₂ 664.2502 [M⁺]; found **664.2526** [M⁺].

Synthesis of 7-*tert*-butyl-1,3-bis(*N*,*N*-diphenylamino)pyrene (3b)

A mixture of 7-*tert*-butyl-1,3-dibromopyrene **2** (300 mg, 0.72 mmol), diphenylamine 2 (366 mg, 2.16 mmol), palladium(II) acetate (8.1 mg, 36 µmol), sodium *tert*-butoxide (416 mg, 4.33 mmol), and tri*tert*-butylphosphine (1M in toluene, 0.07 mL, 70 µmol) in toluene (15 mL) was heated to reflux for 12 h under argon. The mixture was chilled to room temperature and quenched with water (100 mL). The mixture was extracted with dichloromethane (3 × 50 mL), and the combined extracts were washed with water and brine, dried with MgSO₄ and concentrated. The residue was chromatographed over silica gel (Wako C-300, 200 g) with chloroform:hexane (2:8) as eluent to give a light-yellow solid. Recrystallization from ethyl acetate afforded 7-*tert*-butyl-1,3-bis(*N*,*N*-diphenylamino)pyrene **3b** (291 mg, 68%) as light yellow prisms. M.p. 235–236°C; ¹H NMR (300 MHz, CDCl₃): $\delta_{\rm H} = 1.53$ (s, 9H, *t*Bu), 6.90 (t, *J* = 7.2 Hz, 4H, Ar-*H*), 7.04 (d, *J* = 7.5 Hz, 8H, Ar-*H*), 7.16 (d, *J* = 8.0 Hz, 8H, Ar-*H*), 7.67 (s, 1H, pyrene-*H*), 7.86 (d, *J* = 9.2 Hz, 2H, pyrene-*H*), 8.06 (d, *J* = 9.2 Hz, 2H, pyrene-*H*), 8.10 ppm (s, 2H, pyrene-*H*); ¹³C NMR (100 MHz, CDCl₃): $\delta_{\rm C} =$ **31.8**, **35.2**, **121.7**, **121.8**, **122.5**, **123.1**, **123.2**, **126.8**, **127.7**, **127.9**.

129.1, 129.2, 131.1, 141.5, 148.3, 149.8 ppm; **FAB-MS (MALDI-TOF)**: *m*/*z* calcd for C₄₄H₃₆N₂ **592.2865** [M⁺]; found 592.2581 [M⁺].

Synthesis of 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-methylphenyl)amino)pyrene (3c)

A mixture of 7-*tert*-butyl-1,3-dibromopyrene **2** (300 mg, 0.72 mmol), bis(4-methylphenyl)amine (426 mg, 2.16 mmol), palladium(II) acetate (8.1 mg, 36 µmol), sodium *tert*-butoxide (416 mg, 4.33 mmol), and tri-*tert*-butylphosphine (1M in toluene, 0.07 mL, 70 µmol) in toluene (15 mL) was heated to reflux for 12 h under argon. The mixture was chilled to room temperature and quenched with water (100 mL). The mixture was extracted with dichloromethane (3 × 50 mL), and the combined extracts were washed with water and brine, dried with MgSO4 and concentrated. The residue was chromatographed over silica gel (Wako C-300, 200 g) with chloroform:hexane (2:8) as eluent to give a light-yellow solid. Recrystallization from ethyl acetate afforded 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-methylphenyl)amino)pyrene **3c** (387 mg, 82%) as light-yellow prisms. M.p. 224–225°C; ¹H NMR (300 MHz, CDCl₃): $\delta_{\rm H} = 1.52$ (s, 9H, *t*Bu), 2.25 (s, 12H, Me), 6.91 (d, *J* = 8.8 Hz, 8H, Ar-*H*), 6.96 (d, *J* = 8.4 Hz, 8H, Ar-*H*), 7.60 (s, 1H, pyrene-*H*), 7.83 (d, *J* = 9.2 Hz, 2H, pyrene-*H*), 8.06 (d, *J* = 9.7 Hz, 2H, pyrene-*H*), 8.07 ppm (s, 2H, pyrene-*H*); ¹³C NMR (100 MHz, CDCl₃): $\delta_{\rm C} = 20.7$, 31.8, 35.1, 121.7, 122.3, 123.3, 126.4, 127.2, 127.9, 128.9, 129.6, 130.8, 131.1, 142.0, 146.2, 149.6 ppm; FAB-MS (MALDI-TOF): *m*/z calcd for C48H44N₂ 648.3504 [M⁺]; found 648.3521 [M⁺].

Synthesis of 7-tert-butyl-1,3-bis(N,N-bis(4'-tert-butylphenyl)amino)pyrene (3d)

A mixture of 7-*tert*-butyl-1,3-dibromopyrene **2** (300 mg, 0.72 mmol), bis(4-*tert*-butylphenyl)amine (607 mg, 2.16 mmol), palladium(II) acetate (8.1 mg, 36 µmol), sodium *tert*-butoxide (416 mg, 4.33 mmol), and tri-*tert*-butylphosphine (1M in toluene, 0.07 mL, 70 µmol) in toluene (15 mL) was heated to reflux for 12 h under argon. The mixture was chilled to room temperature and quenched with water (100 mL). The mixture was extracted with dichloromethane (3×50 mL), and the combined extracts were washed with water and brine, dried with MgSO4 and concentrated. The residue was chromatographed over silica gel (Wako C-300, 200 g) with chloroform:hexane (3:7) as eluent to give an orange solid. Recrystallization from methanol afforded 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-*tert*-butylphenyl)amino)pyrene **3d** (588 mg, 73%) as yellow prisms. M.p. 229–230°C; ¹H NMR (300 MHz, CDCl₃): $\delta_{\rm H} = 1.27$ (s, 18H, *t*Bu), 1.29 (s, 9H, *t*Bu), 6.99 (t, *J* = 8.1 Hz, 7H, Ar-*H*), 7.07 (d, *J* = 7.6 Hz, 2H, Ar-*H*), 7.18 (t, *J* = 8.1 Hz, 7H, Ar-*H*), 7.75 (d, *J* = 7.6 Hz, 1H, pyrene-*H*), 7.86 (t, *J* = 8.6 Hz, 2H, pyrene-*H*), 8.10 ppm (m, 4H, pyrene-*H*); ¹³C NMR (100 MHz, CDCl₃): $\delta_{\rm C} =$ **31.4**, 31.8, 34.1, 35.1, 121.0, 122.3, 123.2, 123.4, 125.8, 126.9, 127.3, 127.9, 128.9, 129.3, 131.1, 134.7, 141.7, 144.1, 145.7, 149.6 ppm; FAB-MS (MALDI-TOF): *m/z* calcd for C₆₀H₆₈N₂ 816.5383 [M⁺]; found 816.5383 [M⁺].

Synthesis of 7-tert-butyl-1,3-bis(N,N-bis(4'-methoxyphenyl)amino)pyrene (3e)

A mixture of 7-*tert*-butyl-1,3-dibromopyrene **2** (300 mg, 0.72 mmol), bis(4-methoxyphenyl)amine (495 mg, 2.16 mmol), palladium(II) acetate (8.1 mg, 36 µmol), sodium *tert*-butoxide (416 mg, 4.33 mmol), and tri-*tert*-butylphosphine (1M in toluene, 0.07 mL, 70 µmol) in toluene (15 mL) was heated to reflux for 12 h under argon. The mixture was chilled to room temperature and quenched with water (100 mL). The mixture was extracted with dichloromethane (3 × 50 mL), and the combined extracts were washed with water and brine, dried with MgSO4 and concentrated. The residue was chromatographed over silica gel (Wako C-300, 200 g) with chloroform:hexane (8:2) as eluent to give an orange solid. Recrystallization from ethyl acetate afforded 7-*tert*-butyl-1,3-bis(*N*,*N*-bis(4'-methoxyphenyl)amino)pyrene **3e** (370 mg, 72%) as orange prisms. M.p. 260–261°C; ¹H NMR (300 MHz, CDCl₃): $\delta_{\rm H} = 1.52$ (s, 9H, *t*Bu), 3.74 (s, 12H, OMe), 6.72 (d, *J* = 9.2 Hz, 8H, Ar-*H*), 6.92 (d, *J* = 9.0 Hz, 8H, Ar-*H*), 7.48 (s, 1H, pyrene-*H*); ¹³C NMR (100 MHz, CDCl₃): $\delta_{\rm C} = 31.8, 35.1, 55.5, 114.5, 122.1, 123.2, 123.3, 123.4, 125.3, 126.8, 127.6, 128.0, 131.2, 142.5, 142.7, 149.5, 154.5 ppm; FAB-MS (MALDI-TOF):$ *m/z*calcd for C48H44N2O4 712.3301 [M⁺]; found 712.3286 [M⁺].

X-ray Crystallography

A suitable single crystal ($0.87 \times 0.63 \times 0.43 \text{ mm}^3$) was selected and mounted on a Bruker APEX 2 CCD diffractometer equipped with graphite-monochromated Mo-K α radiation for **3e**.²⁶ Data were corrected for Lorentz and polarisation effects and for absorption.²⁶ The structure was solved by direct methods and refined routinely except that atoms C36 > C41 & O(3) were modelled as disordered over two sets of positions with major component 51.3(3)%. Details of the crystal parameters, data collection conditions, and refinement parameters²⁷ for the compound **3e** are summarized in Tables S1–2. Crystallographic data for the structure in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication numbers CCDC 1537633. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [fax: 144-1223-336033 or e-mail: deposit@ccdc.cam.ac.uk].

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.xxxxxx.

NMR spectra, UV-vis spectra, emission spectra (PDF) Crystallographic information for **3e** (cif)

AUTHOR INFORMATION

Corresponding Author

- * E-mail: <u>yamatot@cc.saga-u.ac.jp</u>.
- * E-mail: hyxhn@sina.com

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was performed under the Cooperative Research Program of "Network Joint Research Center for Materials and Devices (Institute for Materials Chemistry and Engineering, Kyushu University)". We would like to thank the OTEC at Saga University and the International Cooperation Projects of Guizhou Province (No. 20137002), National Science Foundation of China (No. 21602014), the Scientific Research Foundation for the Re-turned Overseas Chinese Scholars, the State Education Ministry, the Fund Program for the Scientific Activities of Selected Returned Overseas Professionals of Beijing, The Scientific Research Common Program of Beijing Municipal Commission of Education (KM201510015003), The EPSRC is thanked for an overseas travel grant to C.R.

REFERENCES

1 (a) *Functional Organic Materials - Syntheses, Strategies, and Applications*, Müller, T. J. J., Bunz, U. H. F., Ed.; WileyVHC: Weinheim, **2007**. (b) Zhu, M.; Yang, C. *Chem. Soc. Rev.* **2013**, *42*, 4963–4976. (c) Maggini, L.; Bonifazi, D. *Chem. Soc. Rev.* **2012**, *41*, 211–241. (d) Reineke, S.; Rosenow, T. C.; Lüssem, B.; Leo, K. *Adv. Meter.* **2010**, *22*, 3189–3193. (e) Fan, C.; Zhu, L.; Liu, T.; Jiang, B.; Ma, D.; Qin, J.; Yang, C. *Angew. Chem., Int. Ed.* **2014**, *53*, 2147–2151.

2 (a) Kido, J.; Okamoto, Y. *Chem. Rev.* **2002**, *102*, 2357–2368. (b) Samuel, I. D. W.; Turnbull, G. A. *Chem. Rev.* **2007**, *107*, 1272–1295. (c) Grimsdale, A. C.; Leok Chan, K.; Martin, R. E.; Jokisz, P. G.; Holmes, A. B. *Chem. Rev.* **2009**, *109*, 897–1091.

3 (a) Stender, A. S.; Marchuk, K.; Liu, C.; Sander, S.; Meyer, M. W.; Smith, E. A.; Neupane, B.; Wang, G.; Li, J.; Cheng, J.-X.; Huang, B.; Fang, N. *Chem. Rev.* **2013**, *113*, 2469–2527. (b) Wang, X. H.; Guo, Z. Q.; Zhu, S. Q.; Liu, Y. J.; Shi, P.; Tian, H.; Zhu, W. H. J. Mater. Chem. B **2016**, *4*, 4683–4689.

4 (a) Zhang, R.; Zhao, Y.; Zhang, T. F.; Xu, L.; Ni, Z. H. Dyes Pigments 2016, *130*, 106–115.
(b) Lee, S. Y.; Yasuda, T.; Yang, Y. S.; Zhang, Q. S.; Adachi, C. *Angew. Chem., Int. Ed.* 2014, *53*, 6402–6406. (c) Han, M. G.; Park, K. -B.; Bulliard, X.; Lee, G. H.; Yun, S.; Leem, D. -S.; Heo, C. -

J.; Yagi, T.; Sakurai, R.; Ro, T.; Lim, S. -J.; Sul, S.; Na, K.; Ahn, J.; Jin, Y. W.; Lee, S. ACS Appl. Mater. Interfaces 2016, 8, 26143–26151.

5 (a) Shen, X.; Wang, Y.; Zhao, E.; Yuan, W.; Liu, Y.; Lu, P.; Qin, A.; Ma, Y.; Sun, J.; Tang,
B. Z. J. Phys. Chem. C 2013, 117, 7334–7347. (b) Xue, v; Liu, W.; Qiu, X.; Gao, Y.; Yang, W. J.
Phys. Chem. C 2014, 118, 18668–18675.

6 (a) Zhu, L. M.; Xu, J. C.; Sun, Z.; Fu, B. Q.; Qin, C. Q.; Zeng, L. T.; Hu, X. C. *Chem. Commun.* **2015**, *51*, 1154–1156. (b) Teran, N. B.; He, G. S.; Baev, A.; Shi, Y. R.; Swihart, M. T.; Prasad, P. N.; Marks, T. J.; Reynolds, J. R. *J. Am. Chem. Soc.* **2016**, *138*, 6975–6984.

7 Saigusa, H.; Lim, E. C. Acc. Chem. Res. 1996, 29, 171-178.

8 Hsieh, C. C.; Jiang, C. M.; Chou, P. T. Acc. Chem. Res. 2010, 43, 1364–1374.

9 Levi, L.; Müller, T. J. J. Chem. Soc. Rev. 2016, 45, 2825-2846.

10 (a) Khramov, D. M.; Bielawski, C. W. J. Org. Chem. 2007, 72, 9407–9417. (b) Feng, X.; Tomiyasu, H.; Hu, J. Y.; Wei, X. F.; Redshaw, C.; Elsegood, M. R.J.; Horsburgh, L.; Teat, S. J.; Yamato, T. J. Org. Chem. 2015, 80, 10973–10978. (c) McNamara, L. E.; Liyanage, N.; Peddapuram, A.; Murphy, J. S.; Delcamp, J. H.; Hammer, N. I. J. Org. Chem. 2016, 81, 32–42.

11 Figueira-Duarte, T. M.; Müllen, K. Chem. Rev. 2011, 111, 7260-7314.

12 Misra, R.; Gautam, P.; Mobin, S. M. J. Org. Chem. 2013, 78, 12440-12452.

13 (a) Oseki, Y.; Fujitsuka, M.; Sakamoto, M.; Majima, T. J. Phys. Chem. A 2007, 111, 9781–9788. (b) Yang, S. W.; Elangovan, A.; Hwang, K. C.; Ho, T. I. J. Phys. Chem. B 2005, 109, 16628–16635. (c) Sessler, J. L.; Karnas, E.; Kim, S. K.; Ou, Z. P.; Zhang, M.; Kadish, K. M.; Ohkubo, K.; Fukuzumi, S. J. Am. Chem. Soc. 2008, 130, 15256–15257.

14 Feng, X.; Hu, J. Y.; Redshaw, C.; Yamato, T. Chem. Eur. J. 2016, 22, 11898-11916.

15 Jérome, D.; Schulz, H. J. Advances in Physics 2002, 51, 293-479.

16 (a) Xiao, L. X.; Chen, Z. J.; Qu, B.; Luo, J. X.; Kong, S.; Gong, Q. H.; Kido, J. Adv. Mater.
2011, 23, 926–952. (b) Ying, L.; Ho, C-L.; Wu, H. B.; Cao, Y.; Wong, W-Y. Adv. Mater. 2014, 26, 2459–2473.

17 (a) Xia, H.; Liu, D. Q.; Song, K. S.; Miao, Q. Chem. Sci. 2011, 2, 2402–2406. (b) Wang, C.
Z.; Kihara, R.; Feng. X.; Thuéry, P.; Redshaw, C.; Yamato, T. ChemistrySelect 2017, 2, 1436–1441.

18 (a) Shizu, K.; Tanaka, H.; Uejima, M.; Sato, T.; Tanaka, K.; Kaji, H.; Adachi, C. J. Phys. Chem. C 2015, 119, 1291–1297. (b) Rajamalli, P.; Senthilkumar, N.; Gandeepan, P.; Huang, P. Y.; Huang, M. J.; Ren-Wu, C. Z.; Yang, C. Y.; Chiu, M. J.; Chu, L. K.; Lin, H. W.; Cheng, C. H. J. Am. Chem. Soc. 2016, 138, 628–634.

19 Sasabe, H.; Kido, J. J. Mater. Chem. C 2013, 1, 1699-1707.

20 Feng, X.; Hu, J. Y.; Tomiyasu, H.; Tao, Z.; Redshaw, C.; Elsegood, M. R. J.; Horsburgh, L.; Teat, S. J.; Wei, X. F.; Yamato, T. *RSC Adv.* **2015**, *5*, 8835–8848.

21 (a) Jelley, E. E. Nature **1936**, 138, 1009–1010. (b) Kasha, M.; Rawls, H. R.; Ashraf El-Bayoumi, M. Pure Appl. Chem. **1965**, 11, 371–392. (c) Lanzani, G. Photophysics Behind Photovoltaics and Photonics Wiley, **2002**. (d) Heyne, B. Photochem. Photobiol. Sci. **2016**, 15, 1103– 1114.

22 Snow, A. W. The Porphyrin Handbook, 2003,17. 129–176.

23 (a) Lippert, V. E.; Z. Naturforsch. A: Phys. Sci. 1955, 10, 541–545. (b) Mataga, N.; Kaifu,
Y.; Koizumi, M. Bull. Chem. Soc. Jpn. 1956, 29, 465–470.

24 (a) Jadhav, T.; Dhokale, B.; Patil, Y.; Mobin, S. M.; Misra, R. J. Phys. Chem. C 2016, 120, 24030–24040. (b) Tayade, R. P.; Sekar, N. J. Lumin. 2016, 176, 298–308. (c) Lim, C-H.; Ryan, M. D.; McCarthy, B. G.; Theriot, J. C.; Sartor, S. M.; Damrauer, N. H.; Musgrave, C. B.; Miyake, G. M. J. Am. Chem. Soc. 2017, 139, 348–355.

25 Feng, X.; Hu, J. Y.; Yi, L.; Seto, N.; Tao, Z.; Redshaw, C.; Elsegood, M. R. J.; Yamato, T. *Chem. - Asian J.* **2012**, *7*, 2854–2863.

26 APEX 2 & SAINT. Software for CCD Diffractometers. Madison, USA: Bruker AXS Inc;2012.

27 Sheldrick, G. M. Acta Crystallogr. 2015, C71, 3-8.