

Physical modelling of water, fauna and flora: Knowledge gaps, avenues for future research and infrastructural needs

Robert E. Thomas¹, Matthew F. Johnson², Lynne E. Frostick¹, Daniel R. Parsons¹, Tjeerd J. Bouma³, Jasper T. Dijkstra⁴, Olivier Eiff⁵, Sylvie Gobert⁶, Pierre-Yves Henry⁷, Paul Kemp⁸, Stuart J. McLelland¹, Frederic Y. Moulin⁵, Dag Myrhaug⁷, Alexandra Neyts⁷, Maike Paul⁹, W. Ellis Penning⁴, Stephen P. Rice², Sara Puijalon¹⁰, Adrian Stanica¹¹, Davide Tagliapietra¹², Michal Tal¹³, Alf Tørum¹⁴, Michaelis I. Vousdoukas⁹

1. Abstract

In recent decades, the field of eco-hydraulics has grown to cover wide-ranging research into the complex interactions between aquatic organisms and hydraulics. Understanding of these interactions is important for the management of aquatic environments, prediction of the impacts on them as a result of environmental change, and our ability to exploit their ecosystem services. Physical modelling is a key method for generating understanding of these interactions, and as such is central to our ability to predict future trajectories of aquatic environments. However, many aspects of this field remain poorly understood and the use of physical models within eco-hydraulics requires significant advancement in both methodological application and substantive understanding.

This paper presents a review of the emergent themes from a workshop held amongst participants of the PISCES work package of the EC-funded HYDRALAB IV project and a group of invited eco-hydraulic experts. The identified themes are: 1.) abiotic factors; 2.) adaptation and behaviour; 3.) complexity and feedback; 4.) heterogeneity; and 5.) scale and scaling. The paper examines these themes and identifies how progress on each of them is key to existing and future

¹ Department of Geography, Environment and Earth Sciences, University of Hull, Cottingham Road, Hull. HU6 7RX. UK.

² Department of Geography, Loughborough University, Loughborough, Leicestershire, LE11 3TU, UK.

³ Nederlands Instituut voor Ecologie, Postbus 140, 4400 AC Yerseke, Netherlands.

⁴ Deltares, Rotterdamseweg 185, NL-2629 HD Delft, Netherlands.

⁵ Institut de Mécanique des Fluides, UMR CNRS/INP-UPS 5502, Allée du Professeur Camille SOULA, 31400 Toulouse, France.

⁶ Laboratoire d'Océanologie – Centre MARE, Université de Liège, Allée de la Chimie 3, 4000 Liège (Sart-Tilman), Belgium.

⁷ Department of Marine Technology, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway.

⁸ Engineering and the Environment, University of Southampton, Highfield, Southampton, SO17 1BJ. UK.

⁹ Leibniz Uni Hannover, Forschungszentrum Küste, Merkurstr. 11, 30419 Hannover, Germany.

¹⁰ UMR CNRS 5023 “Ecologie des Hydrosystèmes Naturels et Anthropisés”, Université Lyon 1, 43 Boulevard du 11 novembre 1918, 69622 Villeurbanne Cedex, France.

¹¹ GeoEcoMar, Str. Dimitrie Onciul 23-25, sector 2, 024053 Bucharest, Romania.

¹² CNR, ISMAR, Arsenale – Tesa 104, Castello 2737/F, 30122 Venezia, Italy.

¹³ Aix-Marseille Université, CEREGE UMR 7330, Europôle de l'Arbois, BP 80, 13545 Aix-en-Provence Cedex 04, France.

¹⁴ Department of Civil and Transport Engineering, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway.

efforts to progress our knowledge and understanding of eco-hydraulic interactions. Illustrative examples are drawn from studies on biofilms, plants, and sessile and mobile fauna in shallow water fluvial and shallow? marine environments. Examples of research gaps, suggestions for future research and directions for educational, infrastructural and technological advance are also presented.

Keywords: Biofilms, Biogeomorphology, Eco-hydraulics, Experimental facilities, Flow-biota interactions, Macrozoobenthos, Vegetated flows

2. Introduction

The complex interactions between organisms and hydraulics are fundamental to the understanding and management of aquatic environments. The recent growth of the field of eco-hydraulics (e.g. Nikora 2010, Rice *et al.* 2010a, 2010b) reflects the imperative of predicting the consequences of current and future environmental change (Solomon *et al.* 2007, Parry *et al.* 2007, Thorne *et al.* 2007). Changes such as sea-level rise, increased storminess and conveyance-related flooding (Parry *et al.* 2007, Thorne *et al.* 2007) can, in turn, be expected to force changes to the dynamic interactions between ecology and hydraulics. Physical modelling provides an essential bridge between field observations and theoretical, stochastic and numerical models that attempt to predict the impact of environmental changes on aquatic ecosystems.

There are long-known and obvious limitations to the application of physical modelling to real world situations, many of which are centred on issues of scale. One important issue is the different timescales between the experimental period available in the laboratory and those over which a process might act in the natural environment. This is particularly important to bear in mind when considering the long-term response of aquatic systems to climate change. Another issue is that of scaling the experimental domain and process timescales to the hydraulic facility. This is particularly important in sediment transport and morphodynamic experiments since there is a danger of obtaining the right morphology for the wrong reasons (*cf.* Beven 1989, Favis-Mortlock *et al.* 1996). Added to this, the scaling of both sediment and plant materials is problematic since the cohesive and frictional components of sediment shear strength do not scale linearly (Peakall *et al.* 2007) and neither do the tensile and frictional components of plant root strength (Pollen and Simon 2006). Nevertheless, if the above limitations are mitigated against and controlled during the design of experiments, the incorporation of plants and animals into hydraulic facilities can yield important data and provide significant insight, particularly on the detailed interactions between physical and biological processes.

Recent progress in eco-hydraulic experimentation has been relatively rapid (e.g. Nikora 2010, Rice *et al.* 2010a, Nepf 2012). However, there are still many aspects of this field that are in their infancy and within this context it is significant that the introduction of biota is impossible in many of our hydraulic

facilities. As a result, experiments have to rely on inert surrogates, often plastics or wood, which simplify responses but introduce issues of complexity and representation. Other issues relate to the fact that eco-hydraulic experimental research is truly interdisciplinary, requiring contributions from ecology, environmental sciences and hydraulic engineering/physics. It is challenging to access information across these disciplinary boundaries because of differing research methods, aims, and terminology (Rice *et al.* 2010a). Given the problems it is not surprising that our understanding of the interactions between flow and biota are still at an early stage of development.

2.1. Issues and difficulties associated with understanding the interactions between flows and biota

The measurement of turbulent fluid flow fields is a challenging problem even before considering their interaction with mobile objects, deformable boundaries and/or biology. Turbulent flows are highly variable in all three spatial dimensions and in time and exhibit a range of spatial and temporal scales and quasi-periodicities from very small lengths (the Taylor or Kolmogorov microscales) and short durations to considerably longer lengths (up to the thickness of the boundary layer) and durations. Thus, velocity measurements should be undertaken at high spatial and temporal resolution, but simultaneously cover a large spatial domain and sample for long periods of time. These dichotomous requirements, compounded by data storage limitations, help to make controlled laboratory environments especially attractive venues for the study of turbulent flows.

However, introducing biological agents, such as biofilms, plants, and sessile and mobile vertebrate and invertebrate animals, into the laboratory presents significant additional challenges. Accounting for the additional degrees of freedom afforded by the interactions of turbulent flows with features immersed within the fluid and with the boundaries within which they are constrained is non-trivial. Aquatic fauna and flora are capable of modifying both time- and space-averaged as well as turbulent fluid flows, but conversely, fluid flows are also capable of changing the form and behaviour of deformable and reactionary biota. Furthermore, there are other, potentially unrelated, factors which may drive or at least modify the behaviour of biota.

In attempting to understand the interactions between flows and biota, some parallels may be drawn with the experience of geomorphologists and geologists attempting to understand long-term and large-scale landscape evolution. In his 1991 book “To Interpret the Earth: Ten ways to be wrong”, Stanley Schumm (1991: 35) identified ten difficulties associated with using modern conditions as a basis for extrapolation:

- “1. Time, involving both absolute duration and relative time spans; 2. Space, involving size and scale; 3. Location, the site of concern within a natural system; 4. Convergence (equifinality; Chorley, 1962), the production of similar results from

different processes and causes; 5. Divergence, the production of different results from similar processes and causes; 6. Efficiency, the variable efficiency and work accomplished by a process; 7. Multiplicity, the multiple explanations that combine to influence and cause natural phenomena; 8. Singularity (or Heterogeneity), the natural variability among like things; 9. Sensitivity, the susceptibility of a system to change; and 10. Complexity, the complex behaviour of a system that has been subject to altered conditions”.

Schumm (1991) suggests that these difficulties may be grouped into three broad classes that are based upon 1. Scale and place, 2. Cause and effect, and 3. System response. He notes that time, space and location are all associated with the identification of the required resolution of observations and the necessary spatial and temporal extents of study. In contrast, convergence, divergence, efficiency and multiplicity are associated with variations from a simple cause and effect relationship between two or more factors. Finally, singularity, sensitivity and complexity are problems that involve system characteristics and response to change.

Many of the difficulties identified by Schumm (1991) were also identified by the experts who attended an eco-hydraulic foresight workshop held in Trondheim on 15th September 2011. This was organised as part of HYDRALAB IV, an Integrated Infrastructure Initiative that is part of the European Community’s 7th Framework Programme. This includes PISCES, a joint research activity designed to improve physical modelling of eco-hydraulic systems. The foresight team aim to identify the probable future infrastructure requirements of the next generation of engineers and scientists working within the broad field of environmental hydraulics. Anticipation of those requirements is essential if we are to respond in a timely fashion to the challenges that face society as freshwater and coastal-marine systems respond to environmental change. In the sections that follow, the dominant issues undermining existing and future efforts to further understand eco-hydraulic interactions are described and discussed:

- Abiotic Factors: the detection of, reaction to, and modification of a number of environmental factors that may be dependent on or independent of the flow field by subaqueous plants and animals;
- Adaptation and Behaviour: the adaptation and behavioural modifications of organisms at different temporal scales;
- Complexity and Feedback: complex interactions between organisms and the hydrodynamic environment and the role of feedback, whether positive or negative, in amplifying or moderating organism or environmental response, respectively;

- Heterogeneity: “the natural variability among like things” (Schumm 1991); and
- Scale and Scaling: is it possible to scale down biological (and biomechanical) processes operating at the large scale, are the variables measured at the large scale pertinent at the small scale and does technology permit us to measure the same variable across scales?

Note that while each of these issues may be viewed as being distinct and different, it is also clear that they overlap and interact with one another. In each section below, each issue is briefly discussed and then some potential avenues for research are described. These examples are drawn from studies upon biofilms, plants, mobile benthic invertebrates and sessile macrozoobenthos that populate shallow water fluvial and marine environments, reflecting the areas of expertise of the contributors to the workshop. The improvements to technologies and infrastructures necessary to tackle these issues are then outlined.

3. Abiotic Factors

3.1. Theme Description

The behaviour of subaqueous plants and animals may be modified by a number of environmental factors that may be dependent on or independent of the flow field around them. Biota may sense and react to variations in acoustic, electro-chemical, optical, and thermal stimuli that may be beyond the ability of humans to detect and may presently also be beyond the ability of instruments to detect. Examples of abiotic factors (see the recent review of Bornette and Puijalon 2011) include light, sound, carbon dioxide, oxygen, nutrients (e.g. nitrogen, phosphorus, and sulphur), the presence/absence of heavy minerals, salinity, pH, organic matter (both within the substrate and the water column), sediment composition (both within the substrate and the water column), turbidity and temperature. Some of these factors are inter-related. Furthermore, the responses of biota may be physiological and/or chemical rather than physical and thus may also prove difficult to detect.

3.2. Examples of knowledge gaps

The impact of biological activity upon substrate and suspended sediment physico-chemical characteristics is presently an under-studied area. For example, while there is an extensive body of literature describing the nutrient and oxygen exchange promoted by passive irrigation and pumping of Thalassinidean shrimp (e.g. Aller *et al.* 1983, Suchanek 1983, Ziebis *et al.* 1986, Astall *et al.* 1997, Griffen *et al.* 2004, D’Andrea and DeWitt 2009), few studies have extended to study the chemical composition of the sediment surrounding burrows and the role of chemical exchanges in modifying the chemical balance within the substrate, especially under different wave-current regimes. Furthermore, studies of this type need to be extended to a wider range of species, particularly mobile animals and plants. Macrophytes, for instance, often form in clustered groups of individuals of one or more species on the beds of rivers. The deposition of sediment and organic matter is promoted within

these ‘patches’ (Schoelynck *et al.* 2012), making them biogeochemical hotspots (Schoelynck 2011). There is also growing evidence that Si concentration in aquatic species litter is a major factor driving decomposition rates (Schaller *et al.* 2012a, Schaller *et al.* 2012b), which can impact upon ecosystem productivity and nutrient availability and thus affect vascular plant distribution (Bragazza and Gerdol 2002). Unfortunately, laboratory studies on plants face practical constraints. For example, the root networks of some plants are too anoxic for use in the lab, while others (e.g. *Posidonia oceanica*) are thought to fix oxygen into the substrate, improving growing conditions. However, it is not currently possible to transplant *P. oceanica*, mainly because the uptake of nitrogen by leaves is insufficient to replenish the N lost during natural leaf decay (Lepoint *et al.* 2004). In addition, although *P. oceanica* may fix oxygen, it is still sensitive to substrate oxygen levels, has deep roots and has high stem densities that are difficult to replicate (Gobert *et al.* 2003). Further research is thus needed to identify and isolate the environmental parameters and tolerance ranges to which *P. oceanica* is sensitive and/or to identify other species that may be more resilient. Such studies are complicated by the difficulties presented by chemical instabilities within the substrate, associated with limited sediment thicknesses, planform areas and ground (pore) and overlying water chemical imbalances within experimental facilities. The latter issue may be ameliorated by using riverside flumes and/or unfiltered water. At a broader spatial scale, the *relative* importance of the decay and exchange of organic and inorganic matter, salinity and hydrodynamics in modifying the spatial distribution of individuals, and/or communities of organisms in estuarine and lagoonal environments is yet to be fully established.

Due to the complex and important controls that environmental variables exert over the behaviour and morphology of living organisms, it is intuitive to assume that organisms will be sensitive to changes in these parameters. Organisms may respond to subtle alterations to environmental variables and at levels below those that may cause stress. However, many aquatic species tolerate a wide range of conditions in the field and thus have global distributions covering a range of climates, habitats and environmental conditions. For instance, signal crayfish (*Pacifastacus leniusculus*) are native to the north-western USA, but now thrive throughout Europe and Japan. Many non-marine aquatic plants also have extremely wide distributions, with climatic factors seemingly having minimal effect on their distribution (Santamaria 2002). This implies that many plants and animals can survive a large range of environmental conditions, potentially lessening the challenge of maintaining organisms in flumes. However, whilst great ranges of conditions are ‘survivable’, they may still have important consequences for flume work as organism behaviour and morphology can be largely controlled by environmental conditions. For instance, whilst many species of mollusc have colonies all over the world, those in areas of high hydraulic stress are likely to be morphologically dissimilar to those from sheltered areas (e.g., blue mussels (*Mytilus edulis*): Seed 1969, Jørgensen 1976; limpets (genera *Cymbula* and *Scutellastra*): Branch and Marsh 1978; brown mussels (*Perna perna*): McQuaid *et al.*

2000, McQuaid and Lindsay 2000; Mediterranean mussels (*Mytilus galloprovincialis*): Steffani and Branch 2003, Hammond and Griffiths 2004; fan mussels (*Pinna nobilis*): García-March *et al.* 2007). Similarly, plants within a species can be morphologically very different depending on the environment within which they are growing (e.g., Idestam-Almquist and Kautsky 1995, Strand and Weisner 1996, Doyle 2001, Puijalon and Bornette 2006, Puijalon *et al.* 2008a). Likewise, at the annual scale, external factors other than hydrodynamics may trigger changes to biofilm composition (de Brouwer *et al.* 2000). The effects of light attenuation, clogging of pore spaces, abrasion and fluid drag on biofilm dynamics have still not been isolated. Consequently, there is a need for research that identifies important organism stresses, beyond just the tolerable range, and a discussion of which environmental conditions need to be replicated. In addition, consideration needs to be given to organisms with a wide spatial distribution and the implications this has for the selection of experimental parameters. For example, when using blue mussels should the characteristics of the water (e.g. temperature, salinity, pH) in Norway or Chile be replicated? Much relevant information could be gained from existing biological literature and the answers to many such questions will depend on the specific aims of the study. For most studies, replicating the conditions at the site of interest would be sufficient, but an understanding of which conditions need to be replicated in flumes to ensure organism behaviour is analogous to field equivalents would be beneficial to experimental design.

4. Adaptation and behaviour

4.1. Theme Description

Living organisms adapt to the natural environment in which they are immersed at temporal scales ranging from supra-millennial (genetic mutation or evolution) to sub-second reconfigurations or behavioural adjustments. These adjustments may be classified as avoidance or tolerance and may take the form of changes to growth, movement, feeding and reproduction strategies. These adaptations may be in response to hydraulic drivers (i.e., the mean current, whether unidirectional or multidirectional; waves; turbulence; boundary and/or wall effects) or they may be predominantly in response to abiotic drivers (see section 3). Alternatively, adaptations may result from the superimposition of abiotic drivers on hydraulic drivers. In addition to the adaptations made by individuals, some species also form colonies that may enable both the individuals within the colony and the colony as a whole to obtain a collective advantage (e.g., within colonies the ascidians *Botryllus schlosseri* inhale water independently but exhale into a common exhalant cavity, increasing the flow rate and enhancing the efficiency of waste removal; Vogel 1994). Further, some groups of species tend to associate with each other because of either an observed or a presumed competitive advantage afforded by cohabitation (e.g., goose barnacles (*Capitulum mitella*) require a firm substrate, which may be supplied by the shells of mussels (*Septifer virgatus*). This increases the roughness of

shells, reducing wave-induced shear stress and thus decreasing mussel entrainment; Kawai and Tokeshi 2004). The challenge faced by the scientific community is to observe and measure these adaptations and behavioural changes in an objective manner, both for individual specimens and entire colonies.

4.2. Examples of knowledge gaps

Without detailed biological and ecological knowledge of organism activity and behaviour in field situations, it is impossible to assess the extent to which the behaviour exhibited by an organism removed from its natural habitat is “natural”. Thus, although it is difficult to assess what constitutes a “natural” behaviour of a wild organism in an undisturbed field situation, a first goal must be to obtain baseline information on natural behaviours. Incorporating a field validation component to flume experiments is also an effective way of assessing the realism of the response of biota to stimuli in laboratory flumes as field experiments integrate across biological interactions. Once this information has been obtained for the organism of interest, a second goal is to assess how analogous the behaviour of organisms in flume environments is to field equivalents. In addition, it would be beneficial to understand whether organisms respond differently to a ‘natural stress’, such as a flood event, than to an ‘unnatural stress’ that is specifically associated with artificial environments, such as a lack of space or the imposition of a diurnal signature through pausing experiments each night. If tell-tale signs of these stresses could be identified it would be extremely beneficial to future studies.

Acclimatising organisms to experimental conditions is important so as to avoid mistakenly believing the response of an organism to the new environment is actually the organism response to experimental variables (e.g., Barmuta *et al.* 2001, Battin *et al.* 2003, Hurtado *et al.* 2011, National Research Council Committee for the Update of the Guide for the Care and Use of Laboratory Animals 2011). However, the life-history of an organism partially dictates its response to environmental and biological conditions. Therefore, the environment in which the organism has been kept prior to experimentation is likely to impact upon experimental results. This has a number of implications. First, organisms used in experiments may either be obtained from the field, raised in a laboratory or collected in the field and kept in a laboratory environment until used in experiments. There is evidence that the behaviour of animals raised in laboratories differs from field-collected equivalents (Tiselius *et al.* 1995) but further research is needed to determine whether organisms raised in these different manners behave and respond to environmental stimuli in the same way. Second, maintaining organisms in experimental conditions for extended periods may increase the likelihood of unnatural behaviours by organisms. Signal crayfish (*P. leniusculus*) that had been collected from the field but maintained in laboratory conditions rapidly altered their behaviour (Rice *et al.* 2012), presumably due to the standardisation of conditions and lack of biological interactions (e.g. lack of predation and/or competition). Research is needed to determine whether these findings are applicable to other species.

In addition to research on individual organisms, the impact of communities of organisms, rather than descriptions of the impact of individuals and/or single species on near-bed hydraulics, needs to be explored. For example, growing evidence (e.g. Amyot and Downing 1997, Rowden *et al.* 1998, Widdows *et al.* 1998, 2000a, 2000b, 2004, Andersen 2001, Widdows and Brinsley 2002, Fortino 2006) suggests that communities of animals can control substrate stability, with seasonal cycles in community structure controlling seasonal cycles in substrate stability. In addition, Merz (1984) noted that dense aggregations of polychaete worms can form hummocks. Hummocking has also been observed in dense aggregations of barnacles and mussels, with studies suggesting that the topography of hummocks results in organic matter being transported more evenly across colonies than topographically flat colonies. To our knowledge, the hydrodynamics of hummocked colonies has not been studied directly (although see the studies of flows over self-organised mussel beds and “strings” by Folkard and Gascoigne 2009 and Widdows *et al.* 2009, respectively), despite this evidence that flow alterations are of biological significance. The fact that multiple, unrelated, organisms hummock would suggest that it is of considerable significance for colony organisms. In addition, sessile animals such as ascidians, barnacles and anemones are associated with many organisms that attach to, or live within, reefs and aggregations (e.g. algae, molluscs and sponges) that may add considerable roughness and hence increase the drag acting on their hosts. Also, the presence of other organisms will increase competition which could lead to behavioural and physiological alterations. For example, dense communities of epiphytes (e.g. black fly larvae) may weigh down the leaves of plants and thus deprive their hosts of light, reduce dissolved gas exchange, affect their biomechanics and thus modify how they respond to forcing. Plant structure and morphology is known to change over the order of days/weeks in response to hydrodynamic forcing (Puijalon and Bornette 2004, 2006, Puijalon *et al.* 2005, 2008a, 2008b), but the extent to which these changes provide potential competitive advantages and/or disadvantages (e.g. the invasion of other species if plant size reduces) is yet to be established. Each of these points emphasises that the interactions between communities of organisms and environments are complex and variable and require the expertise of biologists and physical scientists to unravel the complex bi-directional impacts. Organisms respond and react to each other through purely biological interactions, in addition to interactions with each other based on habitat modification (ecosystem engineering). These interactions control the community structure and, consequently, the impacts on the physical environment. However, the nature of the physical environment will also impact the biological interactions between organisms.

5. Complexity and Feedback

5.1. Theme Description

Natural systems are inherently complex. Complexity arises in many different and sometimes unexpected ways, through the myriad interactions between biofilms, plants and animals and the

aquatic environment, including water and sediment physico-chemistry. These often non-linear interactions are further complicated by feedback, whether positive or negative. King (1970: 147) describes the two types of feedback:

“...positive feedback results in the operating process further extending the change it has induced in the dependent variable... Negative feedback, on the other hand, causes a self-regulating effect that reverses the change induced by the action of the process. The first effect is self-generating, the second is self-regulating.”

In recent years, much work has been done on feedbacks in ‘biogeomorphology’ and/or ‘ecogeomorphology’ (e.g., Fisher *et al.* 2007, Reinhardt *et al.* 2010, Rice *et al.* 2010b, Hession *et al.* 2010, Darby 2010, Wheaton *et al.* 2011, Schoelynck *et al.* 2012).

5.2. Examples of knowledge gaps

To date, experimental studies have focused on a small group of species for each ecosystem. In some circumstances (e.g. sea grasses; Fonseca and Fisher 1986, James and Barko 2000), transfer of results across species is limited and thus work needs to be expanded to a broader range of species, with targeted studies to identify organism characteristics that allow comparison of eco-hydraulic interactions across species. In addition, work is needed to identify the roles of individuals within diverse ecosystem functions (see section 4), including their responses to hydrodynamic forcing.

Feedback operates at a range of spatial and temporal scales. For example, most biofilms, plants and benthic animals reconfigure in response to hydrodynamic loading. Changes in the projected areas of organisms then reduce the hydrodynamic loading upon the organisms (negative feedback) and simultaneously impact upon the flow field. Adjustment mechanisms and the effects of those mechanisms vary by species, and furthermore, vary dynamically in response to turbulent fluctuations (Hedden *et al.* 1995). Thus, as argued by Lauder (2011), in order to properly quantify the interactions between turbulent fluid flows and the biota immersed in them, we must make simultaneous measurements of the flow field and the four-dimensional motion of biota. To date, most concurrent measurements of biological motion and velocities have been made around fish. For example, Sakakibara *et al.* (2004) used both conventional 2D and stereoscopic (3D) Particle Image Velocimetry (PIV) to quantify the flow field and shedding vortex structures around the tail fin of a goldfish, while employing a three-dimensional shape identification technique to estimate the location and shape of the fish relative to the measurement planes. Very recently, Cameron *et al.* (in review) developed a 3D PIV system to quantify the flow field and coupled it with a weighted cross spectral density-based algorithm to detect the 2D motion of a *Ranunculus penicillatus* patch in the field. Nevertheless, it is not easy to reconstruct organism topologies or three-dimensional turbulent flow fields from planar data. Recent advances in high speed camera and synchronisation technologies to capture hundreds of

positions at five to ten times the frequency of organism motion, allied to advances in tomographic, holographic, defocusing (Arroyo and Hinsch 2008) and scanning (Albagnac *et al.* 2013) 3D/3C PIV systems that measure temporal variations of all three velocity components within a finite volume, should help circumvent this problem. Scanning techniques are particularly well-adapted for high-resolution hydraulic measurements and do not impose additional optical access constraints.

At a broader scale, mobile animals alter the topographic roughness of the bed (e.g., Jones and Jago 1993, Ziebis *et al.* 1996), which in turn alters the velocity profile. Evidence suggests (Wright *et al.* 1997, Peine *et al.* 2009) that in most low-energy, fine-grained substrates, animals are the principle phenomenon altering roughness. However, the hydrodynamic impact of the roughness generated by mobile animals has rarely been quantified. For example, it has been widely noted that deposition of fine-grained sediment is promoted in pits that are constructed by a range of animals, including worms, bivalves, crustaceans and gastropods (Yager *et al.* 1993), but to our knowledge, no study details the alteration to flow characteristics created by biogenically constructed pits. Similarly, Thalassinidean shrimp burrows can cover large areas in cones and funnels with diameters hundreds of millimetres wide (Dworschak 1987). Each burrow is also subject to both passive irrigation and pumping by resident shrimp. Therefore, it seems likely that the hydrodynamic conditions over shrimp colonies are altered, but little is known of this alteration, especially in comparison to the biological literature focusing on nutrient and oxygen transfers, ecosystem engineering and functional behaviour of Thalassinidean shrimp.

At an even broader scale, vegetation colonising the intertidal zone has been found to reduce erosion, enhance deposition and stabilise the coastline, thus improving growing conditions (e.g. Bouma *et al.* 2005). Bouma *et al.* (2005) described both areal density (stems m⁻²) and scale-dependent feedbacks that were manifest through altered sedimentation patterns. Whether living plants or dead leaves, seagrass meadows act as ecosystem engineers (*sensu* Jones *et al.* 1994, van der Heide *et al.* 2007). In the Mediterranean, leaf litter commonly gets washed up on beaches, with up to 2 m of vertical accumulation, significantly reducing wave impact and erosion caused by waves and currents (Simeone and De Falco 2012). However, it remains unclear whether i) flow and turbulence fields are always modified in a way that reduces erosion and/or enhances deposition and ii) there are particular species, assemblages of species, stem, leaf and stand configurations and/or areal extents needed to sustain (Coops *et al.* 1996) and attenuate hydrodynamic forcing. In addition, further study is needed to document and quantify the complex response of the coastal ecosystem when leaf litter is removed. To what extent do these actions promote the erosion of beaches, increase sedimentation down-current, and in turn reduce the density of sea grass meadows?

6. Heterogeneity

6.1. Theme Description

In this paper, the term “heterogeneity” is considered *sensu* Schumm (1991): “the natural variability among like things”. In other words, unless (part of) an individual of a particular species has been reproduced asexually (i.e. cloned), it cannot be expected to be either physically or behaviourally identical to another (part of an) individual of the same species. For example, the structure and function of the stems and leaves of individual *R. penicillatus* plants vary depending upon the extent of submergence. Seagrasses exhibit spatio-temporal variations in stem areal density at the scale entire meadows (e.g. Kelly *et al.* 2001) but are also formed of heterogeneous patches of variable size and shape that may force heterogeneity in the mean and turbulent flow fields (Maltese *et al.* 2007). Likewise, while many landscape features appear to look and behave in the same manner, and indeed may share physical characteristics, none can be expected to be identical. In the 1:80 scale braided river experiments of Tal and Paola (2007, 2010), alfalfa seeds were initially distributed uniformly across a flume bed composed of uniform sediment with a constant water table elevation and with constant, uniform lighting. However, the resulting shoot growth patterns, and the behaviour of the braided channels, were highly heterogeneous; heterogeneity in form emerged from apparent homogeneity (or at most, very subtle heterogeneities (e.g. elevation or seed location (planform) differences of the order of a few grains in size)) because of heterogeneity in process (flow and sediment dynamics). Heterogeneity may thus arise in a number of different ways, but a key challenge is how experimentalists account for it not only within experimental design but also while interpreting experimental results.

6.2. Examples of knowledge gaps

In nature, the aquatic environment at a particular location and at an instant in time is controlled by many interacting external factors. A change to one of those factors, such as variability in upstream or downstream flow conditions induced by diurnal, seasonal or annual changes in tides, waves and/or currents, results in changes to the local hydraulics. Conversely, within a laboratory flume these external factors, or boundary conditions, are imposed depending upon the specific aims of the experimental study. Boundary conditions imposed at the inlet, outlet, bed and flume walls propagate and dissipate within flumes. At the inlet, biases introduced by pumps and incompletely removed by dampers may have little resemblance to real world conditions and may render experimental results meaningless. Likewise, in contrast to the real world, secondary currents of the second kind (Prandtl 1952) are generated adjacent to where the flume walls meet the flume bed. Thus, a key question is how inherited homogeneities and heterogeneities in boundary conditions propagate and dissipate and what impact they have upon the validity of experimental results.

To date, laboratory experiments on the hydrodynamics of vegetation have generally been performed under constant flow depths with spatially uniform vegetation patterns in a monoculture or with

artificial plants where all surrogates are identical. Similarly, the hydrodynamics of polychaete tube lawns has mainly been studied using surrogate tubes arranged on a regular grid (e.g. Friedrichs *et al.* 2000) in contrast to natural lawns that form clumps and irregular patterns, creating a more heterogeneous environment. This approach has the advantage that the number of variables is reduced and input parameters can be controlled more easily. Conversely, it has the disadvantage that it does not replicate the natural variability in morphology and dynamic behaviour between individuals within a species or between different species that make up a community. For example, plant morphology is variable in time (seasonality) and space and the structure of vegetation patches varies (e.g. in height, density, vitality, number of leaves per shoot) even within species depending on their locations (Puijalón *et al.* 2008a). The combination of this heterogeneity, both at the scale of the individual and at the scale of the community, along with the unsteadiness of the hydrodynamic regime makes analysing flow-organism interaction a very complex scientific problem. While this heterogeneity has been recognised and investigated in detail from a biological perspective (e.g. Neumeier 2005, Harder *et al.* 2006, Möller 2006, Stewart 2006, Puijalón *et al.* 2008a, Feagin *et al.* 2011, Möller *et al.* 2011, Miler *et al.* 2012), its effect on the physical environment has not yet been fully addressed. It is therefore uncertain whether heterogeneity within species needs to be taken into account or whether the use of a representative morphology is sufficient for studies of organism-flow-interactions.

As stated previously, while some species of mollusc may be present globally, those in areas of high hydraulic stress are likely to be morphologically dissimilar to those from sheltered areas and it would be useful to establish the extent to which these morphological adaptations benefit an individual mollusc and also influence the hydraulics within the surrounding fluid.

7. Scale and Scaling

7.1. Theme Description

This theme encompasses two key ideas: First, do biological (and biomechanical) processes scale in the same way as geomorphological and hydraulic processes? i.e., can biological processes operating at the large scale in the real world be scaled such that behaviours are replicated in physical experiments conducted at smaller scales? Second, are we measuring the correct variables at the most appropriate scales? To what extent are the variables we measure at the large scale the pertinent variables at a smaller scale? Do present technologies permit us to measure the same variable across scales? The answers to these questions are dependent upon which parameters can be controlled and measured in both the field and in the experimental set up. This presents us with difficulties because, for example, biocosms enable researchers to have good control of abiotic variables but poor control of hydraulic variables while experimental flume facilities enable there to be good control of hydraulic variables but not abiotic variables. Furthermore, velocity measurement techniques have finite sampling volume sizes and at smaller scales, the amount of spatial averaging incorporated within a velocity

measurement may be so large as to smear significant spatio-temporal patterns. For example, a modern laboratory Acoustic Doppler Velocimeter (ADV) has a typical sampling volume of 85 mm³. This corresponds to a cylinder of 6 mm in diameter and 3 mm in height (Nortek AS 2009) and is probably too coarse for measurements in turbulent flows with integral length scales of the same order as found near the vicinity of biofilms or in vegetal canopies with thin stems (see for instance the criticism of the interpretation of ADV measurements of López and García 2001 by Barkdoll 2002). Even hot-film anemometry is generally limited to spot measurements within 2 mm of the object of interest (e.g. Biggs *et al.* 1998).

7.2. Examples of knowledge gaps

It is often necessary to scale biota and hence employ surrogates because in some situations it is not feasible to use 1:1 scale physical models. Many studies have scaled physical processes using Froude or Reynolds numbers as the basis for comparison (e.g. ASCE 1942, Yalin 1971, Peakall *et al.* 1996, Julien 2002, Wilson *et al.* 2003). However, when researching at the interface with ecology, it is also necessary to scale biological factors. The studies of Gran and Paola (2001) and Tal and Paola (2007, 2010) scaled the stem diameter of live surrogate riparian trees (alfalfa), but it may also be necessary to scale other parameters such as the elasticity, root-mass and stem or population density (see Nikora 2010 for some suggestions of appropriate similarity numbers). Many biological parameters may not scale linearly and for others, the concept of scaling is difficult to envisage. For instance, when scaling population density of live surrogates is it necessary to scale competition and how can that be achieved? Does a direct relationship even exist between population density and competition (between either individual organisms or species), and/or is it possible to identify other reasonable metrics for competition? If not, it may be necessary to neglect some biological parameters, with implications for the reliability of experimental conclusions.

Eco-hydraulic processes and hydraulic-organism interactions operate over a wide range of spatio-temporal scales. While linkages between microscale turbulence and biological response (i.e. avoidance or tolerance through sheltering, reconfiguration and/or streamlining) can be expected to be present, the extent to which this is the case is still unknown. Thus, if our interest is in organism response, to what extent is a detailed description of turbulence needed? The answer to this question is a function of the ability of the organism to reconfigure (i.e., its stiffness or flexibility), the scale of interest and also the scientific approach of the investigator (e.g. reductionist or holistic). If our interest concerns inflexible organisms at the channel- or coastline-scale, then presumably the answer is that a detailed description of turbulence is not warranted and only a relative time-averaged value is needed (e.g. turbulent fluctuation normalised by the time-averaged velocity; Nikora *et al.* 2003). However, if our interest concerns very flexible structures at the patch scale, the converse is likely to be true and microscale turbulence may have a significant impact upon the behaviour of individual

organisms. The ideal spatio-temporal resolution of measurements is also related to the scale of interest and, indeed, to the approach of the investigator. Given presently available technologies, perhaps the key question is how to upscale detailed measurements of turbulence signals at a single place in space and over relatively short periods of time (of the order of minutes to hours to adequately characterise the full range of turbulent fluctuations; Soulsby 1980), first to describe the response of individual organisms and second to describe higher level responses of the ecosystem as a whole.

There has been a great deal of research on the impacts of (especially) sessile animals on the physical environment. However, much of this previous research has a biological focus at a scale of relevance to organisms. Research at intermediate- and large-scales is largely missing from the existing literature. This is of importance both because of the bioengineering potential of aggregations of bivalves for protecting coasts and for assessing the significance of sessile animals at larger scales. For instance, mussels have been found to significantly alter near-bed hydraulics in flumes with a maximum flow depth of 1 m (Butman *et al.* 1994, Van Duren *et al.* 2006, Peine *et al.* 2011), but the significance of mussel beds in natural flows that can be tens of metres deep is unknown. There is also significant uncertainty associated with upscaling bioturbation fluxes obtained from laboratory experiments on a small number of organisms to the broader ecosystem scale, and the significance of bioturbation (in terms of sediment, nutrient and other fluxes) relative to physical processes remains largely unknown. In one exception, Grant (1983) found that, during high ebb flow, rays disturbed up to $24.2 \text{ cm}^3 \text{ m}^{-2}$ of sediment, worms displaced $1.8 \text{ cm}^3 \text{ m}^{-2}$ and amphipods disturbed $60\text{-}70 \text{ cm}^3 \text{ m}^{-2}$. These are significant quantities, especially when accumulated; however they are minor compared to tidal sediment displacement which was measured to be between 6.1×10^3 and $1.2 \times 10^4 \text{ cm}^3 \text{ m}^{-2}$ during the high ebb flow. Similar studies need to be developed for other animal-related processes in order to assess their significance in comparison to acknowledged physical processes.

Most studies investigating the impact of vegetation on coastal hydrodynamics have been performed over short durations and in the summer months, when above-ground biomass is high and incident wave conditions are slight to moderate (e.g. Paul and Amos 2011, Jadhav and Chen 2012). Field studies on unidirectional and channelled flows have also tended to focus on low or moderate discharges, due to difficulties associated with planning field campaigns to measure highly unpredictable events (e.g. Bakry *et al.* 1992 and references therein). Results obtained from such studies cannot easily be extrapolated to the annual scale for perennial species (e.g. Widdows *et al.* 2008a, 2008b) and it is questionable whether results obtained in this manner can be upscaled to extreme or catastrophic events (Feagin *et al.* 2009; Koch *et al.* 2009). Attempts have been made to investigate the effect of vegetation on extreme events in laboratory studies, but these experiments have been performed in small-scale model basins, a fact which raises doubts as to how well they express field conditions due to distorted scaling and edge effects (Lynett 2007). It is therefore

desirable to undertake full-scale laboratory or field studies under extreme conditions to establish how present knowledge can be transferred to extreme events such as large magnitude, low frequency floods or storm surges (e.g., see Rueben *et al.* 2011 or Song and Irish in prep.).

8. Technological and infrastructural needs

In order to address the outstanding issues identified above, the participants in the Trondheim workshop noted a series of limitations of existing infrastructures and measurement technologies and made a number of suggestions for infrastructural and technological development. It was recognised that technological developments tend to be stepwise and that it is thus necessary to anticipate the refinement of existing techniques and technologies. However, it was also recognised that there must be an awareness of potentially new and/or ground-breaking technologies and techniques that may lead to a revolution in our understanding, rather than evolution.

8.1. Importance of interdisciplinary research

Perhaps the most obvious starting point for progress is the improvement of “human technology”. Crucially, to date, there has been a dearth of truly interdisciplinary studies (and interdisciplinary discussions) involving engineers, applied mathematicians, microbiologists, ecologists and fluid dynamicists. Disciplinary backgrounds lead scientists in disparate fields to ask fundamentally different questions (see Rice *et al.* 2010a and references therein). For example, microbiologists may focus on biofilm microbial or algal composition and succession, and relate it to biomass and function (nutrient degradation, carbon, nitrogen, and/or phosphorus cycles), but may provide very little quantitative information on either the properties of the flow or the mechanical properties (three-dimensional structure, elasticity, porosity) of the biofilm. Hydraulics and fluid mechanics researchers provide a better description of the local flow conditions but ignore those variables studied by microbiologists and struggle to describe the properties of the turbulent boundary layer and mass transfer due to the lack of information on the biomechanical properties of the underlying biofilms. The characterisation of these biomechanical properties will likely require the application of existing techniques developed in other fields or scientific communities (e.g., cryosections or Confocal Laser-Scanning Microscopy (CLSM) used by Battin *et al.* 2003 or 3D X-ray tomography used by Iltis *et al.* 2011) or the development of new techniques (e.g., the use of rotating electrodes by Boulêtreau *et al.* (2011) to make biofilm thickness and elasticity measurements). Further, these biomechanical data are required for input into numerical models that may help to develop a more complete understanding of scale-dependent interactions and feedbacks. High resolution small-scale models can then be used to parameterise large-scale models to bridge gaps between processes operating at different scales (see also section 7.2). However, numerical models are limited without adequate parameterisation and validation datasets. Many experimental programmes suggest such data as an anticipated outcome, but few consider the spatio-temporal resolution of the model mesh during experimental design. To

minimise the need for interpolation and extrapolation, measurements should be undertaken at a resolution and precision that is commensurate with that of the numerical model and at positions in space and instants in time that are coincident with the model mesh. Thus, interdisciplinary discussion and collaboration, within and between research institutes, consultancies and management agencies, are key first steps for identifying and understanding the essential behaviours of flora and fauna under hydraulic stress. The result of such discussion and collaboration should be the identification and development of common themes, problems and questions across disparate fields and applications.

8.2. Limitations of existing facilities

Before describing potential areas for improvement, it is perhaps beneficial to first outline some basic limitations of existing technologies and facilities. Some of these limitations follow directly from the lack of interdisciplinary collaboration during the design and construction of experimental facilities. For example, most studies conducted in fluid dynamics laboratories have been geared towards investigating responses to hydrodynamic forcing and have ignored behavioural changes and longer-term responses triggered by abiotic stresses (perhaps due to toxic effects or long term fitness effects). Therefore, many experimental facilities can only house species that are relatively insensitive to abiotic stimuli and do not permit researchers to control some key abiotic factors that are needed for healthy organisms and/or natural behaviours (e.g. temperature or oxygen content). Furthermore, most hydraulics facilities have little control over the source of water- many use domestic or commercial water supply networks that have been chemically treated to alter or reduce the bio-geochemical activity that is crucial for the natural function of ecological systems. Changing the bio-geochemical constituents of the water, whether ground- or surface-water, may also have implications when sediment is incorporated in experiments. Adding sediment to a flume is not trivial and ensuring that it is adequately recirculated, without artificially sorting it, through pumps is already challenging even before one considers the control of nutrient fluxes and biotically mediated interactions between the water and the sediment. These fluxes are sensitive to the chemical signatures of the surface water, pore-water and the constituent components of the sediment bed itself. Adequately studying these fluxes, behaviours and responses requires large, deep facilities where key parameters can be controlled for long periods of time. Such facilities are often prohibitively expensive to construct and run and so an additional consideration is another human factor: funding.

8.3. Desirable improvements to facilities

Addressing the limitations identified in section 8.2 will first require many of the issues and themes described in sections 3 to 7 to be tackled by the scientific community. For example, while there is undoubtedly a need for new facilities that are flexible and capable of employing ranges of widths and depths and with holding facilities, etc... that ensure organism integrity, what range of widths and depths is required? It is known that boundary and territorial conditions are important for animals and

plants, but how wide does a flume need to be to house a certain number of individuals and/or species? To answer this question, it will be necessary to improve direct observations of animals and plants in the natural environment, perhaps exploiting microsensors and nanotechnologies to develop location sensors, or using smaller tracers for plants and animals. Alternatively, perhaps it is possible to use chemical indicators to detect and measure the stress caused to organisms by different stimuli. Whichever methodologies are adopted, it will also be necessary to develop behavioural screening devices to identify and isolate organism responses to specific stimuli. In addition, the depth of the flume is a key limiting factor, both from the perspective of maximum water depth, but also maximum sediment thickness. Benthic organisms are likely to be sensitive to hyporheic flow, and thus a flume must be deep enough to ensure flow through a porous bed, not preferential flow along the flume floor.

While hydraulic flume facilities have excellent control over flow rates and velocity fields they generally have limited control over abiotic factors. Conversely, the biocosms employed in aquaculture research have greater control over abiotic factors, but have little or no control over flow/wave conditions. Biocosms are generally installed *in situ* as an enclosure within the field environment. Therefore, problems associated with modifications to the bio-geochemical properties of the ground- and surface-water are minimised. There are a handful of facilities across the world (e.g., the St. Anthony Falls Laboratory, Minneapolis, USA; the Silverstream flume near Christchurch, New Zealand; and the Stroud Water Research Center, Pennsylvania, USA) that extract natural stream water to ensure that bio-geochemical properties are unchanged, but more hydraulic flumes need to be built with this provision. It is thus desirable to transfer knowledge and technologies to produce hybrid facilities capable of ensuring control over both hydraulic and abiotic variables. This should make it possible to study the behaviour of sensitive species, such as sponges, that need better control of abiotic factors. In addition, almost all fish species and many aquatic invertebrates are sensitive to underwater sounds and vibrations. Although acoustic ecology is a well-established area of research, relatively little is known about the response of aquatic organisms to environmental noises. However, animals are known to respond to anthropogenic sounds and noise is regularly and successfully used to manipulate the movement of fish in rivers in relation to engineered features, such as turbines and fish passages (Knudsen *et al.* 1994, Popper and Hastings 2009). Therefore, it is likely that aquatic organisms will be sensitive to the vibration and noise produced by pumps and also ambient laboratory noise. Improvements are thus needed to reduce these disturbances, either through improvements to pump technologies and/or by sound-proofing flume walls.

However, in all likelihood, the single most important factor limiting advancement of eco-hydraulic experimentation is time. It is presently not feasible for researchers to adequately control key parameters, run experiments and take detailed measurements for prolonged periods of time. First, it is not straightforward to ensure continuous, reliable operation of complex equipment such as pumps,

lasers, high-speed cameras etc... Furthermore, measurement techniques such as Laser Doppler Anemometry (LDA) and PIV are incredibly memory intensive and have associated high data storage requirements (of the order of 80GB of data per minute with a modern 100Hz, 5MP camera) and it is difficult to prevent memory buffers from becoming full, causing data loss. After image acquisition and storage, PIV requires significant computing power to process images, cross-correlate between images and extract velocity fields. Thus, improvements also need to be made to computing facilities in order to extract and post-process the resulting data. Solutions to all these issues are necessarily expensive. This has implications for funding streams. In particular, the transnational access components of HYDRALAB IV, whereby the European Commission provides funds to large or unique affiliated laboratories to enable researchers from other countries to perform experiments in those installations may provide a model for new funding mechanisms. Within each member state, funding bodies may provide funds such that researchers from other institutions within the member state can gain access to facilities that are capable of performing experiments for prolonged periods of time.

9. Conclusion

This review has described the outcomes of discussions between contributors to the PISCES work package of HYDRALAB IV and a panel of invited experts to identify coherent themes undermining existing and future efforts to further understand eco-hydraulic interactions. Five dominant issues were defined and discussed: abiotic factors; adaptation and behaviour; complexity and feedback; heterogeneity; and scale and scaling. Examples of areas where knowledge is currently lacking, and of avenues for future research, have been presented, with reference to biofilms, plants, sessile and mobile fauna in shallow water fluvial and marine environments. Finally, limitations in existing practices and experimental facilities have been identified and suggestions on future improvements outlined.

It is our contention that physical modelling, informed by interdisciplinary discussion and collaboration, will play a crucial role in advancing our knowledge of organism-fluid interactions. The exploitation of infrastructural and technological advances will be central to this advance. Results obtained from studies performed over short durations and under low to moderate hydraulic forcing (e.g. incident wave conditions or discharges) cannot easily be extrapolated to the seasonal nor annual scale, nor up-scaled adequately to extreme or catastrophic events. There is therefore a desperate need for large, deep, and flexible hydraulic facilities that can permit the study of biogeochemical, biological, ecological and sedimentary interactions under extreme conditions (e.g., floods or storm surges) for extended periods. Furthermore, knowledge and technology will need to be transferred between disparate fields to design and construct hybrid facilities capable of ensuring control and measurement of both hydraulic and abiotic variables. For maximum flexibility, such a facility should

be capable of sourcing natural stream and/or unfiltered water to ensure that bio-geochemical properties are unchanged and simultaneously also be capable of using salt water. Because of construction, maintenance and staffing costs, it is likely that this would inevitably be a shared facility and would require the implementation of imaginative funding mechanisms, such as the transnational access components of HYDRALAB IV that enable researchers from European countries to perform experiments in large or unique affiliated laboratories.

10. Acknowledgements

The work described in this publication was supported by the European Community's 7th Framework Programme through a grant to the budget of the Integrated Infrastructure Initiative HYDRALAB IV, Contract no. 261520. This document reflects only the authors' views and not those of the European Community. This work may rely on data from sources external to the HYDRALAB project Consortium. Members of the Consortium do not accept liability for loss or damage suffered by any third party as a result of errors or inaccuracies in such data. The information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose. The user thereof uses the information at its sole risk and neither the European Community nor any member of the HYDRALAB Consortium is liable for any use that may be made of the information. The authors wish to thank Drs. Walter Bertoldi, Chris Bromley, Andrew Folkard, Marwan Hassan, Anne Lightbody, Ponnambalam Rameshwaran, Jonas Schoelynck, Desirée Tullos and Catherine Wilson, who all read an earlier draft of this manuscript and contributed useful comments and discussions.

11. References

- Albagnac, J., Moulin, F.Y., Eiff, O., Lacaze, L., Brancher, P. (2013). A three-dimensional experimental investigation of the structure of the spanwise vortex generated by a shallow vortex dipole. *Environ. Fluid Mech.* DOI: 10.1007/s10652-013-9291-6.
- Aller, R.C., Yingst, J.Y., Ullman, W.J. (1983). Comparative biogeochemistry of water in intertidal *Onuphis* (Polychaeta) and *Upogebia* (Crustacea) burrows: temporal patterns and causes. *J. Marine Res.* **41**, 571–604.
- Amyot, J.-P., Downing, J.A. (1997). Seasonal variation in vertical and horizontal movement of the freshwater bivalve *Elliptio complanata* (Mollusca: Unionidae). *Freshwater Biol.* **37**, 345–354.
- Andersen, T.J. (2001). Seasonal variation in erodability of two temperate, microtidal mudflats. *Estuar. Coast. Shelf Sci.* **53**, 1–12.

- Arroyo, M.P., Hinsch, K.D. (2008). Recent developments of PIV towards 3D measurements. In *Particle Image Velocimetry: New developments and recent applications. Topics in Applied Physics* **112**. A. Schroeder, C.E. Willert, eds. Springer, Berlin/Heidelberg, Germany, 127–154.
- ASCE. (1942). *Hydraulic Models, American Society of Civil Engineers Manual of Practice* **25**. American Society of Civil Engineers, New York, NY, USA.
- Astall, C.M., Taylor, A.C., Atkinson, R.J.A. (1997). Behavioural and physiological implications of a burrow-dwelling lifestyle for two species of *Upogebiid* mud-shrimp (Crustacea: Thalassinidea). *Estuar. Coast. Shelf Sci.* **44**, 155–168.
- Bakry, M.F., Gates, T.K., Khattab, A.F. (1992). Field-measured hydraulic resistance characteristics in vegetation-infested canals. *J. Irrig. Drain. Eng.* **118**(2), 256–274.
- Barkdoll, B.D. (2002). Discussion of “Mean flow and turbulence structure of open-channel flow through non-emergent vegetation” by Fabián López and Marcelo H. García. *J. Hydraulic Eng.* **128**(11), 1032.
- Barmuta, L.A., McKenny, C.E.A., Swain, R. (2001). The response of lotic mayfly *Nousia* sp. (Ephemeroptera: Leptophlebiidae) to moving water and light of different wavelengths. *Freshwater Biol.* **46**, 567–573.
- Battin, T.J., Kaplan, L.A., Newbold, J.D., Cheng, X., Hansen, C. (2003). Effects of current velocity on the nascent architecture of stream microbial biofilms. *Appl. Environ. Microb.* **69**, 5443–5452.
- Beven, K.J. (1989). Changing ideas in hydrology: the case of physically-based models. *J. Hydrology* **105**(1-2), 157–172.
- Biggs, B.J.F., Goring, D.G., Nikora, V.I. (1998). Subsidy and stress responses of stream periphyton to gradients in water velocity as a function of community growth form. *J. Phycol.* **34**, 598–607.
- Bornette, G., Puijalon, S. (2011). Response of aquatic plants to abiotic factors: a review. *Aquat. Sci.* **73**, 1–14.
- Boulêtreau, S., Charcosset, J.-Y., Gamby, J., Lyautey, E., Mastrotillo, S., Azémar, F., Moulin, F., Tribollet, B., Garabétian, F. (2011). Rotating disk electrodes to assess river biofilm thickness and elasticity. *Water Res.* **45**, 1347–1357.
- Bouma, T.J., de Vries, M.B., Low, E., Peralta, G., Táncoz, I.C., van de Koppel, J., Herman, P.M.J. (2005). Trade-offs related to ecosystem engineering: A case study on stiffness of emerging macrophytes. *Ecology* **86**, 2187–2199.

- Bragazza, L., Gerdol, R. (2002). Are nutrient availability and acidity-alkalinity gradients related in Sphagnum-dominated peatlands? *J. Veg. Sci.* **13**(4), 473–482.
- Branch, G.M., Marsh, A.C. (1978). Tenacity and shell shape in six *Patella* species: adaptive features. *J. Exp. Mar. Biol. Ecol.* **34**, 111–130.
- Butman, C.A., Fréchet, M., Geyer, W.R., Starczak, V.R. (1994). Flume experiments on food supply to the blue mussel *Mytilus edulis* L. as a function of boundary-layer flow. *Limnol. Oceanogr.* **39**, 1755–1768.
- Cameron, S.M., Nikora, V.I., Albayrak, I., Miler, O., Stewart, M., Siniscalchi, F. (in review). Interactions between aquatic plants and turbulent flow: A field study using stereoscopic PIV. *J. Fluid Mech.*
- Chorley, R.J. (1962). Geomorphology and general systems theory. *US Geological Survey Professional Paper 500-B*. US Government Printing Office, Washington, D.C., USA.
- Coops, H., Geilen, N., Verheij, H.J., Boeters, R., van der Velde, G. (1996). Interactions between waves, bank erosion and emergent vegetation: an experimental study in a wave tank. *Aquat. Bot.* **53**, 187–198.
- D’Andrea, A.F., DeWitt, T.H. (2009). Geochemical ecosystem engineering by the mud shrimp *Upogebia pugettensis* (Crustacea: Thalassinidae) in Yaquina Bay, Oregon: density-dependent effects on organic matter remineralization and nutrient cycling. *Limnol. Oceanogr.* **54**, 1911–1932.
- Darby, S.E. (2010). Reappraising the geomorphology–ecology link. *Earth Surf. Process. Landforms* **35**(3), 368–371.
- de Brouwer, J.F.C., Bjelic, S., De Deckere, E.M.G.T., Stal, L.J. (2000). Interplay between biology and sedimentology in a mudflat (Biezelingse Ham, Westerschelde, The Netherlands). *Cont. Shelf Res.* **20**, 1159–1177.
- Doyle, R.D. (2001). Effects of waves on the early growth of *Vallisneria americana*. *Freshwater Biol.* **46**, 389–397.
- Dworschak, P.C. (1987). Burrows of *Solecurtus strigilatus* (Linne) and *S. multistriatus* (Scacchi). *Senck. Marit.* **19**, 131–147.
- Favis-Mortock, D.T., Quinton, J.N., Dickinson, W.T. (1996). The GTCE validation of soil erosion models for global change studies. *J. Soil Water Conserv.* **51**(5), 397–403.

- Feagin, R.A., Lozada-Bernard, M., Ravens, T.M., Möller, I., Yeager, K.M., Baird, A.H. (2009). Does vegetation prevent wave erosion of salt marsh edges? *P. Nat. Acad. Sci. USA* **106**, 10109–10113.
- Feagin, R.A., Irish, J.L., Möller, I., Williams, A.M., Colón-Rivera, R.J., Mousavi, M.E. (2011). Short communication: Engineering properties of wetland plants with application to wave attenuation. *Coastal Eng.* **58**(3), 251–255.
- Fisher, S.G., Heffernan, J.B., Sponseller, R.A., Welter, J.R. (2007). Functional ecomorphology: Feedbacks between form and function in fluvial landscape ecosystems. *Geomorphology* **89**(1–2), 84–96.
- Folkard, A.M., Gascoigne, J.C. (2009). Hydrodynamics of discontinuous mussel beds: Laboratory flume simulations. *J. Sea Res.* **62**(4), 250–257.
- Fonseca, M.S., Fisher, J.S. (1986). A comparison of canopy friction and sediment movement between four species of seagrass with reference to their ecology and restoration. *Mar. Ecol.-Prog. Ser.* **29**, 15–22.
- Fortino, K. (2006). Effect of season on the impact of ecosystem engineers in the New River, NC. *Hydrobiologia* **559**, 463–466.
- Friedrichs, M., Graf, G., Springer, B. (2000). Skimming flow induced over a simulated polychaete tube lawn at low population densities. *Mar. Ecol.-Prog. Ser.* **192**, 219–228.
- García-March, J.R., Pérez-Rojas, L., García-Carrascosa, A.M. (2007). Influence of hydrodynamic forces on population structure of *Pinna nobilis* L., 1758 (Mollusca: Bivalvia): the critical combination of drag force, water depth, shell size and orientation. *J. Exp. Mar. Biol. Ecol.* **342**, 202–212.
- Gobert, S., Kyramarios, M., Lepoint, G., Pergent-Martini, C., Bouquegneau, J.-M. (2003). Variations at different spatial scales of *Posidonia oceanica* (L.) Delilebeds; effects on the physico-chemical parameters of the sediment. *Oceanol. Acta* **26**, 199–207.
- Gran, K., Paola, C. (2001). Riparian vegetation controls on braided stream dynamics. *Water Resour. Res.* **37**, 3275–3283.
- Grant, J. (1983). The relative magnitude of biological and physical sediment reworking in an intertidal community. *J. Marine Res.* **41**, 673–689.
- Griffen, B.D., deWitt, T.H., Langdon, C. (2004). Particle removal rates by the mud shrimp *Upogebia pugettensis*, its burrow, and a commensal clam: effects on estuarine phytoplankton abundance. *Mar. Ecol.-Prog. Ser.* **269**, 223–236.

- Hammond, W., Griffiths, C.L. (2004). Influence of wave exposure on South African mussel beds and their associated infaunal communities. *Mar. Biol.* **144**, 547–552.
- Harder, D., Hurd, C., Speck, T. (2006). Comparison of mechanical properties of four large, wave-exposed seaweeds. *Am. J. Bot.* **93**(10), 1426–1432.
- Hedden, R.L., Fredericksen, T.S., Williams, S.A. (1995). Modeling the effect of crown shedding and streamlining on the survival of loblolly pine exposed to acute wind. *Can. J. Forest Res.* **25**, 704–712.
- Hession, W.C., Curran, J.C., Resler, L.M., Wynn, T.M. (2010). Preface: geomorphology and vegetation: interactions, dependencies, and feedback loops. *Geomorphology* **116**(3–4), 203–205.
- Hurtado, M.Á., Mirella da Silva, P., Le Goïc, N., Palacios, E., Soudant, P. (2011). Effect of acclimatization on hemocyte functional characteristics of the Pacific oyster (*Crassostrea gigas*) and carpet shell clam (*Ruditapes decussatus*). *Fish Shellfish Immun.* **31**(6), 978–984.
- Idestam-Almquist, J., Kautsky, L. (1995). Plastic responses in morphology of *Potamogeton pectinatus* L. to sediment and above sediment conditions at two sites in the northern Baltic proper. *Aquat. Bot.* **52**, 205–216.
- Iltis, G.C., Armstrong, R.T., Jansik, D.P., Wood, B.D., Wildenschild, D. (2011). Imaging biofilm architecture within porous media using synchrotron-based X-ray computed microtomography. *Water Resour. Res.* **47**, W02601.
- Jadhav, R., Chen, Q. (2012). Field investigation of wave dissipation over salt marsh vegetation during tropical cyclone. Proc. 33rd Int. Conf. *Coastal Engineering* Santander Spain, waves.41, P. Lynett, J. McKee Smith, eds. ICCE, Los Angeles, CA, USA.
- James, W.F., Barko, J.W. (2000). Sediment resuspension dynamics in canopy- and meadow-forming submersed macrophyte communities. *U.S. Army Corps of Engineers, Engineer Research and Development Center Report ERDC/EL SR-00-8*. U.S. Army Corps of Engineers, Vicksburg, MS, USA.
- Jones, C.G., Lawton, J.H., Shachak, M. (1994). Organisms as ecosystem engineers. *Oikos* **69**, 373–86.
- Jones, S.E., Jago, C.F. (1993). *In situ* assessment of modification of sediment properties by burrowing invertebrates. *Mar. Biol.* **115**, 133–142.
- Jørgensen, C.B. (1976). Growth efficiencies and factors controlling size in some mytilid bivalves, especially *Mytilus edulis* L.: review and interpretation. *Ophelia* **15**, 175–192.

- Julien, P.Y. (2002). *River Mechanics*. Cambridge University Press, Cambridge, UK.
- Kawai, T., Tokeshi, M. (2004). Variable modes of facilitation in the upper intertidal: goose barnacles and mussels. *Mar. Ecol.-Prog. Ser.* **272**, 203–213.
- Kelly, N.M., Fonseca, M., Whitfield, P. (2001). Predictive mapping for management and conservation of seagrass beds in North Carolina. *Aquat. Conserv.* **11**, 437–451.
- King, C.A.M. (1970). Feedback relationships in geomorphology. *Geogr. Ann. A* **52**(3-4), 147–159.
- Knudsen, F.R., Enger, P.S., Sand, O. (1994). Avoidance responses to low frequency sound in downstream migrating Atlantic salmon smolt, *Salmo salar*. *J. Fish Biol.* **45**(2), 227–233.
- Koch, E.W., Barbier, E.B., Silliman, B.R., Reed, D.J., Perillo, G.M.E., Hacker, S.D., Granek, E.F., Primavera, J.H., Muthiga, N., Polansky, S., Halpern, B.S., Kennedy, C.J., Kappel, C.V., Wolanski, E. (2009). Non-linearity in ecosystem services: temporal and spatial variability in coastal protection. *Front. Ecol. Environ.* **7**, 29–37.
- Lauder, G.V. (2011). Swimming hydrodynamics: ten questions and the technical approaches needed to resolve them. *Exp. Fluids* **51**, 23–35.
- Lepoint, G., Vangeluwe, D., Eisinger, M., Paster, M., van Treeck, P., Bouquegneau, J.-M., Gobert, S. (2004). Nitrogen dynamics in *Posidonia oceanica* cuttings: implications for transplantation experiments. *Mar. Pollut. Bull.* **48**, 465–470.
- López, F., García, M.H. (2001). Mean flow and turbulence structure of open-channel flow through non-emergent vegetation. *J. Hydraulic Eng.* **127**(5), 392–402.
- Lynett, P. (2007). Effect of a shallow water obstruction on long wave runup and overland flow velocity. *J. Waterway, Port, Coastal, Ocean Eng.* **133**(6), 455–462.
- Maltese, A., Cox, E., Folkard, A.M., Ciraolo, G., La Loggia, G., and Lombardo, G. (2007). Laboratory measurements of flow and turbulence in discontinuous distributions of ligulate seagrass. *J. Hydraul. Eng.* **133**(7), 750–760.
- McQuaid, C.D., Lindsay, T.L. (2000). Effect of wave exposure on growth and mortality rates of the mussel *Perna perna*: Bottom-up regulation of intertidal population. *Mar. Ecol.-Prog. Ser.* **206**, 147–154.
- McQuaid, C.D., Lindsay, J.R., Lindsay, T.L. (2000). Interactive effects of wave exposure and tidal height on population structure of the mussel *Perna perna*. *Mar. Biol.* **137**, 925–932.

Merz, R.A. (1984). Self-generated versus environmentally produced feeding currents: a comparison for the sabellid polychaete *Eudistylia vancouveri*. *Biol. Bull.* **167**, 200–209.

Miler, O., Albayrak, I., Nikora, V., O'Hare, M. (2012). Biomechanical properties of aquatic plants and their effects on plant–flow interactions in streams and rivers. *Aquat. Sci.* **74**(1), 31–44.

Möller, I. (2006). Quantifying saltmarsh vegetation and its effect on wave height dissipation: Results from a UK East coast saltmarsh. *Estuar. Coast. Shelf Sci.* **69**, 337–351.

Möller, I., Mantilla-Contreras, J., Spencer, T., Hayes, A. (2011). Micro-tidal coastal reed beds: Hydro-morphological insights and observations on wave transformation from the southern Baltic Sea. *Estuar. Coast. Shelf Sci.* **92**, 424–436.

National Research Council Committee for the Update of the Guide for the Care and Use of Laboratory Animals. (2011). *Guide for the Care and Use of Laboratory Animals, Eighth Edition*. The National Academies Press, Washington, D.C., USA.

Nepf, H.M. (2012). Flow and transport in regions with aquatic vegetation. *Annu. Rev. Fluid Mech.* **44**, 123–142.

Neumeier, U. (2005). Quantification of vertical density variations of salt-marsh vegetation. *Estuar. Coast. Shelf Sci.* **63**, 489–496.

Nikora, V. (2010). Hydrodynamics of aquatic ecosystems: an interface between ecology, biomechanics and environmental fluid mechanics. *River. Res. Applic.* **26**, 367–384.

Nikora, V.I., Aberle, J., Biggs, B.J.F., Jowett, I.G., Sykes, J.R.E. (2003). Effects of fish size, time-to-fatigue and turbulence on swimming performance: a case study of *Galazias maculatus*. *J. Fish Biol.* **63**, 1365–1382.

Nortek AS. (2009). *Vectrino velocimeter manual*. Nortek AS, Oslo, Norway.

Parry, M.L., Canziani, O.F., Palutikof, J.P., van der Linden, P.J., Hanson, C.E. (eds). (2007). *Climate Change 2007: Impacts, Adaptation and Vulnerability*. Cambridge University Press, Cambridge, UK.

Paul, M., Amos, C.L. (2011). Spatial and seasonal variation in wave attenuation over *Zostera noltii*. *J. Geophys. Res. - Oceans* **116**(C8), C08019.

Peakall, J., Ashworth, P., Best, J. (1996). Physical modelling in fluvial geomorphology: Principles, applications and unresolved issues. In *The Scientific Nature of Geomorphology*. B.L. Rhoads, C.E. Thorn, eds. John Wiley & Sons, Chichester, UK, pp. 221–253.

- Peakall, J., Ashworth, P.J., Best, J.L. (2007). Meander-bend evolution, alluvial architecture, and the role of cohesion in sinuous river channels: A flume study. *J. Sediment. Res.* **77**, 197–212.
- Peine, F., Friedrichs, M., Graf, G. (2009). Potential influence of tubicolous worms on the bottom roughness length z_0 in the south-western Baltic Sea. *J. Exp. Mar. Biol. Ecol.* **374**, 1–11.
- Peine, F., Bobertz, B., Graf, G. (2011). Influence of the blue mussel *Mytilus edulis* (Linnaeus) on the bottom roughness length (z_0) in the south-western Baltic Sea. *Baltica* **18**, 13–22.
- Pollen, N., Simon, A. (2006). Geotechnical implications for the use of Alfalfa in experimental studies of alluvial-channel morphology and planform. *Eos Trans. AGU* **87**(52), Fall Meet. Suppl., Abstract H21G-1455.
- Popper, A.N., Hastings, M.C. (2009). The effects of anthropogenic sources of sound on fishes. *J. Fish Biol.* **75**(3), 455–489.
- Prandtl, L. (1952). *Essentials of fluid dynamics: with applications to hydraulics, aeronautics, meteorology and other subjects*. Blackie, London, UK.
- Puijalon, S., Bornette, G. (2004). Morphological variation of two taxonomically distant plant species along a natural flow velocity gradient. *New Phytol.* **163**, 651–660.
- Puijalon, S., Bornette, G. (2006). Phenotypic plasticity and mechanical stress: biomass partitioning and clonal growth of an aquatic plant species. *Am. J. Bot.* **93**, 1090–1099.
- Puijalon, S., Sagnes, P., Bornette, G. (2005). Adaptations to increasing hydraulic stress: morphology, hydrodynamics and fitness of two higher aquatic plant species. *J. Exp. Bot.* **56**, 777–786
- Puijalon, S., Léna, J.-P., Rivière, N., Champagne, J.Y., Rostan, J.C., Bornette, G. (2008a). Phenotypic plasticity in response to mechanical stress: hydrodynamic performance and fitness of 4 aquatic plant species. *New Phytol.* **177**, 907–917.
- Puijalon, S., Bouma, T.J., van Groenendael, J., Bornette, G. (2008b). Clonal plasticity of aquatic plant species submitted to mechanical stress: escape vs. resistance strategy. *Ann. Bot.-London* **102**, 989–996.
- Reinhardt, L., Jerolmack, D., Cardinale, B.J., Vanacker, V., Wright, J. (2010). Dynamic interactions of life and its landscape: feedbacks at the interface of geomorphology and ecology. *Earth Surf. Process. Landforms* **35**(1), 78–101.

- Rice, S.P., Lancaster, J., Kemp, P. (2010a). Experimentation at the interface of fluvial geomorphology, stream ecology and hydraulic engineering and the development of an effective, interdisciplinary river science. *Earth Surf. Process. Landforms* **35**(1), 64–77.
- Rice, S.P., Little, S., Wood, P.J., Moir, H.J., Vericat, D. (2010b). The relative contributions of ecology and hydraulics to ecohydraulics. *River Res. Applic.* **26**(4), 363–366.
- Rice, S.P., Johnson, M.F., Reid, I. (2012). Animals and the geomorphology of gravel-bed rivers. In *Gravel Bed Rivers: Processes, Tools, Environments*. M.A. Church, P. Biron, A.G. Roy, eds. John Wiley and Sons, New York, USA, 225–241.
- Rowden, A.A., Jago, C.F., Jones, S.E. (1998). Influence of benthic macrofauna on the geotechnical and geophysical properties of surficial sediment, North Sea. *Cont. Shelf Res.* **18**, 1347–1363.
- Rueben, M., Holman, R., Cox, D., Shin, S., Killian, J., Stanley, J. (2011). Optical measurements of tsunami inundation through an urban waterfront modeled in a large-scale laboratory basin. *Coastal Eng.* **58**(3), 229–238.
- Sakakibara, J., Nakagawa, M., Yoshida, M. (2004). Stereo-PIV study of flow around a maneuvering fish. *Exp. Fluids* **36**, 282–293.
- Santamaria, L. (2002). Why are most aquatic plants widely distributed? Dispersal, clonal growth and small-scale heterogeneity in a stressful environment. *Acta Oecol.* **23**, 137–154.
- Schaller, J., Brackhage, C., Dudel, E.G. (2012a). Silicon availability changes structural carbon ratio and phenol content of grasses. *Environ. Exp. Bot.* **77**, 283–287.
- Schaller, J., Brackhage, C., Gessner, M.O., Bäüker, E., Dudel, E.G. (2012b). Silicon supply modifies C:N:P stoichiometry and growth of *Phragmites australis*. *Plant Biology* **14**(2), 392–396.
- Schoelynck, J. (2011). Macrophyte patches as biogeochemical hotspots: what is the impact on river water quality? *PhD Thesis*, Departement Biologie, Universiteit Antwerpen, Belgium.
- Schoelynck, J., De Groote, T., Bal, K., Vandenbruwaene, W., Meire, P., Temmerman, S. (2012). Self-organised patchiness and scale-dependent bio-geomorphic feedbacks in aquatic river vegetation. *Ecography* **35**(8), 760–768.
- Schumm, S.A. (1991). *To Interpret the Earth: Ten Ways to be Wrong*. Cambridge University Press, New York, USA.

- Seed, R. (1969). The ecology of *Mytilus edulis* L. (Lamellibranchiata) on exposed rocky shores. 2. Growth and mortality. *Oecologia* **3**, 317–350.
- Simeone, S., De Falco, G. (2012). Morphology and composition of beach-cast *Posidonia oceanica* litter on beaches with different exposures. *Geomorphology* **151–152**, 224–233.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (eds). (2007). *Climate Change 2007: The Physical Science Basis*. Cambridge University Press, Cambridge, UK.
- Song, Y.K., Irish, J.L. (In prep). Inundation by long waves in the presence of discontinuous macro-roughness with application to tsunami runup in forests. *J. Geophys. Res. - Oceans*.
- Soulsby, R.L. (1980). Selecting record length and digitization rate for near-bed turbulence measurements. *J. Phys. Oceanogr.* **10**, 208–219.
- Steffani, C.N., Branch, G.M. (2003). Growth rate, condition, and shell shape of *Mytilus galloprovincialis*: responses to wave exposure. *Mar. Ecol.-Prog. Ser.* **246**, 197–209.
- Stewart, H.L. (2006). Hydrodynamic consequences of flexural stiffness and buoyancy for seaweeds: a study using physical models. *J. Exp. Biol.* **209**(11), 2170–2181.
- Strand, J.A., Weisner, E.B. (1996). Wave exposure related growth of epiphyton: implications for the distribution of submerged macrophytes in eutrophic lakes. *Hydrobiologia* **325**, 113–119.
- Suchanek, T.H. (1983). Control of seagrass communities and sediment distribution by *Callinassa* (Crustacea, Thalassinidea) bioturbation. *J. Marine Res.* **41**, 281–298.
- Tal, M., Paola, C. (2007). Dynamic single-thread channels maintained by the interaction of flow and vegetation. *Geology* **35**, 347–350.
- Tal, M., Paola, C. (2010). Effects of vegetation on channel morphodynamics: results and insights from laboratory experiments. *Earth Surf. Process. Landforms* **35**(9), 1014–1028.
- Thorne, C.R., Evans, E.P., Penning-Rowsell, E.C. (eds). (2007). *Future flooding and coastal erosion risks*. Thomas Telford, London, UK.
- Tiselius, P., Hansen, B., Jonsson, P., Kiorboe, T., Nielsen, G., Piontkovski, S., Saiz, E. (1995). Can we use laboratory-reared copepods for experiments? A comparison of feeding behaviour and reproduction between a field and a laboratory population of *Acartia tonsa*. *ICES J. Mar. Sci.* **52**, 369–376.

van der Heide, T., van Nes, E.H., Geerling, G.W., Smolders, A.J.P., Bouma, T.J., van Katwijk, M.M. (2007). Positive feedbacks in seagrass ecosystems: Implications for success in conservation and restoration. *Ecosystems* **10**, 1311–1322.

van Duren, L.A., Herman, P.M.J., Sandee, A.J.J., Heip, C.H.R. (2006). Effects of mussel filtering activity on boundary layer structure. *J. Sea Res.* **55**, 3–14.

Vogel, S. (1994). *Life in Moving Fluid*. 2nd ed. Princeton University Press, Princeton, New Jersey, USA.

Wheaton, J.M., Gibbins, C., Wainwright, J., Larsen, L., McElroy, B. (2011). Preface: Multiscale Feedbacks in Ecogeomorphology. *Geomorphology* **126**(3–4), 265–268.

Widdows, J., Brinsley, M.D. (2002). Impact of biotic and abiotic processes on sediment dynamics and the consequences to the structure and functioning of the intertidal zone. *J. Sea Res.* **48**, 143–156.

Widdows, J., Brinsley, M.D., Bowley, N., Barrett, C. (1998). A benthic annular flume for *in situ* measurement of suspension feeding/biodeposition rates and erosion potential of intertidal cohesive sediments. *Estuar. Coast. Shelf Sci.* **46**, 27–38.

Widdows, J., Brinsley, M.D., Salkeld, P.N., Lucas, C.H. (2000a). Influence of biota on spatial and temporal variation in sediment erodability and material flux on a tidal flat (Westerschelde, The Netherlands). *Mar. Ecol.-Prog. Ser.* **194**, 23–37.

Widdows, J., Brown, S., Brinsley, M.D., Salkeld, P.N., Elliott, M. (2000b). Temporal changes in intertidal sediment erodability: influence of biological and climatic factors. *Cont. Shelf Res.* **20**, 1275–1289.

Widdows, J., Blauw, A., Heip, C.H.R., Herman, P.M.J., Lucas, C.H., Middelburg, J.J., Schmidt, S., Brinsley, M.D., Twisk, F., Verbeek, H. (2004). Role of physical and biological processes in sediment dynamics of a tidal flat in Westerschelde Estuary, SW Netherlands. *Mar. Ecol.-Prog. Ser.* **274**, 41–56

Widdows, J., Pope, N.D., Brinsley, M.D., Asmus, H., Asmus, R. (2008a). Effect of seagrass beds (*Zostera noltii* and *Z. marina*) on near-bed hydrodynamics and sediment resuspension. *Mar. Ecol.-Prog. Ser.* **358**, 125–136.

Widdows, J., Pope, N., Brinsley, M. (2008b). Effect of *Spartina anglica* stems on nearbed hydrodynamics, sediment erodability and morphological changes on an intertidal mudflat. *Mar. Ecol.-Prog. Ser.* **362**, 45–57.

Widdows, J., Pope, N.D., Brinsley, M.D., Gascoigne, J., Kaiser, M.J. (2009). Influence of self-organised structures on near-bed hydrodynamics and sediment dynamics within a mussel (*Mytilus edulis*) bed in the Menai Strait. *J. Exp. Mar. Biol. Ecol.* **379**, 92–100.

Wilson, C.A.M.E., Stoesser, T., Bates, P.D., Batemann Pinzen, A. (2003). Open channel flow through different forms of submerged flexible vegetation. *J. Hydraulic Eng.* **29**(11), 847–853.

Wright, L.D., Schaffner, L.C., Maa, J.P.-Y. (1997). Biological mediation of bottom boundary layer processes and sediment suspension in the lower Chesapeake Bay. *Mar. Geol.* **141**(1–4), 27–50.

Yager, P.L., Nowell, A.R.M., Jumars, P.A. (1993). Enhanced deposition to pits: a local food source for benthos. *J. Marine Res.* **51**, 209–236.

Yalin, M.S. (1971). *Theory of Hydraulic Models*. Macmillan, London, UK.

Ziebis, W., Forster, S., Huettel, M., Jorgensen, B.B. (1996). Complex burrows of the mud shrimp *Callinassa truncata* and their geochemical impact in the sea bed. *Nature* **382**, 619–622.