



DR. ANDREW JOHN SIMPSON (Orcid ID : 0000-0002-0480-1134)

Article type : Letter to the Editor

Exercise-induced bronchoconstriction: a survey of diagnostic practice in secondary care across the United Kingdom

Simpson AJ¹, Drake SM², Fowler SJ².

1. Department of Sport, Health and Exercise Science, University of Hull
2. Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The University of Manchester; Manchester Academic Health Science Centre and NIHR Manchester Biomedical Research Centre, Manchester University Hospitals NHS Foundation Trust, Manchester, United Kingdom

Corresponding author:

Dr Andrew Simpson, A.Simpson2@hull.ac.uk

Department of Sport, Health and Exercise Science

University of Hull

Cottingham Road

Hull

HU6 7RX

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14305](https://doi.org/10.1111/ALL.14305)

This article is protected by copyright. All rights reserved

To the Editor,

Exercise-induced bronchoconstriction (EIB) is highly prevalent in children and adults with asthma, and elite athletes. There is strong evidence to support that bronchial provocation tests that act indirectly are essential to obtain an accurate diagnosis of the condition.¹ Despite these recommendations, people with respiratory symptoms on exertion are regularly diagnosed with EIB based on symptoms alone and treated in primary care.² In secondary care, debate remains regarding the optimal diagnostic protocol.³ Further, it is unclear which diagnostic tests are commonly utilized and, importantly, the extent to which practitioners adhere to bronchial provocation guidelines in secondary care.

In this survey of diagnostic practice in secondary care, we aim to identify current indirect bronchial provocation challenges in use across the United Kingdom for the diagnosis of EIB and, where applicable, assess adherence to exercise bronchial provocation guidelines.

Methods

We conducted a survey of EIB diagnostic practices in hospitals with respiratory physiology laboratories, identified through the Association for Respiratory Technology and Physiology (ARTP). Forty-three hospitals were identified and approached via e-mail and/or phone. Details of the first-line indirect bronchial provocation test and standard operating procedures (SOP) for exercise challenge tests were requested. SOPs were subsequently examined for adherence to five criteria specified in the American Thoracic Society exercise bronchial provocation guidelines,^{1,4} as detailed in Table 1. Data regarding EIB diagnosis thresholds for exercise provocation testing were also obtained.

Results

Forty of 43 centers responded; 29 (73%) of these stated that they offered one or more indirect bronchial provocation tests for the diagnosis of EIB. As a first line diagnostic test, 48% of respondents conducted mannitol challenges, 45% exercise provocation challenges, 3% cardiopulmonary exercise testing and 3% histamine challenge testing.

Exercise bronchial provocation SOPs were provided by nine centers (21%) and their adherence to ATS guidelines assessed. All nine SOPs referred to measuring exercise intensity via heart rate at an appropriate intensity and all but one (89%) followed the guidance on exercise duration (figure 1). Spirometry timings did not follow ATS criteria in 67% of SOPs (figure 1). Importantly, only two of the SOPs (22%) included guidance on appropriate environmental conditions (figure 1). No center provided details sufficient

to indicate full adherence to medication restrictions were routinely implemented in the SOPs; the authors were therefore unable ascertain with any certainty whether medication restrictions were considered. The criterion for EIB diagnosis varied between centers; 56% specified a fall in FEV₁ of ≥10%, with the remaining SOPs stating a ≥15% fall in FEV₁ as the threshold for diagnosis following exercise provocation.

Discussion

Due to the poor sensitivity and specificity of respiratory symptoms to diagnose EIB,⁵an indirect bronchial provocation challenge is recommended.¹Strong evidence suggests that the prevalence of EIB will vary with the type of challenge and the conditions under which the challenge is performed.^{1,3}Thus, strict adherence to provocation guidelines is essential if results are to be valid and comparable across centers. Here, we report that mannitol and bronchial provocation with exercise are utilized in the vast majority of UK hospitals for the diagnosis of EIB; both of which are considered appropriate for the diagnosis of the condition.¹Interestingly, eucapnic voluntary hyperpnoea, which is often referred to as the 'gold standard' for the diagnosis of the condition in athletes, was not utilized by any of the participating centers. Adherence to mannitol provocation guidelines was not considered further, due to the nature of this bronchial provocation challenge being typically well standardized.

The lack of standardization for exercise challenge testing however is of concern. Many variables including exercise intensity, duration, humidity, temperature and time since medication will affect the airway response to exercise in individuals with EIB.⁶Of these factors, our data suggest that medication restrictions were often not detailed and environmental conditions (i.e., temperature and humidity of inspired air) were rarely considered in the SOPs. As the inhalation of dry air is the main contributing factor in the pathophysiology of EIB,⁷lack of consideration of this variable will undoubtedly result in missed diagnoses. Recommendations to overcome this problem include the use of an environmental chamber, which can reduce and control the temperature and humidity of inspired air. However this method is often limited to research facilities or sport science departments. Or, more practical for a hospital setting, the inhalation of medical grade air which is dry.⁴Crucially, our data indicated that these recommendations are scarcely adhered to in clinical practice. As data were retrieved directly from the SOPs provided, we do not foresee any problems with recall bias in the study, however we acknowledge a risk of non-response bias in the sample.

The problem of poor standardization of exercise challenge testing goes beyond clinical practice and may affect the diagnostic guidelines themselves. Indeed, previous research suggest that environmental

Accepted Article
conditions are rarely reported or considered in diagnostic studies for EIB,⁸ even those used to inform clinical guidelines. For example, NICE guidelines on asthma diagnosis, monitoring and management state that exercise challenge tests should not be used for the diagnosis of asthma in persons aged 17 and over;⁹ a recommendation based on findings from five studies. It is therefore highly likely that inferences from poorly performed exercise provocation challenges are informing clinical guidelines.

To conclude, our data show that poor adherence to exercise bronchial provocation guidelines is commonplace in secondary care. The consequence is potentially widespread under-detection of EIB. More effort must be made to educate practitioners in the importance of controlling environmental conditions and ensure strict adherence to exercise provocation guidelines. Where adherence to the environmental criteria are not feasible, a surrogate indirect bronchial provocation challenge (e.g., mannitol, eucapnic voluntary hyperpnoea) should be performed.

Acknowledgements

SD and SJF are supported by the NIHR Manchester Biomedical Research Centre.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Author contributions

AJS (guarantor), SMD and SJF planned the research study. SMD collected the data. AJS, SMD SJF analyzed results. AJS drafted the manuscript. AJS, SMD and SJF approved the final version of the manuscript.

References

1. Parsons JP, Hallstrand TS, Mastronarde JG, Kaminsky DA, Rundell KW, Hull JH, et al. An official American Thoracic Society clinical practice guideline: exercise-induced bronchoconstriction. *Am J Respir Crit Care Med*. 2013;187(9):1016-27.
2. Hull JH, Hull PJ, Parsons JP, Dickinson JW, Ansley L. Approach to the diagnosis and management of suspected exercise-induced bronchoconstriction by primary care physicians. *BMC Pulm Med*. 2009;9:29.
3. Jackson A, Allen H, Hull JH, Hopker J, Backhouse SH, Price OJ, et al. Diagnosing exercise-induced bronchoconstriction: Over-or under-detection? *Allergy*. 2019:00:1–4.
4. Crapo RO, Casaburi AL, Coates PI, et al. Guidelines for methacholine and exercise challenge testing—1999. *Am J Respir Crit Care Med*, 2000;161:309-329
5. Simpson AJ, Romer LM, Kippelen P. Self-reported symptoms after induced and inhibited bronchoconstriction in athletes. *Med Sci Sports Exerc*, 2015;47:2005-2013
6. Anderson SD, Kippelen P. Assessment of EIB: What you need to know to optimize test results. *Immunol Allergy Clin North Am*, 2013;33:363-380
7. Anderson SD, Daviskas E. The mechanism of exercise-induced asthma is *J Allergy Clin Immunol*, 2000;106:453–459
8. Stickland MK, Spooner CH, Dryden DM, Rowe BH. The need for standardization in exercise challenge testing for exercise-induced asthma/bronchoconstriction. *J Allergy Clin Immunol*, 2010;126:878-880
9. Asthma: diagnosis, monitoring and chronic asthma management, NICE NG80, November 2017. <https://www.nice.org.uk/guidance/ng80> (accessed 27 September 2019).

Table 1. Exercise bronchial provocation specifications from exercise challenge testing guidelines^{1,4}

Guideline domain	Guideline Specification
Medication restriction	Short-acting bronchodilators, 8h; long acting bronchodilators, 48h; cromolyn sodium, 8h; nedocromil, 48h; anti-histamines, 3d; leukotriene modifiers, 24 h.
Exercise intensity	Monitored via heart rate (HR) or ventilation (VE). Target HR >80% of predicted max (220 – age in years). VE 40-60% of predicted max maximum voluntary ventilation (FEV ₁ x 35).
Exercise duration	Total duration 6-8 minutes with 4-6 minutes at exercise intensity described above.
Environmental conditions (inspired air)	A nose clip should be worn and inhaled water content <10mg/l or <25°C with <50% relative humidity. Alternatively, dry air delivered from medical grade compressed gas source.
Post-challenge spirometry timings	Spirometry performed pre-test and at 5, 10, 15, 20 and 30 minutes post-exercise.

FEV₁, Forced Expiratory Volume in 1 second.

Figure 1. Adherence to exercise bronchial provocation guideline specifications^{1,4}

