

1 **Perfluorinated alkyl acids in the serum and follicular fluid of UK women with and without**  
2 **polycystic ovarian syndrome undergoing fertility treatment and associations with hormonal and**  
3 **metabolic parameters**

4 Heffernan AL<sup>1,2\*</sup>, Cunningham TK,<sup>3,4\*</sup> Drage DS,<sup>1,5</sup> Aylward LL,<sup>1,6</sup> Thompson K,<sup>1</sup> Vijayasarathy S,<sup>1</sup>  
5 Mueller JF<sup>1</sup>, Atkin SL,<sup>7</sup> Sathyapalan T<sup>3#</sup>

6  
7 1. Queensland Alliance for Environmental Health Sciences, The University of Queensland, Australia

8 2. The Florey Institute of Neuroscience and Mental Health, Melbourne, Australia

9 3. Centre for Cardiovascular and Metabolic Research, Hull York Medical School, University of Hull

10 4. The Hull IVF Unit, Women's and Childrens' Hospital, Hull Royal Infirmary, Hull, UK;

11 5. School of Geography, Earth and Environmental Sciences, University of Birmingham, UK

12 6. Summit Toxicology, LLP, USA

13 7. Weill Cornell Medicine Qatar, Education City, Qatar Foundation, Doha, Qatar

14 8. The Hull IVF Unit, Womens and Childrens Hospital, Hull Royal Infirmary, Hull, UK

15

16 \*Joint first authors

17

18 # Corresponding Author

19 T Sathyapalan

20 Michael White Diabetes Centre, Hull Royal Infirmary, Hull, UK, HU3 2RW

21 Tel +441482675312

22 Fax +441482675370

23 Email [thozhukat.sathyapalan@hyms.ac.uk](mailto:thozhukat.sathyapalan@hyms.ac.uk)

24 **Abstract**

25 Women undergoing treatment for infertility could be a sensitive subpopulation for endocrine effects  
26 of exposure to perfluorinated alkyl acids (PFAAs), persistent organic pollutants with potential  
27 endocrine activity. Women with polycystic ovarian syndrome (PCOS, n=30) and age- and BMI-matched  
28 controls (n=29) were recruited from a UK fertility clinic in 2015. Paired serum and follicular fluid  
29 samples were collected and analysed for 13 PFAAs. Sex steroid and thyroid hormones, metabolic  
30 markers, and serum biochemical parameters were measured and assessed for associations with serum  
31 PFAAs. Four PFAAs were detected in all serum and follicular fluid samples and concentrations in the  
32 two matrices were highly correlated ( $R^2 > 0.95$ ): perfluorooctane sulfonate (PFOS), perfluorooctanoic  
33 acid (PFOA), perfluorohexane sulfonate (PFHxS), and perfluorononanoic acid (PFNA). Serum PFOS was  
34 positively associated with age ( $p < 0.05$ ) and was higher in PCOS cases than controls (geometric mean  
35 3.9 vs. 3.1 ng/mL,  $p < 0.05$ ) and in women with irregular vs. regular menstrual cycles ( $p < 0.05$ ). When  
36 adjusted for PCOS case status and serum albumin, serum testosterone was positively associated with  
37 PFOA, and sex hormone binding globulin was positively associated with PFOS ( $p < 0.05$ ); no other  
38 associations between sex steroid or thyroid hormones and PFAA concentrations were observed.  
39 Fasting glucose was significantly positively associated with PFOA, adjusted for age, PCOS status, and  
40 serum albumin ( $p < 0.05$ ). Serum insulin and HbA1c were positively associated with BMI ( $p < 0.01$ ), but  
41 not with PFAAs. Serum PFAA concentrations can be used as surrogates for follicular fluid  
42 concentrations due to the high correlations observed. Limited associations between serum PFAAs and  
43 sex steroid hormones and fasting glucose were observed. Associations were modified by serum  
44 albumin, which can influence serum PFAA concentrations, and these interrelationships should be  
45 considered in assessing endocrine associations for PFAAs.

46

47 **Key words:** polycystic ovary syndrome; IVF; PFAA, endocrine disrupting chemicals; perfluorinated alkyl  
48 acids; PFOS; PFNA

## 49 Introduction

50 Perfluorinated alkyl acids (PFAAs; perfluorinated chemicals (PFCs)) consist of a fluorinated  
51 hydrophobic alkyl chain with a hydrophilic end group, and are used widely as surfactants in household  
52 and industrial applications such as textile treatments, food packaging, and as aqueous film-forming  
53 foams. The dominant exposure pathway for humans is diet, particularly meat and fish, and via breast  
54 milk for infants ([Gebbinck et al., 2015](#); [Haug et al., 2010](#); [Kärman et al., 2007](#)). PFAAs are persistent  
55 and bioaccumulative. Elimination of PFAAs depends on chain length and they sequester particularly in  
56 the liver and kidney. They are non-covalently bound to protein in serum, particularly serum albumin  
57 ([Andersen et al., 2008](#); [Bischel et al., 2010](#)). Serum elimination half-life is approximately 3.8 and 5.4  
58 years for perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA), respectively ([Li et al.,](#)  
59 [2017b](#); [Olsen et al., 2007](#)). PFAAs can cross the placenta ([Kim et al., 2011](#)), and have been associated  
60 with adverse effects on fertility, birth outcomes, and early development in humans ([Bach et al., 2015](#);  
61 [Goudarzi et al., 2016](#); [Lyngso et al., 2014](#); [Olsen et al., 2009](#)). PFOS and PFOA have intrinsic estrogenic  
62 activity and anti-estrogenic effects *in vitro* ([Henry and Fair, 2013](#)), and PFOS is capable of modulating  
63 steroidogenesis ([Kraugerud et al., 2011](#)). *In vivo*, PFAAs were associated with increased breast cancer  
64 risk in Inuit women, perhaps related to their estrogenic effects ([Bonefeld-Jorgensen et al., 2011](#)), while  
65 other studies have shown increased serum PFAAs associated with an earlier menopause, and with  
66 PFOS being inversely associated with estradiol levels ([Knox et al., 2011](#)). Because PFAAs are eliminated  
67 via both menstruation and renal elimination, it may be difficult to assess and interpret relationships  
68 between serum PFAA concentrations and outcomes such as birth weight, which can be affected by  
69 glomerular filtration rates, or timing of menopause, which can influence PFAA levels due to decreased  
70 elimination of PFAAs post-menopause ([Ruark et al., 2017](#); [Verner et al., 2015](#)).

71

72 Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders and affects 6-20%  
73 of reproductive-aged women ([Bozdag et al., 2016](#); [March et al., 2010](#); [Teede et al., 2013](#); [Yildiz et al.,](#)  
74 [2012](#)) with clinical manifestations of irregular menstruation, hyperandrogenism and/or polycystic  
75 ovaries ([Bozdag et al., 2016](#); [PCOS Consensus Workshop, 2004](#)). PCOS is associated with infertility,  
76 hirsutism, and acne ([Ehrmann, 2005](#); [Norman et al., 2007](#)). Thus, PCOS patients may be a sensitive  
77 subpopulation for compounds that alter endocrine outcomes. The previously reported association of  
78 PFAAs with menstrual irregularity and infertility ([Lyngso et al., 2014](#); [Velez et al., 2015](#)) may have been  
79 influenced by the inclusion of women with PCOS in the studies. The aim of the study was to examine  
80 correlation of serum and follicular fluid measures of PFAAs, and to explore the associations of PFAAs  
81 with hormonal parameters in women with and without PCOS, and undergoing fertility treatment.

82

## 83 **Materials and Methods**

84 This prospective cohort study was performed within the Hull IVF Unit, UK following approval by The  
85 Yorkshire and The Humber NRES ethical committee, UK (approval number 02/03/043). The PCOS  
86 subjects were recruited sequentially in 2015, using the revised 2003 criteria from the Rotterdam  
87 ESHRE/ASRM sponsored PCOS consensus workshop group, indicating PCOS to be present if any 2 out  
88 of 3 criteria were met: menstrual disturbance (oligo or amenorrhoea), clinical and/or biochemical  
89 signs of androgenism or polycystic ovaries on ultrasound ([PCOS Consensus Workshop, 2004](#)).  
90 Inclusion criteria were age 20-45 years, BMI  $\leq 35$  and undergoing *in vitro* fertilisation. Patients with  
91 known immunological disease, diabetes, renal or liver insufficiency, acute or chronic infections, or  
92 inflammatory diseases were excluded from the study. No comparative study on which to base formal  
93 power calculations was available; therefore, power and sample size for pilot studies has been  
94 reviewed ([Birkett and Day, 1994](#)) that concluded that a minimum of 20 degrees-of-freedom was  
95 required to estimate effect size and variability. Hence, we planned to recruit 25 patients per group  
96 with an additional 5 patients allowing for drop-outs and covariate adjustment. A total of 59 women  
97 were recruited into the study, 30 PCOS cases and 29 control subjects matched for age and weight.

98

### 99 *Sample Collection*

100 The subjects fasted from midnight and had a fasting blood sample taken on day 21 of the luteal phase  
101 of the cycle before commencing their IVF treatment. Fasting venous blood samples were collected,  
102 separated by centrifugation at 3500 x g for 15 min at 4°C, and the aliquots stored at -80°C within 1  
103 hour of collection. Plasma glucose was measured using a Synchron LX20 analyzer (Beckman-Coulter),  
104 and serum insulin was assayed using a competitive chemiluminescent immunoassay performed using  
105 the DPC Immulite 2000 analyzer (Euro/DPC, Llanberis, UK). C reactive protein (CRP) was measured  
106 enzymatically using a Synchron LX20 analyzer (Beckman-Coulter, UK). Estradiol and all thyroid assays  
107 were performed on an Abbott Architect i4000 immunoassay analyser (Abbott Diagnostics Division,  
108 UK). Serum testosterone and androstenedione were measured by liquid chromatography tandem  
109 mass spectrometry (LC/MS/MS; Acquity UPLC-Quattro Premier XE-MS, Waters, Manchester, UK). Sex  
110 hormone binding globulin (SHBG) was measured by an immunometric assay with fluorescence  
111 detection (DPC Immulite 2000 analyzer; upper limit 2.0 nmol/l). Glycosylated hemoglobin A1c (HbA1c)  
112 measurements were made using ion-exchange chromatography.

113

### 114 *Analysis for PFAAs*

115 Samples were analysed for 13 PFAAs including PFOS, PFOA, perfluorohexane sulfonate (PFHxS),  
116 perfluorononanoic acid (PFNA), and perfluorodecanoic acid (PFDA) (Table 1). 200  $\mu$ L serum or follicular

117 fluid was transferred to a 2 mL Eppendorf tube, followed by addition of the internal standards.  
118 Proteins were precipitated with acetonitrile, centrifuged, filtered (2 µm GHP membrane; Pall, East  
119 Hills, NY, USA), and concentrated under gentle stream of nitrogen. Samples were reconstituted in  
120 5mM ammonium acetate in water prior to analysis via high performance liquid chromatography  
121 tandem mass spectrometry (HPLC-MS/MS) using a Nexera HPLC (Shimadzu Corp., Kyoto, Japan)  
122 coupled to API5500 QTRAP mass spectrometer (Sciex, Melbourne, Australia) with electrospray  
123 ionization (ESI) interface operating in negative mode. Chromatographic separation of the analytes was  
124 achieved with a Gemini C<sub>18</sub> column (50x2.0 mm, 4 µm; Phenomenex, Torrance, CA), maintained at 45  
125 °C, with a flow rate of 0.3 mL/min and injection volume of 5 µL. Mobile phases consisted of  
126 methanol:water (1:99, v/v) (A), and methanol:water (95:5, v/v) (B), with 5mM ammonium acetate in  
127 both phases. An isolator column (Phenomenex) was included inline directly after the mobile phase  
128 mixing chamber to delay the elution of solvent-derived background PFAA contamination. Data  
129 acquisition and processing was carried out using Analyst® TF 1.6 and MultiQuant™ software (Sciex).  
130 Further details of reagents, standards, and mass spectrometry settings are provided in the  
131 Supplementary Material.

132

### 133 *Statistics.*

134 Descriptive data are presented as mean ± SD for continuous data and n (%) for categorical data. T-  
135 tests were used to compare means where appropriate. A p-value of <0.05 was considered to indicate  
136 statistical significance except for exploratory Pearson correlation coefficient evaluations for regression  
137 modelling development (p<0.1).

138

139 Measured serum PFAA, hormone concentrations, and metabolic markers were assessed for normality  
140 and ln-transformed where appropriate. Estimated glomerular filtration rate (eGFR) was calculated  
141 using the Modification of Diet in Renal Disease (MDRD) study method ([Levey et al., 1999](#)). Insulin  
142 resistance (IR) was calculated from basal glucose and insulin concentration using the homeostasis  
143 model assessment (HOMA) ((Insulin x glucose)/22.5)([Matthews et al., 1985](#)). Free androgen index  
144 (FAI) was calculated as 100 times the ratio of serum testosterone and SHBG concentrations. A sum  
145 PFAA (ΣPFAA) variable by calculated by adding the molar concentrations of the four frequently  
146 detected PFAA compounds (i.e. sum of PFOS, PFOA, PFHxS, PFNA).

147

148 Pairwise Pearson correlation coefficients and significance were examined as an initial step in assessing  
149 potential associations and identifying potential covariates. Multivariable linear regression was used  
150 to assess predictors for PFAA concentrations and potential associations between measured hormone  
151 concentrations or metabolic endpoints and serum PFAA concentrations. Tobit regressions were used

152 to examine oestradiol concentrations due to left censoring of the data (concentrations below 75  
153 pmol/L were not quantified). Statistical analyses were conducted using Stata (IC 12.1, Stata Corp.,  
154 College Station, TX).

155

## 156 **Results**

157 Cases and controls were similar in age, BMI, and age at menarche (Table 2). Measured hormone  
158 concentrations were not available for all cases and controls (see Table 2). PCOS cases more frequently  
159 had irregular menstrual cycles (87% vs. 14%,  $p < 0.001$ ). PCOS cases were more likely to be taking  
160 metformin (47% vs. 0%), and had significantly lower average fasting glucose concentrations than  
161 controls (4.4 nmol/L vs. 4.9 nmol/L,  $p < 0.01$ ), though HbA1c did not differ to controls.

162

163 PCOS cases had higher androgen levels with significantly elevated FAI and androstenedione compared  
164 to controls, though testosterone and oestradiol did not differ. PCOS cases also had lower eGFR on  
165 average than controls (88.3 vs. 97.4 mL/min/ 1.73m<sup>2</sup>,  $p < 0.05$ ). Serum insulin, HOMA-IR and CRP did  
166 not differ between PCOS and controls.

167

168 Detection frequencies and descriptive statistics for serum and follicular fluid PFAA concentrations are  
169 presented in Table 1. Four PFAAs were detected in all serum and follicular fluid samples: PFOS, PFOA,  
170 PFHxS, and PFNA, and all were significantly correlated with one another. Detection frequencies for  
171 PFDA were 76%; and <50% for PFPeA, PFBS, PFHpA and PFUnDA, (49, 7, 17 and 36%, respectively). In  
172 general, PFOS was present at the highest concentrations, followed by PFOA, PFHxS, and PFNA.  
173 Geometric mean serum concentrations of PFOS were significantly greater in PCOS cases than controls  
174 (Table 1, Figure 1). Other serum PFAAs were not significantly different between PCOS and controls,  
175 and geometric mean follicular fluid concentrations were not different between groups (Table 2), due  
176 to greater variation in measured PFAS concentrations. For the four frequently detected PFAAs,  
177 concentrations in follicular fluid were highly correlated with serum concentrations ( $R^2 > 0.95$  for all  
178 four, Figure 2). The mean ratios of follicular fluid to serum concentrations were 0.59, 0.78, 0.86, and  
179 0.77 for PFOS, PFOA, PFHxS, and PFNA, respectively (Table 1).

180

181 Patients with irregular menstrual cycles (from both PCOS and control groups;  $n=30$ , GM: 4.16 ng/mL)  
182 had significantly higher PFOS concentrations than those with regular cycles ( $n=29$ , GM 3.25 ng/mL)  
183 ( $p=0.011$ , 2-tailed t-test; Figure 3), but the PFOS concentration was not associated with the degree of  
184 irregularity in PCOS patients (not shown). No associations between other PFAA concentrations and

185 menstrual cycle regularity were observed. No associations between parity (nulliparous versus  
186 primiparous) and PFAA concentrations were observed (data not shown).

187

188 We examined pair-wise Pearson correlations for the ln-transformed concentrations of the frequently  
189 detected PFAAs and age, BMI, serum albumin, and ln-transformed eGFR (Supplementary Information,  
190 Tables S1 to S3). No correlations with BMI were observed for any of the PFAAs. PFOS and PFNA were  
191 negatively correlated with eGFR and positively correlated with serum albumin. Based on the  
192 correlation matrix, we examined predictors for each PFAA concentration in multivariable regressions  
193 with dependent variables of age, ln-transformed eGFR, serum albumin, and PCOS status. PFOS was  
194 significantly positively associated with age (approximately 0.1 ng/ml increase per year of age) and  
195 status as a PCOS case vs. control. PFNA was negatively associated with eGFR. No other statistically  
196 significant predictors for PFAA concentrations were identified.

197

198 Associations between metabolic endpoints (fasting glucose, serum insulin, HbA1C, and HOMA-IR) and  
199 serum PFAAs were assessed considering potential confounders. In pairwise correlations, fasting  
200 glucose was positively correlated with age and negatively correlated with serum albumin and status  
201 as a PCOS case; these variables were retained in multivariable regressions examining potential  
202 associations between fasting glucose and ln-transformed PFAA concentrations. Significant positive  
203 associations were observed between fasting glucose and ln-transformed PFOA ( $\beta=0.18$ , 95% CI  
204 0.01,0.36,  $p=0.035$ ; i.e. 1.2 ng/mL increase in PFOA for every 1nmol/L increase in glucose) and ln-  
205 transformed  $\sum PFAA$  ( $\beta=0.18$ , 95% CI 0.00,0.37,  $p=0.05$ ; i.e. 1.2  $\mu\text{mol/L}$  increase in  $\sum PFAA$  for every  
206 1nmol/L increase in glucose). Serum insulin, HbA1c, and HOMA-IR were all significantly positively  
207 associated with BMI, but no significant associations with any of the PFAAs were observed. These  
208 results were not affected by inclusion of metformin in use in the analysis or stratification by metformin  
209 use.

210

211 We examined associations between ln-transformed concentrations of each of the four frequently  
212 detected PFAAs and steroid hormone concentrations, SHBG concentrations, and FAI, adjusting for  
213 status as a PCOS case vs. control, as well as for serum albumin concentrations (Table 3).  $\sum PFAA$  was  
214 assessed in association with hormone concentrations. Serum testosterone concentrations were  
215 positively associated with ln-transformed PFOA and with  $\sum PFAA$ , though not with the degree of  
216 testosterone elevation. SHBG concentrations were significantly positively associated with PFOS and  
217 with  $\sum PFAA$ . No other significant associations between measured steroid hormone levels and serum  
218 PFAA concentrations were observed in adjusted models (Table 3).

219

220 No significant associations between any of the frequently detected PFAAs and TSH, fT3, or fT4 were  
221 observed. Both fT3 and fT4 were significantly negatively associated with serum albumin, consistent  
222 with non-specific binding of total T3 and T4 to serum protein (not shown).

223

## 224 **Discussion**

225 The potential effects of PFAAs on reproductive and thyroid hormones and potential influences on  
226 reproductive health are of interest given previous reports linking PFAAs to alterations in endocrine  
227 activity and function ([Bach et al., 2016](#); [Coperchini et al., 2017](#)). The profile of PCOS also includes  
228 alterations in reproductive hormones; thus, this population might represent a sensitive subpopulation  
229 for chemical exposures that also influence hormone concentrations.

230

231 The four frequently detected PFAAs were correlated with each other in this study, consistent with  
232 previous reports ([Calafat et al., 2007](#); [Ye et al., 2017](#)). Concentrations in follicular fluid of each PFAA  
233 were strongly associated with the corresponding serum concentrations, with average ratios ranging  
234 from approximately .59 to .86 for the four frequently detected PFAAs (Table 1). These ratios are  
235 similar to those reported by McCoy et al. (2017) ([McCoy et al., 2017](#)) for PFOA, PFHxs, and PFNA, but  
236 somewhat lower than the value reported for PFOS (0.59 in the current study vs. 0.82 in McCoy et al.  
237 2017). The lower concentrations in follicular fluid relative to serum might reflect a lower total protein  
238 concentration in follicular fluid relative to serum ([Leroy et al., 2004](#)), which is pertinent because PFAAs  
239 are known to be protein bound ([Andersen et al., 2008](#); [Bischel et al., 2010](#)). The high correlations  
240 between PFAA concentrations in follicular fluid and serum suggest that measures of PFAAs in serum  
241 likely are good surrogates for examining potential dose-effect relationships on the ovary.

242

243 The assessment of associations between PFAA levels and sex steroid hormones resulted in limited  
244 findings of a positive association between testosterone and PFOA and the molar sum of PFAAs after  
245 adjusting for serum albumin concentrations and PCOS case status; no significant findings remained  
246 after adjustment for the other hormone-PFAA combinations (Table 3). When the data were restricted  
247 to PCOS cases alone there was no linear relationship of PFOS to increased testosterone. This may be  
248 due to a combination of the small sample size (n= 27 PCOS cases with measured testosterone) and the  
249 high degree of variation in testosterone levels in the PCOS cases. In a recent systematic review, Bach  
250 et al. (2016) reported that associations between reproductive hormone levels and PFAA exposures  
251 was mixed ([Bach et al., 2016](#)).

252



253 TSH, free T3, and free T4 were not associated with any of the frequently detected PFAAs or the sum  
254 of these PFAAs in this study. We did observe a negative correlation between serum albumin and free  
255 T4 and free T3. Previous studies have reported mixed results regarding associations between PFAAs  
256 and thyroid hormones. Crawford et al. (2017) in a study of women without infertility found no  
257 associations between TSH and PFAAs, and reported a positive association between free T4 and PFNA  
258 ([Crawford et al., 2017](#)). Lin et al. (2013) found a similar relationship between free T4 and PFNA in  
259 adolescents and young adults from the NHANES survey ([Lin et al., 2013](#)). Chan et al. 2011 found no  
260 associations between serum PFAA concentrations and hypothyroxinemia in 974 pregnant women  
261 ([Chan et al., 2011](#)). Lewis et al. (2015) found a positive association between free T4 and serum  
262 concentrations of all four PFAAs considered here in women of reproductive age in the NHANES 2011-  
263 2012 survey, but other thyroid hormone concentrations were not associated with PFAAs in women of  
264 reproductive age ([Lewis et al., 2015](#)). Overall, studies have reported a mixed pattern of associations  
265 between PFAAs and thyroid hormone concentrations ([de Cock et al., 2014](#); [Jain, 2013](#); [Ji et al., 2012](#);  
266 [Kato et al., 2016](#); [Li et al., 2017a](#); [Shah-Kulkarni et al., 2016](#); [Tsai et al., 2017](#)).

267  
268 PCOS cases had higher geometric mean concentrations of PFOS than controls, but concentrations of  
269 other PFAAs were similar between cases and controls. Vagi et al. (2014) found elevated PFOS and  
270 PFOA concentrations in another study of PCOS cases relative to controls ([Vagi et al., 2014](#)). The current  
271 study also found an association of PFOS concentrations with menstrual irregularity, similar to one  
272 previous study ([Lyngso et al., 2014](#)). Menstruation may be an important elimination pathway for  
273 PFAAs, and has been hypothesized to be responsible for lower PFAA concentrations in females  
274 compared to males ([Lorber et al., 2015](#); [Wong et al., 2014](#)) and in post-menopausal women ([Lorber et  
275 al., 2015](#)). In PCOS women who may suffer from oligomenorrhea or amenorrhea, and thus menstruate  
276 less frequently than controls, the same exposure dose could result in higher serum PFAAs  
277 concentrations ([Vagi et al., 2014](#)). However, for the PCOS women, there was no significant difference  
278 in PFOS concentration between cases with cycles greater than or less than 40 days. Similarly, in this  
279 dataset, PFAAs other than PFOS were not significantly associated with menstrual cycle regularity.

280  
281 In addition, we found that PCOS cases had significantly lower eGFR than controls. Evidence from the  
282 literature suggestd this may be due to inflammation and reflected in a higher CRP ([Gozukara et al.,  
283 2015](#)); however, in our study, CRP did not differ between PCOS and controls. . Renal elimination is  
284 another pathway for elimination of PFAA compounds ([Han et al., 2012](#); [Verner et al., 2015](#)). Thus,  
285 lower eGFR for PCOS cases may result in higher serum PFAA concentrations for the same external  
286 exposure level, which was observed for PFOS, and potentially compounded by menstrual irregularity,

287 thereby reducing PFAA elimination. Previous cross-sectional studies considered the possibility that  
288 PFAAs may negatively impact renal function, resulting in decreased eGFR ([Kataria et al., 2015](#); [Watkins  
289 et al., 2013](#)). In this dataset, PFOS and PFNA were inversely correlated with eGFR ( $p < 0.05$ ); PFOA and  
290 PFHxS were not significantly correlated with eGFR. Thus, decreased eGFR in PCOS cases may result in  
291 some increase in serum concentrations of selected PFAA compounds in these cases compared to  
292 controls, but this relationship may be compound-specific.

293

294 Recent temporal studies report decreasing serum concentrations of PFOS and PFOA, and a  
295 corresponding increase of alternative PFAAs used as replacement chemicals, likely due to action from  
296 manufacturers and legislators to phase out PFOS and PFOA ([Land et al., 2015](#); [US EPA, 2015](#)). In  
297 comparison with the biomonitoring literature, PFOS concentrations in this UK cohort were lower than  
298 most previously reported pregnancy cohorts globally (reviewed in ([Miralles-Marco and Harrad, 2015](#))),  
299 whereas PFOA and PFHxS were higher (Table S4).

300

301 The strengths of the study lie in the age and BMI matched population, with measurement of a range  
302 of hormone and metabolic markers. In addition, PCOS patients may represent a sensitive  
303 subpopulation for effects of endocrine active substances. The study was limited by the small sample  
304 size and by missing data for some hormone measurements. A relatively large number of statistical  
305 evaluations were conducted (approximately 12 outcome measures by four PFAAs and the molar sum  
306 of the four PFAAs, plus assessments of predictors for PFAA concentrations), suggesting that some  
307 findings might be expected to be observed by chance.

308

309 We found high correlations between PFAAs in follicular fluid and serum, supporting the use of serum  
310 as a relevant matrix for biomonitoring for PFAAs for assessment of potential ovarian responses.  
311 Concentrations of PFOS, but not other PFAAs, were higher in the PCOS cases than in controls in this  
312 study. We found evidence of a positive associations between PFOA concentrations and summed PFAA  
313 concentrations and testosterone, and between PFOS and summed PFAAs and SHBG concentrations in  
314 the PCOS cases and controls in this study. The relationships to summed PFAAs appear to be largely  
315 driven by the contributions from the individual significant predictors; that is, the relationships to  
316 summed PFAAs are not stronger in magnitude or significance than the relationships to PFOA or PFOS.  
317 Fasting glucose was positively associated with PFOA concentrations and with summed PFAAs, but not  
318 other PFAAs. Again, the relationship with summed PFAAs appears to be largely due to the specific  
319 association with PFOA.

320

321 We identified a number of factors that should be considered in the evaluation of PFAA concentrations  
322 and potential associations with reproductive outcomes, steroid hormone concentrations, and related  
323 endpoints. Characteristics such as menstrual irregularity, parity, eGFR, and serum albumin levels may  
324 influence serum PFAA concentrations due to elimination mechanisms or physical/chemical properties  
325 of these compounds, and some of these characteristics may also be altered in populations under study  
326 for reproductive outcomes. In addition, serum albumin concentrations can influence both measured  
327 serum hormone concentrations and serum concentrations of PFAAs. Non-covalent binding of sex  
328 steroid hormones to serum protein is recognized as a factor affecting transport and metabolism of  
329 these hormones ([Egloff et al., 1981](#); [Pardridge, 1986](#)), and we observed positive associations between  
330 serum albumin concentrations and the measured levels of the sex steroid hormones. Similarly, T3,  
331 and T4 are protein bound in serum ([Koulouri et al., 2013](#)). We observed a negative correlation  
332 between free T3 and free T4 and albumin in this dataset (Table S2), consistent with this protein  
333 binding. PFAAs are also protein-bound in serum. These interrelationships suggest that characteristics  
334 that influence PFAA elimination or serum concentrations, and which may also be associated with  
335 outcome variables, be carefully considered as covariates in future assessments of associations  
336 between hormone concentrations, reproductive outcomes, and serum PFAAs.

337

### 338 **Acknowledgements**

339 ALH receives funding support from the Australian Research Council (ARC) and the National Health and  
340 Medical Research Council (APP1106911); JFM is funded by an ARC Future Fellowship (FF120100546).

341

### 342 **References**

343 Andersen, M.E., Butenhoff, J.L., Chang, S.C., Farrar, D.G., Kennedy, G.L., Jr., Lau, C., Olsen, G.W.,  
344 Seed, J., Wallace, K.B., 2008. Perfluoroalkyl acids and related chemistries--toxicokinetics and  
345 modes of action. *Toxicol Sci* 102, 3-14.

346 Bach, C.C., Bech, B.H., Brix, N., Nohr, E.A., Bonde, J.P., Henriksen, T.B., 2015. Perfluoroalkyl and  
347 polyfluoroalkyl substances and human fetal growth: a systematic review. *Critical reviews in*  
348 *toxicology* 45, 53-67.

349 Bach, C.C., Vested, A., Jorgensen, K.T., Bonde, J.P., Henriksen, T.B., Toft, G., 2016. Perfluoroalkyl and  
350 polyfluoroalkyl substances and measures of human fertility: a systematic review. *Critical*  
351 *reviews in toxicology* 46, 735-755.

352 Birkett, M.A., Day, S.J., 1994. Internal pilot studies for estimating sample size. *Stat Med* 13, 2455-  
353 2463.

354 Bischel, H.N., Macmanus-Spencer, L.A., Luthy, R.G., 2010. Noncovalent interactions of long-chain  
355 perfluoroalkyl acids with serum albumin. *Environ Sci Technol* 44, 5263-5269.

356 Bonefeld-Jorgensen, E.C., Long, M., Bossi, R., Ayotte, P., Asmund, G., Kruger, T., Ghisari, M., Mulvad,  
357 G., Kern, P., Nzulumiki, P., Dewailly, E., 2011. Perfluorinated compounds are related to breast  
358 cancer risk in Greenlandic Inuit: a case control study. *Environ Health* 10, 88.

359 Bozdag, G., Mumusoglu, S., Zengin, D., Karabulut, E., Yildiz, B.O., 2016. The prevalence and  
360 phenotypic features of polycystic ovary syndrome: a systematic review and meta-analysis. *Hum  
361 Reprod* 31, 2841-2855.

362 Calafat, A.M., Kuklenyik, Z., Reidy, J.A., Caudill, S.P., Tully, J.S., Needham, L.L., 2007. Serum  
363 Concentrations of 11 Polyfluoroalkyl Compounds in the U.S. Population: Data from the National  
364 Health and Nutrition Examination Survey (NHANES) 1999–2000. *Environmental Science &  
365 Technology* 41, 2237-2242.

366 Chan, E., Burstyn, I., Cherry, N., Bamforth, F., Martin, J.W., 2011. Perfluorinated acids and  
367 hypothyroxinemia in pregnant women. *Environmental research* 111, 559-564.

368 Coperchini, F., Awwad, O., Rotondi, M., Santini, F., Imbriani, M., Chiovato, L., 2017. Thyroid  
369 disruption by perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA). *Journal of  
370 endocrinological investigation* 40, 105-121.

371 Crawford, N.M., Fenton, S.E., Strynar, M., Hines, E.P., Pritchard, D.A., Steiner, A.Z., 2017. Effects of  
372 perfluorinated chemicals on thyroid function, markers of ovarian reserve, and natural fertility.  
373 *Reproductive toxicology (Elmsford, N.Y.)* 69, 53-59.

374 de Cock, M., de Boer, M.R., Lamoree, M., Legler, J., van de Bor, M., 2014. Prenatal exposure to  
375 endocrine disrupting chemicals in relation to thyroid hormone levels in infants – a Dutch  
376 prospective cohort study. *Environmental health : a global access science source* 13, 106.

377 Egloff, M., Savoure, N., Tardivel-Lacombe, J., Massart, C., Nicol, M., Degrelle, H., 1981. Influence of  
378 sex hormone binding globulin and serum albumin on the conversion of androstenedione to  
379 testosterone by human erythrocytes. *Acta endocrinologica* 96, 136-140.

380 Ehrmann, D.A., 2005. Polycystic ovary syndrome. *N Engl J Med* 352, 1223-1236.

381 Gebbink, W.A., Berger, U., Cousins, I.T., 2015. Estimating human exposure to PFOS isomers and PFCA  
382 homologues: the relative importance of direct and indirect (precursor) exposure. *Environment  
383 international* 74, 160-169.

384 Goudarzi, H., Nakajima, S., Ikeno, T., Sasaki, S., Kobayashi, S., Miyashita, C., Ito, S., Araki, A.,  
385 Nakazawa, H., Kishi, R., 2016. Prenatal exposure to perfluorinated chemicals and  
386 neurodevelopment in early infancy: The Hokkaido Study. *The Science of the total environment*  
387 541, 1002-1010.

388 Gozukara, I.O., Gozukara, K.H., Kucur, S.K., Karakilic, E.U., Keskin, H., Akdeniz, D., Aksoy, A.N.,  
389 Carlioglu, A., 2015. Association of Glomerular Filtration Rate with Inflammation in Polycystic  
390 Ovary Syndrome. *International journal of fertility & sterility* 9, 176-182.

391 Han, X., Nabb, D.L., Russell, M.H., Kennedy, G.L., Rickard, R.W., 2012. Renal elimination of  
392 perfluorocarboxylates (PFCAs). *Chemical research in toxicology* 25, 35-46.

393 Haug, L.S., Thomsen, C., Brantsæter, A.L., Kvaalem, H.E., Haugen, M., Becher, G., Alexander, J.,  
394 Meltzer, H.M., Knutsen, H.K., 2010. Diet and particularly seafood are major sources of  
395 perfluorinated compounds in humans. *Environment International* 36, 772-778.

396 Henry, N.D., Fair, P.A., 2013. Comparison of in vitro cytotoxicity, estrogenicity and anti-estrogenicity  
397 of triclosan, perfluorooctane sulfonate and perfluorooctanoic acid. *J Appl Toxicol* 33, 265-272.

398 Jain, R.B., 2013. Association between thyroid profile and perfluoroalkyl acids: data from NHNAES  
399 2007-2008. *Environmental research* 126, 51-59.

400 Ji, K., Kim, S., Kho, Y., Paek, D., Sakong, J., Ha, J., Kim, S., Choi, K., 2012. Serum concentrations of  
401 major perfluorinated compounds among the general population in Korea: dietary sources and  
402 potential impact on thyroid hormones. *Environment international* 45, 78-85.

403 Kärrman, A., Ericson, I., van Bavel, B., Darnerud, P.O., Aune, M., Glynn, A., Lignell, S., Lindström, G.,  
404 2007. Exposure of Perfluorinated Chemicals through Lactation: Levels of Matched Human Milk  
405 and Serum and a Temporal Trend, 1996–2004, in Sweden. *Environmental Health Perspectives*  
406 115, 226-230.

407 Kataria, A., Trachtman, H., Malaga-Dieguez, L., Trasande, L., 2015. Association between  
408 perfluoroalkyl acids and kidney function in a cross-sectional study of adolescents.  
409 *Environmental Health* 14, 89.

410 Kato, S., Itoh, S., Yuasa, M., Baba, T., Miyashita, C., Sasaki, S., Nakajima, S., Uno, A., Nakazawa, H.,  
411 Iwasaki, Y., Okada, E., Kishi, R., 2016. Association of perfluorinated chemical exposure in utero  
412 with maternal and infant thyroid hormone levels in the Sapporo cohort of Hokkaido Study on  
413 the Environment and Children's Health. *Environmental health and preventive medicine* 21, 334-  
414 344.

415 Kim, S., Choi, K., Ji, K., Seo, J., Kho, Y., Park, J., Kim, S., Park, S., Hwang, I., Jeon, J., Yang, H., Giesy, J.P.,  
416 2011. Trans-Placental Transfer of Thirteen Perfluorinated Compounds and Relations with Fetal  
417 Thyroid Hormones. *Environmental Science & Technology* 45, 7465-7472.

418 Knox, S.S., Jackson, T., Javins, B., Frisbee, S.J., Shankar, A., Ducatman, A.M., 2011. Implications of  
419 early menopause in women exposed to perfluorocarbons. *The Journal of clinical endocrinology*  
420 and metabolism 96, 1747-1753.

421 Koulouri, O., Moran, C., Halsall, D., Chatterjee, K., Gurnell, M., 2013. Pitfalls in the measurement and  
422 interpretation of thyroid function tests. *Best practice & research. Clinical endocrinology &*  
423 *metabolism* 27, 745-762.

424 Kraugerud, M., Zimmer, K.E., Ropstad, E., Verhaegen, S., 2011. Perfluorinated compounds  
425 differentially affect steroidogenesis and viability in the human adrenocortical carcinoma  
426 (H295R) in vitro cell assay. *Toxicol Lett* 205, 62-68.

427 Land, M., de Wit, C.A., Cousins, I.T., Herzke, D., Johansson, J., Martin, J.W., 2015. What is the effect  
428 of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of  
429 perfluoroalkyl acids and their precursors in the environment? A systematic review protocol.  
430 *Environmental Evidence* 4, 3.

431 Leroy, J.L., Vanholder, T., Delanghe, J.R., Opsomer, G., Van Soom, A., Bols, P.E., Dewulf, J., de Kruif,  
432 A., 2004. Metabolic changes in follicular fluid of the dominant follicle in high-yielding dairy cows  
433 early post partum. *Theriogenology* 62, 1131-1143.

434 Levey, A.S., Bosch, J.P., Lewis, J.B., Greene, T., Rogers, N., Roth, D., 1999. A more accurate method to  
435 estimate glomerular filtration rate from serum creatinine: a new prediction equation.  
436 *Modification of Diet in Renal Disease Study Group. Annals of internal medicine* 130, 461-470.

437 Lewis, R.C., Johns, L.E., Meeker, J.D., 2015. Serum Biomarkers of Exposure to Perfluoroalkyl  
438 Substances in Relation to Serum Testosterone and Measures of Thyroid Function among Adults  
439 and Adolescents from NHANES 2011–2012. *International Journal of Environmental Research*  
440 *and Public Health* 12, 6098-6114.

441 Li, Y., Cheng, Y., Xie, Z., Zeng, F., 2017a. Perfluorinated alkyl substances in serum of the southern  
442 Chinese general population and potential impact on thyroid hormones. *Scientific reports* 7,  
443 43380.

444 Li, Y., Fletcher, T., Mucs, D., Scott, K., Lindh, C.H., Tallving, P., Jakobsson, K., 2017b. Half-lives of  
445 PFOS, PFHxS and PFOA after end of exposure to contaminated drinking water. *Occup and*  
446 *Environ Med.*

447 Lin, C.Y., Wen, L.L., Lin, L.Y., Wen, T.W., Lien, G.W., Hsu, S.H., Chien, K.L., Liao, C.C., Sung, F.C., Chen,  
448 P.C., Su, T.C., 2013. The associations between serum perfluorinated chemicals and thyroid  
449 function in adolescents and young adults. *Journal of hazardous materials* 244-245, 637-644.

450 Lorber, M., Eaglesham, G.E., Hobson, P., Toms, L.M., Mueller, J.F., Thompson, J.S., 2015. The effect  
451 of ongoing blood loss on human serum concentrations of perfluorinated acids. *Chemosphere*  
452 118, 170-177.

453 Lyngso, J., Ramlau-Hansen, C.H., Hoyer, B.B., Stovring, H., Bonde, J.P., Jonsson, B.A., Lindh, C.H.,  
454 Pedersen, H.S., Ludwicki, J.K., Zvezdai, V., Toft, G., 2014. Menstrual cycle characteristics in

455 fertile women from Greenland, Poland and Ukraine exposed to perfluorinated chemicals: a  
456 cross-sectional study. *Human reproduction (Oxford, England)* 29, 359-367.

457 March, W.A., Moore, V.M., Willson, K.J., Phillips, D.I., Norman, R.J., Davies, M.J., 2010. The  
458 prevalence of polycystic ovary syndrome in a community sample assessed under contrasting  
459 diagnostic criteria. *Hum Reprod* 25, 544-551.

460 Matthews, D.R., Hosker, J.P., Rudenski, A.S., Naylor, B.A., Treacher, D.F., Turner, R.C., 1985.  
461 Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma  
462 glucose and insulin concentrations in man. *Diabetologia* 28, 412-419.

463 McCoy, J.A., Bangma, J.T., Reiner, J.L., Bowden, J.A., Schnorr, J., Slowey, M., O'Leary, T., Guillette,  
464 L.J., Jr., Parrott, B.B., 2017. Associations between perfluorinated alkyl acids in blood and ovarian  
465 follicular fluid and ovarian function in women undergoing assisted reproductive treatment. *The*  
466 *Science of the total environment* 605-606, 9-17.

467 Miralles-Marco, A., Harrad, S., 2015. Perfluorooctane sulfonate: a review of human exposure,  
468 biomonitoring and the environmental forensics utility of its chirality and isomer distribution.  
469 *Environment international* 77, 148-159.

470 Norman, R.J., Dewailly, D., Legro, R.S., Hickey, T.E., 2007. Polycystic ovary syndrome. *Lancet* 370,  
471 685-697.

472 Olsen, G.W., Burris, J.M., Ehresman, D.J., Froehlich, J.W., Seacat, A.M., Butenhoff, J.L., Zobel, L.R.,  
473 2007. Half-Life of Serum Elimination of Perfluorooctanesulfonate, Perfluorohexanesulfonate,  
474 and Perfluorooctanoate in Retired Fluorochemical Production Workers. *Environmental Health*  
475 *Perspectives* 115, 1298-1305.

476 Olsen, G.W., Butenhoff, J.L., Zobel, L.R., 2009. Perfluoroalkyl chemicals and human fetal  
477 development: an epidemiologic review with clinical and toxicological perspectives.  
478 *Reproductive toxicology (Elmsford, N.Y.)* 27, 212-230.

479 Pardridge, W.M., 1986. Serum bioavailability of sex steroid hormones. *Clinics in endocrinology and*  
480 *metabolism* 15, 259-278.

481 PCOS Consensus Workshop, 2004. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop,  
482 Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic  
483 ovary syndrome. *Fertility and sterility* 81, 19-25.

484 Ruark, C.D., Song, G., Yoon, M., Verner, M.A., Andersen, M.E., Clewell, H.J., 3rd, Longnecker, M.P.,  
485 2017. Quantitative bias analysis for epidemiological associations of perfluoroalkyl substance  
486 serum concentrations and early onset of menopause. *Environment international* 99, 245-254.

487 Shah-Kulkarni, S., Kim, B.-M., Hong, Y.-C., Kim, H.S., Kwon, E.J., Park, H., Kim, Y.J., Ha, E.-H., 2016.  
488 Prenatal exposure to perfluorinated compounds affects thyroid hormone levels in newborn  
489 girls. *Environment international* 94, 607-613.

490 Teede, H.J., Joham, A.E., Paul, E., Moran, L.J., Loxton, D., Jolley, D., Lombard, C., 2013. Longitudinal  
491 weight gain in women identified with polycystic ovary syndrome: results of an observational  
492 study in young women. *Obesity (Silver Spring)* 21, 1526-1532.

493 Tsai, M.-S., Lin, C.-C., Chen, M.-H., Hsieh, W.-S., Chen, P.-C., 2017. Perfluoroalkyl substances and  
494 thyroid hormones in cord blood. *Environmental Pollution* 222, 543-548.

495 US EPA, 2015. PFOA Stewardship Program Baseline Summary Report. Accessed 5 Sept 2017.  
496 Available from: [https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/pfoa-](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/pfoa-stewardship-program-baseline-year-summary-report)  
497 [stewardship-program-baseline-year-summary-report](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/pfoa-stewardship-program-baseline-year-summary-report).

498 Vagi, S.J., Azziz-Baumgartner, E., Sjodin, A., Calafat, A.M., Dumesic, D., Gonzalez, L., Kato, K., Silva,  
499 M.J., Ye, X., Azziz, R., 2014. Exploring the potential association between brominated diphenyl  
500 ethers, polychlorinated biphenyls, organochlorine pesticides, perfluorinated compounds,  
501 phthalates, and bisphenol A in polycystic ovary syndrome: a case-control study. *BMC endocrine*  
502 *disorders* 14, 86.

503 Velez, M.P., Arbuckle, T.E., Fraser, W.D., 2015. Maternal exposure to perfluorinated chemicals and  
504 reduced fecundity: the MIREC study. *Hum Reprod* 30, 701-709.

505 Verner, M.A., Luccisano, A.E., Morken, N.H., Yoon, M., Wu, H., McDougall, R., Maisonet, M., Marcus,  
506 M., Kishi, R., Miyashita, C., Chen, M.H., Hsieh, W.S., Andersen, M.E., Clewell, H.J., 3rd,  
507 Longnecker, M.P., 2015. Associations of Perfluoroalkyl Substances (PFAS) with Lower Birth  
508 Weight: An Evaluation of Potential Confounding by Glomerular Filtration Rate Using a  
509 Physiologically Based Pharmacokinetic Model (PBPK). *Environmental health perspectives* 123,  
510 1317-1324.

511 Watkins, D.J., Josson, J., Elston, B., Bartell, S.M., Shin, H.-M., Vieira, V.M., Savitz, D.A., Fletcher, T.,  
512 Wellenius, G.A., 2013. Exposure to perfluoroalkyl acids and markers of kidney function among  
513 children and adolescents living near a chemical plant. *Environmental health perspectives* 121,  
514 625-630.

515 Wong, F., MacLeod, M., Mueller, J.F., Cousins, I.T., 2014. Enhanced elimination of perfluorooctane  
516 sulfonic acid by menstruating women: evidence from population-based pharmacokinetic  
517 modeling. *Environmental science & technology* 48, 8807-8814.

518 Ye, X., Kato, K., Wong, L.Y., Jia, T., Kalathil, A., Latremouille, J., Calafat, A.M., 2017. Per- and  
519 polyfluoroalkyl substances in sera from children 3 to 11 years of age participating in the



520 National Health and Nutrition Examination Survey 2013-2014. International journal of hygiene  
521 and environmental health.

522 Yildiz, B.O., Bozdog, G., Yapici, Z., Esinler, I., Yarali, H., 2012. Prevalence, phenotype and  
523 cardiometabolic risk of polycystic ovary syndrome under different diagnostic criteria. Human  
524 reproduction 27, 3067-3073.

525

526

## Tables and Figures

**Table 1.** Summary of PFAA serum concentrations in PCOS case-control study (ng/mL)

|        | Detection frequency, n (%) |         | Geometric Mean |      | Range     |           | Average ratio | LOR   |     |
|--------|----------------------------|---------|----------------|------|-----------|-----------|---------------|-------|-----|
|        | Serum                      | FF      | Serum          | FF   | Serum     | FF        | FF:Serum      | Serum | FF  |
| PFOS   | 59                         | 58      | 3.46           | 2.00 | 0.93-7.71 | 0.60-4.29 | 0.59          | 0.5   | 0.2 |
| PFOA   | 59                         | 58      | 2.39           | 1.82 | 0.5-8.16  | 0.43-6.64 | 0.78          | 0.1   | 0.1 |
| PFHxS  | 59                         | 58      | 1.04           | 0.88 | 0.2-10.2  | 0.1-9.07  | 0.86          | 0.05  | 0.1 |
| PFNA   | 59                         | 58      | 0.57           | 0.41 | 0.2-1.79  | 0.1-1.43  | 0.77          | 0.2   | 0.1 |
| PFDA   | 45 (76)                    | 14 (24) | 0.31           | -    | <LOR-     | <LOR-     | -             | 0.2   | 0.2 |
| PFPeA  | 29 (49)                    | 0       | -              | -    | <LOR-     | -         | -             | 0.5   | 0.3 |
| PFUnDA | 21 (36)                    | 0       | -              | -    | <LOR-     | -         | -             | 0.2   | 0.7 |
| PFHpA  | 10 (17)                    | 9 (15)  | -              | -    | <LOR-     | <LOR-     | -             | 0.1   | 0.1 |
| PFBS   | 4 (6.8)                    | 12 (21) | -              | -    | <LOR-     | <LOR-     | -             | 0.2   | 0.2 |
| PFBA   | 0                          | 0       | -              | -    | -         | -         | -             | 0.5   | 0.2 |
| PFHxA  | 0                          | 0       | -              | -    | -         | -         | -             | 0.5   | 0.1 |
| PFDS   | 0                          | 0       | -              | -    | -         | -         | -             | 0.5   | 0.1 |
| PFDoDA | 0                          | 0       | -              | -    | -         | -         | -             | 0.5   | 0.4 |

*FF, follicular fluid; LOR, limit of reporting; PFOS, perfluorooctane sulfonate; PFOA, perfluorooctanoic acid; PFHxS, perfluorohexane sulfonate; PFNA, perfluorononanoic acid; PFDA, perfluorodecanoic acid; PFPeA, perfluoropentanoic acid; PFUnDA, perfluoroundecanoic acid; PFHpA, perfluoroheptanoic acid; PFBS, perfluorobutanesulfonate; PFBA, perfluorobutanoic acid; PFHxA, perfluorohexanoic acid; PFDS, Perfluorodecane sulfonate; PFDoDA, perfluorododecanoic acid*

**Table 2.** Demographics, hormone and biochemistry endpoints, and serum and follicular PFAA concentrations for PCOS patients and controls.

|  | Control (n=28)<br>Mean $\pm$ SD | PCOS (n=31)<br>Mean $\pm$ SD |
|--|---------------------------------|------------------------------|
| Age (years)                              | 32.9 $\pm$ 4.6                  | 30.7 $\pm$ 4.6               |
| Body mass index (kg/m <sup>2</sup> )     | 25.6 $\pm$ 3.7                  | 25.9 $\pm$ 3.8               |
| Menarche (years)                         | 13.1 $\pm$ 1.8                  | 12.8 $\pm$ 1.3               |
| Irregular menstrual cycle (%)            | 14%                             | 87%***                       |
| Nulliparous (%)                          | 97%                             | 83%                          |
| Insulin ( $\mu$ U/ml)                    | 7.9 $\pm$ 4.1                   | 7.9 $\pm$ 4.6                |
| Fasting glucose (nmol/L)                 | 4.9 $\pm$ 0.4                   | 4.4 $\pm$ 0.8**              |
| HbA1C (mmol/mol)                         | 30.9 $\pm$ 6.5 (n=27)           | 31.8 $\pm$ 3.0 (n=28)        |
| HOMA-IR                                  | 1.8 $\pm$ 1.0                   | 1.9 $\pm$ 1.6                |
| Metformin use (%)                        | 0%                              | 47%***                       |
| SHBG (nmol/L)                            | 104.2 $\pm$ 80.3 (n=28)         | 71.7 $\pm$ 62.2 (n=28)       |
| Testosterone (nmol/L)                    | 0.85 $\pm$ 0.56 (n=27)          | 1.04 $\pm$ 0.37 (n=27)       |
| Free androgen index (FAI)                | 1.44 $\pm$ 1.47 (n=27)          | 3.32 $\pm$ 4.08 (n=27)*      |
| Estradiol (pmol/L)                       | 398.0 $\pm$ 423.4 (n=27)        | 259.1 $\pm$ 276.0 (n=26)     |
| Androstenedione (nmol/L)                 | 2.7 $\pm$ 1.4 (n=23)            | 4.0 $\pm$ 1.5 (n=24)**       |
| TSH (mU/L)                               | 2.3 $\pm$ 1.0 (n=27)            | 2.0 (0.8) (n=28)             |
| Free T3 (pmol/L)                         | 4.8 $\pm$ 0.7 (n=26)            | 4.8 $\pm$ 0.7 (n=24)         |
| Free T4 (pmol/L)                         | 11.2 $\pm$ 1.3 (n=26)           | 11.4 $\pm$ 2.2 (n=24)        |
| eGFR (mL/min 1.73m <sup>-2</sup> )       | 97.4 $\pm$ 18.9 (n=28)          | 88.3 $\pm$ 10.1 (n=28)*      |
| C Reactive Protein (mg L <sup>-1</sup> ) | 2.34 $\pm$ 2.34 (n=27)          | 2.77 $\pm$ 2.57 (n=28)       |
| Albumin (g/L)                            | 40.1 $\pm$ 2.9 (n=28)           | 40.6 $\pm$ 3.1 (n=28)        |
| Serum <sup>a</sup>                       |                                 |                              |
| PFOS, (ng/mL)                            | 3.1 (2.6-3.6)                   | 3.9 (3.4-4.4)*               |
| PFOA, (ng/mL)                            | 2.4 (1.9-2.9)                   | 2.4 (2.0-2.9)                |
| PFHxS, (ng/mL)                           | 0.9 (0.8-1.2)                   | 1.1 (0.9-1.4)                |
| PFNA, (ng/mL)                            | 0.5 (0.4-0.6)                   | 0.6 (0.5-0.7)                |
| Follicular fluid <sup>a</sup>            |                                 |                              |
| PFOS, (ng/mL)                            | 1.8 (1.6-2.1)                   | 2.2 (1.9-2.5)                |

---

|                |               |               |
|----------------|---------------|---------------|
| PFOA, (ng/mL)  | 1.9 (1.6-2.3) | 1.7 (1.4-2.1) |
| PFHxS, (ng/mL) | 0.8 (0.6-1.0) | 0.9 (0.7-1.2) |
| PFNA, (ng/mL)  | 0.4 (0.3-0.5) | 0.4 (0.3-0.5) |

---


\*p<0.05, \*\*p<0.01, \*\*\*p<0.001; <sup>a</sup> Geometric mean (95% CI)




---

|                             |                    |                    |              |              |                |
|-----------------------------|--------------------|--------------------|--------------|--------------|----------------|
| ln(PFNA, ng/ml)             | 0.29 (0.17)        | 0.38 (0.24)        | -0.09 (0.34) | 0.97 (0.57)  | 85.15 (157.92) |
|                             | <i>0.093</i>       | <i>0.124</i>       | <i>0.795</i> | <i>0.094</i> | <i>0.592</i>   |
| ln( $\sum$ PFAA,<br>umol/L) | <b>0.41 (0.18)</b> | <b>0.61 (0.28)</b> | -0.21 (0.36) | 1.11 (0.63)  | 88.04 (163.56) |
|                             | <b>0.024</b>       | <b>0.035</b>       | <i>0.572</i> | <i>0.084</i> | <i>0.593</i>   |

---




**Figure 1.** Plot of serum concentration (ng/mL, n=59) of PFAAs with detection frequencies >50%. Box indicates interquartile range, whiskers indicate 5<sup>th</sup> and 95<sup>th</sup> %, horizontal line indicates median. Significantly different geometric means (p<0.05) denoted by asterisk.



**Figure 2.** Correlation of PFOS (A), PFOA (B), PFHxS (C) and PFNA (D) in serum and follicular fluid. Shaded area represents 95% confidence interval of regression line forced through the origin (red);  $R^2$  of weighted linear regression shown in bottom right corner.





**Figure 3:** Perfluorooctane sulfonate (PFOS) concentration in women with regular versus irregular menstrual cycles ( $p < 0.05$  for difference of means). Horizontal line represents the geometric mean.